

LOGARITHMIC STABLE MAPS TO DELIGNE–FALTINGS PAIRS I

QILE CHEN

CONTENTS

1. Introduction	1
2. Algebraicity of the stack of log maps	4
3. Minimal logarithmic maps to rank one Deligne-Faltings log pairs	16
4. The stack of minimal log maps as category fibered over \mathbf{LogSch}^{fs}	27
5. The boundedness theorem for minimal log stable maps	31
6. The weak valuative criterion for minimal log stable maps	36
Appendix A. Prerequisites on logarithmic geometry	45
Appendix B. Logarithmic curves and their stacks	49
References	52

1. INTRODUCTION

1.1. Background on relative Gromov-Witten theory. Gromov-Witten theory relative to a smooth divisor was introduced during the past decade, for the purpose of proving a degeneration formula, in the symplectic setting by A.M. Li and Y. Ruan [LR01], and about the same time by E.N. Ionel and T. Parker [IP03]. On the algebraic side, this was worked out by Jun Li [Li01, Li02]. This approach uses the idea of expanded degenerations, which was introduced by Ziv Ran [Ran87]. A related idea of admissible covers was introduced even earlier by Harris and Mumford [HM82].

Recently, the idea of expanded degeneration was systematically studied using orbifold techniques by D. Abramovich and B. Fantechi [AF], and an elegant proof of degeneration formula was given there.

On the other hand, the idea of admissible covers was revisited by Mochizuki [Moc95] using logarithmic geometry. Along the similar idea, B. Kim defined the logarithmic stable maps [Kim09], by putting certain log structures on Jun Li’s predeformable maps. Then he proved that the stack parameterizing such maps is a proper DM stack, and has an explicit perfect obstruction theory using the work of M. Olsson [Ols03a, Ols05]. A degeneration formula under Kim’s setting is proved in [Che].

Another approach using logarithmic structures without expansions was first proposed by Bernd Siebert in 2001 [Sie01]. The goal here is also to obtain the degeneration formula, but in a more general situation, such as simple normal crossings divisors. However, the program has been on hold for a while, since Mark Gross and Bernd Siebert were working on other projects in mirror symmetry. Only recently they have taken up the unfinished project of Siebert jointly. In particular, they succeeded in finding a definition of basic log maps, a

Date: February 8, 2011.

Research partially supported by funds from NSF award DMS-0901278.

crucial ingredient for a good moduli theory of stable log maps to a fixed target with Zariski log structures [GS]. Their definition builds on insights from tropical geometry, obtained by probing the stack of log maps using the standard log point, and is compatible with the minimality introduced in this paper.

A different approach using exploded manifolds to studying holomorphic curves was recently introduced by Brett Parker in [Par09a], [Par09b], and [Par09c]. It also aimed at defining and computing relative and degenerated Gromov-Witten theories in general situation. The theory of exploded manifolds uses a generalized version of tropical curves, and is closely related to logarithmic geometries — the explosion functors, which is central to this theory, can be phrased in terms of certain kind of base change in log geometry, see [Par09a, Section 5]. It was pointed out by Mark Gross that this approach is parallel, and possibly equivalent to the logarithmic approach.

1.2. The approach and main results of this paper. The goal of this paper is to develop the relative Gromov-Witten theory along the logarithmic approach proposed by Bernd Siebert. However, we use somewhat different methods. Instead of using tropical geometry, and probing the stack using standard log point, we associated to each log map a marked graph as in Section 3.3, which allows us to define the right base log structure. We now describe our methods as follows.

The target we will consider in this paper is a projective variety X equipped with a rank-1 Deligne-Faltings log structure \mathcal{M}_X on X as in Definition 3.1.1, which comes from a line bundle L on X , with a morphism of sheaves $s : L \rightarrow \mathcal{O}_X$. In particular, if L is the ideal sheaf of a smooth divisor $D \subset X$, and s is the natural inclusion, then this will recover the relative case. See Section A.2 for more details on DF log structures. Denote by $X^{\log} = (X, \mathcal{M}_X)$ the corresponding log scheme. Instead of considering usual stable maps to the expansions of X , we investigate the usual stable maps to X , with fixed log structure \mathcal{M}_X on the target, and suitable log structures on the source curves.

In the subsequent paper [AC], we will consider target X with generalized Deligne-Faltings log structure \mathcal{M}_X , namely where there exists a global map $P \rightarrow \overline{\mathcal{M}}_X$ from a toric monoid P , which locally lifts to a chart. In particular, this covers many interesting cases, such as a variety with a simple normal crossings divisor, or a simple normal crossings degeneration of a variety with simple normal crossings singularities. It does not cover the case of a normal crossings divisor which is not simple. We hope one can also cover this using the descent argument along this approach.

A key point of this paper is the observation made in Section 3.2, which describes the log map on the level of characteristic monoids. This leads us to the notions of marked graphs 3.3.2 and minimality 3.5.1. Such minimality condition can be explained as the “minimal requirements” which a log map needs to satisfy. Then our minimal log stable maps are defined to be usual stable maps with the minimal log structures. Denote by $\mathcal{K}_\Gamma(X^{\log})$ the category fibered over the category of schemes, which for any scheme T associates the groupoid of minimal log stable maps over T with numerical data Γ . We refer to Section 3.6 for the precise meaning of $\mathcal{K}_\Gamma(X^{\log})$. The main result of this paper is the following:

Theorem 1.2.1. *The fibered category $\mathcal{K}_\Gamma(X^{\log})$ is a proper Deligne-Mumford stack. Furthermore, the natural map $\mathcal{K}_\Gamma(X^{\log}) \rightarrow \mathcal{K}_{g,n}(X, \beta)$ by removing the log structures from minimal log stable maps is representable and finite.*

Remark 1.2.2. In fact, the stack $\mathcal{K}_\Gamma(X^{log})$ carries a universal minimal log structure, which will be denoted by $\mathcal{M}_{\mathcal{K}_\Gamma(X^{log})}$. Thus the pair $(\mathcal{K}_\Gamma(X^{log}), \mathcal{M}_{\mathcal{K}_\Gamma(X^{log})})$ can be viewed as a log stack in the sense of Olsson, see Section A.3. By applying the standard technique in [BF97], and replacing the usual cotangent complex by logarithmic cotangent complex [Ols05], one can produce a perfect obstruction theory of $\mathcal{K}_\Gamma(X^{log})$ relative to $\mathfrak{M}_{g,n}^{pre}$, the stack of log prestable curves defined in Section B.3. We will discuss this perfect obstruction theory and the corresponding virtual fundamental class in another paper.

Up to now, we only introduce $\mathcal{K}_\Gamma(X^{log})$ as category fibered over \mathfrak{Sch} , the category of schemes. Denote by \mathfrak{LogSch}^{fs} the category of fine and saturated log schemes as introduced in Section A.1. The following result exhibits another important aspect of our construction:

Theorem 1.2.3. *The pair $(\mathcal{K}_\Gamma(X^{log}), \mathcal{M}_{\mathcal{K}_\Gamma(X^{log})})$ defines a category fibered over \mathfrak{LogSch}^{fs} , which for any fs log scheme (S, \mathcal{M}_S) associates the category of log stable maps over (S, \mathcal{M}_S) .*

The above categorical interpretation allows us to work systematically with fs log schemes rather than usual schemes. This point of view will be a useful tool in [AC], where we reduce the case with generalized DF-log structure on the target to the case of this paper.

1.3. Outline of the paper. In Section 2, we fix a morphism of fs log schemes $X^{log} \rightarrow B^{log}$, define an auxiliary category $\mathcal{LM}_{g,n}(X^{log}/B^{log})$ of all log maps with target X, fibered over schemes, and show that it is an algebraic stack in the sense of Artin. This stack is unbounded and serves mainly as a construction technique. This will be achieved by verifying Artin’s criteria [Art74, 5.1]. Here the deformation theory of our log maps will be given by the log cotangent complex developed in [Ols05].

Section 3 is aimed at the construction of minimal log maps. In fact, for each log map over a geometric point with fs log structure, we can associate a marked graph, see Construction 3.4.1. These graphs will be used to describe the minimality condition. Then we show that minimality is an open condition, see Proposition 3.5.2. This implies the algebraicity of the stack of minimal log maps using the results of Section 2. In Section 3.7 we show that there are at most finitely many minimal log stable maps over a fixed underlying stable map with a fixed marked graph. The finiteness of automorphisms of minimal log stable maps over a geometric point is proved in Proposition 3.8.1.

Section 4 is devoted to proving Theorem 1.2.3. This will follow naturally from the universal property of minimal log maps in Proposition 4.1.1.

In Section 5, we will show that $\mathcal{K}_\Gamma(X^{log})$ is of finite type by stratifying the stack, and bounding the stratum associated to each marked graph. Indeed, we will prove the boundedness of $\mathcal{K}_\Gamma(X^{log})$ relative to the stack of usual stable maps. One issue here is in constructing all maps of log structures for a given graph. We will turn this into constructing isomorphisms of corresponding line bundles.

The weak valuative criterion of $\mathcal{K}_\Gamma(X^{log})$ is proved in Section 6. In fact, the universal property of minimality produces an extension of *minimal* log stable map, once we can find *any* extension of log stable maps, not necessarily minimal. For separateness, we first show that the marked graph is uniquely determined by the generic fiber. Then we introduce a new map $\bar{\beta}$ as in (6.4.3), which helps us compare any two possible extensions, see Lemma 6.5.1. This leads us to the uniqueness of the extension. In the end, we give a proof of Theorem 1.2.1, and show that the stack of minimal log stable maps is representable and finite over the stack of usual stable maps.

Finally, we have two appendices collecting various results of logarithmic geometry and logarithmic curves, as we need in this paper. The notion of log pre-stable curve is introduced in Definition B.2.3.

1.4. Conventions. Throughout this paper, we will work over an algebraically closed field of characteristic 0, denoted by \mathbb{C} .

The word locally always means étale locally, and neighborhood always means étale neighborhood, unless otherwise specified.

Given a scheme or algebraic stack X , a point $p \in X$ means a closed point unless otherwise specified. We denote by \bar{p} an algebraic closure of p .

We usually use X^{log} to denote a log scheme (X, \mathcal{M}_X) if no confusion could arise. The map $\exp : \mathcal{M}_X \rightarrow \mathcal{O}_X$ is reserved for the structure map of \mathcal{M}_X . Given a section $u \in \mathcal{O}_X$, we sometimes use $\log u$ to denote the corresponding section in \mathcal{M}_X if no confusion could arise.

The category of schemes, fine log schemes, and fs log schemes are denoted by \mathfrak{Sch} , \mathfrak{LogSch} , and \mathfrak{LogSch}^{fs} respectively. See Section A.1 for the precise definitions.

The letter ξ (respectively ξ^{log}) is reserved for log maps over a usual scheme (respectively log scheme). Given a log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over S as in Remark 3.1.5, we will denote by \mathcal{M}_C the corresponding log structure on C , if no confusion could arise.

1.5. Acknowledgements. I am very grateful to my advisor Dan Abramovich, who has suggested several interesting problems to me, including the main problem of this paper. He has given me continuous support and encouragement, and his suggestions have greatly influenced the shape of this paper. I am grateful to Mark Gross, Bernd Siebert, and Martin Olsson who pointed out several mistakes in the previous draft of this paper, and also gave many nice suggestions and comments. I would also like to thank Davesh Maulik, William Gillam, and Jonathan Wise for their helpful conversations.

2. ALGEBRICITY OF THE STACK OF LOG MAPS

In this section, we prove that the stack $\mathcal{LM}_{g,n}(X^{log}/B^{log})$ parameterizing log maps as in Definition 2.1.5, is algebraic by checking Artin's criteria [Art74, 5.1]. The result in this section is only used to prove the stack of minimal log stable maps $\mathcal{K}_{n,g}^{mst}(X^{log}, \beta)$ as in Definition 3.6.5 is algebraic. The reader may wish to assume the result of Theorem 2.1.10, and skip to the next section.

2.1. Log maps over \mathfrak{LogSch} and over \mathfrak{Sch} .

Conventions 2.1.1. In this section, we fix a separated, finite type, log flat morphism of log schemes $\pi : X^{log} \rightarrow B^{log}$. Denote by B and X the underlying schemes of B^{log} and X^{log} respectively. Let \mathcal{M}_B and \mathcal{M}_X be the log structure on B^{log} and X^{log} respectively. Given any B -scheme S , denote by $(X_S, \mathcal{M}_{X_S}^{X_S/S}) \rightarrow (S, \mathcal{M}_S^{X_S/S})$ the pull-back of $X^{log} \rightarrow B^{log}$ over S .

As an analogue of usual pre-stable maps, we introduce our log maps over log schemes as follows:

Definition 2.1.2. A log map ξ^{log} over a fine B^{log} -log scheme (S, \mathcal{M}_S) is a commutative diagram of log schemes:

$$(2.1.1) \quad \begin{array}{ccc} (C, \mathcal{M}_C) & \xrightarrow{f} & (X_S, \mathcal{M}_{X_S}) \\ & \searrow & \swarrow \\ & (S, \mathcal{M}_S) & \end{array}$$

where

- (1) The arrow $(C, \mathcal{M}_C) \rightarrow (S, \mathcal{M}_S)$ is given by the log curve $(C \rightarrow S, \mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S)$ as in Definition B.2.2.
- (2) The arrow $(X_S, \mathcal{M}_{X_S}) \rightarrow (S, \mathcal{M}_S)$ is obtained from the following cartesian diagram of log schemes:

$$\begin{array}{ccc} (X_S, \mathcal{M}_X) & \longrightarrow & X^{log} \\ \downarrow & & \downarrow \pi \\ (S, \mathcal{M}_S) & \longrightarrow & B^{log}. \end{array}$$

Given a log map of fine log schemes $g : (T, \mathcal{M}_T) \rightarrow (S, \mathcal{M}_S)$, we define the pull-back $g^*\xi^{log}$ to be the log map ξ_T^{log} over (T, \mathcal{M}_T) obtained by pulling back (2.1.1) via the log map g .

The above definition gives a category of log maps fibered over $\mathbf{LogSch}_{B^{log}}$, the category of fine log schemes over B^{log} . This category allows pull-back via arbitrary log maps, hence changes the base log structures. In another word, it only parametrizes the “log maps”, without the information of log structures on the base. This is the category of most interest to us.

However, algebraic stacks are built over the category of schemes, rather than the category of log schemes. In order to have the algebraic structure, we need to introduce another fibered category over \mathbf{Sch}_B , the category of B -schemes. This leads to the following definition:

Definition 2.1.3. A log map ξ over a B -scheme S consists of a fine log scheme (S, \mathcal{M}_S) over B^{log} , and a log map ξ^{log} over (S, \mathcal{M}_S) as in Definition 2.1.2. Usually we denote it by

$$\xi = (C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S^{X_S/S} \rightarrow \mathcal{M}_S, \mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S, f),$$

where $\mathcal{M}_S^{X_S/S}$ is the pull-back of \mathcal{M}_B via the structure map $S \rightarrow B$.

Consider another B -scheme T , and a B -scheme morphism $T \rightarrow S$. Then we have an induced strict arrow $g : (T, \mathcal{M}_T) \rightarrow (S, \mathcal{M}_S)$, where $\mathcal{M}_T := g^*(\mathcal{M}_S)$. The pull-back ξ_T of ξ via $T \rightarrow S$ is given by the log scheme (T, \mathcal{M}_T) , and the log map $\xi_T^{log} = g^*\xi^{log}$ over (T, \mathcal{M}_T) . In the rest of this section, if no confusion could arise, we will use $(C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S, f)$ to denote the log map ξ over S .

Since isomorphisms are central to the structure of stacks, we spell out the resulting notion of an isomorphism of log maps over schemes:

Definition 2.1.4. Consider two log maps $\xi_1 = (C_1 \rightarrow S, X_S \rightarrow S, \mathcal{M}_1, f_1)$ and $\xi_2 = (C_2 \rightarrow S, X_S \rightarrow S, \mathcal{M}_2, f_2)$ over a scheme S . An isomorphism $\xi_1 \rightarrow \xi_2$ over S is given by a triple

(ρ, θ, γ) fitting in the following commutative diagram of log schemes:

$$(2.1.2) \quad \begin{array}{ccc} (C_1, \mathcal{M}_{C,1}) & \longrightarrow & (X_S, \mathcal{M}_{X,1}) \\ \downarrow \rho & \searrow & \swarrow & \downarrow \gamma \\ & (S, \mathcal{M}_1) & \\ & \downarrow \theta & \\ (C, \mathcal{M}_{C,2}) & \longrightarrow & (X_S, \mathcal{M}_{X,2}) \\ & \searrow & \swarrow \\ & (S, \mathcal{M}_2) & \end{array}$$

where

- (1) The pair (ρ, θ) is an arrow of log curves $(C_1 \rightarrow S, \mathcal{M}_1) \rightarrow (C_2 \rightarrow S, \mathcal{M}_2)$ as in Definition B.3.1.
- (2) The arrow θ is an isomorphism of log schemes over B^{log} fitting in the following commutative diagram:

$$\begin{array}{ccc} (S, \mathcal{M}_1) & \xrightarrow{\theta} & (S, \mathcal{M}_2) \\ & \searrow & \swarrow \\ & B^{log} & \end{array}$$

- (3) The arrow γ is obtained from the following cartesian diagram of log schemes:

$$\begin{array}{ccc} (X_S, \mathcal{M}_{X_S,1}) & \xrightarrow{\gamma} & (X_S, \mathcal{M}_{X_S,2}) \\ \downarrow & & \downarrow \\ (S, \mathcal{M}_1) & \xrightarrow{\theta} & (S, \mathcal{M}_2). \end{array}$$

Note that the underlying map $\underline{\theta}$ and $\underline{\gamma}$ are both identities of the corresponding underlying schemes.

Denote by $\mathcal{I}som_S(\xi_1, \xi_2)$ the functor over S , which for any S -scheme $T \rightarrow S$ associates the set of isomorphisms of $\xi_{T,1}$ and $\xi_{T,2}$ over T , where $\xi_{T,1}$ and $\xi_{T,2}$ are the pull-back of ξ_1 and ξ_2 via $T \rightarrow S$ respectively. Denote by $\mathcal{A}ut_S(\xi)$ the functor of automorphisms of ξ over S .

Definition 2.1.5. Denote by $\mathcal{LM}_{g,n}(X^{log}/B^{log})$ the fibered category over \mathfrak{Sch}_B , which for any $S \rightarrow B$ associates the groupoid of log maps ξ over S , with the underlying pre-stable curves of genus g , and n -markings. For simplicity of notation, in this section we will use \mathcal{LM} to denote $\mathcal{LM}_{g,n}(X^{log}/B^{log})$.

Remark 2.1.6. By Definition 2.1.3, we only allow pull-backs of log maps via strict log maps in \mathcal{LM} , hence do not change the log structures. Thus, given a scheme S , the groupoid $\mathcal{LM}(S)$ contains all possible log structures \mathcal{M}_S on S with log maps over (S, \mathcal{M}_S) . This is a huge stack, as it parametrizes in particular all possible log structures on the base. One would like to consider a smaller stack parameterizing only log maps without the information of the base log structures. It will be shown in Section 4 that if we work over fs log schemes

rather than the usual category of schemes for the base, then the stack we want is $\mathcal{K}_{g,n}^{min}(X^{log})$ as introduced in Section 3.

Denote by $\mathfrak{M}_{g,n}$ the algebraic stack of genus g , n -marked pre-stable curves with the canonical log structure as in Section B.1. Consider the new algebraic stack

$$\mathfrak{B} = \mathcal{L}og_{\mathfrak{M}_{g,n} \times B^{log}},$$

where the fibered product is in the log sense, and $\mathcal{L}og_{\bullet}$ is the log stack introduced in Section A.3. By Theorem A.3.2, the stack \mathfrak{B} is algebraic over B .

Remark 2.1.7. We give the moduli interpretation of \mathfrak{B} as follows. For any B -scheme S , an object $\zeta \in \mathfrak{B}(S)$ is a diagram

$$(2.1.3) \quad \begin{array}{ccc} (C, \mathcal{M}_C) & & (X_S, \mathcal{M}_{X_S}) \\ & \searrow & \swarrow \\ & (S, \mathcal{M}) & \end{array}$$

where the left arrow is a family of genus g , n -marked log curves given by the induced map $(S, \mathcal{M}_S) \rightarrow \mathfrak{M}_{g,n}$, and the right arrow is given by the induced map $(S, \mathcal{M}_S) \rightarrow B^{log}$. Consider two objects ζ_1 and ζ_2 in $\mathfrak{B}(S)$. An arrow $\zeta_1 \rightarrow \zeta_2$ over the scheme S is a triple (ρ, θ, γ) given by the following diagram

$$(2.1.4) \quad \begin{array}{ccccc} (C_1, \mathcal{M}_{C,1}) & & & & (X_S, \mathcal{M}_{X_S,1}) \\ & \searrow & & & \swarrow \\ & & (S, \mathcal{M}_1) & & \\ \rho \downarrow & & \downarrow \theta & & \downarrow \gamma \\ (C, \mathcal{M}_{C,2}) & & & & (X_S, \mathcal{M}_{X_S,2}) \\ & \searrow & & & \swarrow \\ & & (S, \mathcal{M}_2) & & \end{array}$$

where the square on the left is an isomorphism of log curves, and the square on the right satisfies the condition in Definition 2.1.4(2) and (3).

Observation 2.1.8. Note that there is a natural morphism of fibered categories $\mathcal{LM} \rightarrow \mathfrak{B}$ by removing the log map f as in Definition 2.1.3. Note that any non-trivial isomorphism of a log map is a non-trivial isomorphism of the corresponding log source and target. This implies that \mathcal{LM} is a pre-sheaf over \mathfrak{B} .

We denote by $\mathcal{K}_{g,n}^{pre}(X/B)$ the stack of usual pre-stable maps with the source given by genus g , n -marked pre-stable curves, and family of targets given by $X \rightarrow B$. This is an algebraic stack over B . For simplicity of notations, we will denote this stack by \mathcal{K} .

Observation 2.1.9. Note that we have a natural arrow $\mathcal{LM} \rightarrow \mathcal{K}$ by removing all log structures. Given a log map ξ , denote by $\underline{\xi}$ the corresponding object in \mathcal{K} .

Our main result of this section is the following:

Theorem 2.1.10. *The fibered category \mathcal{LM} is an algebraic stack.*

Proof. The rest of this section is devote to the proof of this theorem. The representability of the diagonal $\mathcal{LM} \rightarrow \mathcal{LM} \times \mathcal{LM}$ is proved in Section 2.2. By Observation 2.1.8, we have a natural map $\mathcal{LM} \rightarrow \mathfrak{B}$ to the algebraic stack \mathfrak{B} . Thus, to produce a smooth cover for \mathcal{LM} , it is enough to check Artin's criteria [Art74, 5.1] relative to \mathfrak{B} . This will be done from Section 2.3 to 2.7. \square

2.2. Representability of the isomorphism functors of log maps.

Proposition 2.2.1. *Consider two log maps ξ_1 and ξ_2 over a B -scheme S as in Definition 2.1.4. The functor $\mathcal{I}som_S(\xi_1, \xi_2)$ is represented by an algebraic space of finite type over S .*

Proof. Using the notations as in Definition 2.1.4, Remark 2.1.7, and Observation 2.1.9, we form the following commutative diagram:

$$(2.2.1) \quad \begin{array}{ccccc} \mathcal{I}som_S(\xi_1, \xi_2) & & & & \\ & \searrow^{\phi_2} & & & \\ & & I & \xrightarrow{\quad} & \mathcal{I}som_S(\underline{\xi}_1, \underline{\xi}_2) \\ & \searrow^{\phi_3} & \downarrow & & \downarrow^{\psi_2} \\ & & \mathcal{I}som_S(\zeta_1, \zeta_2) & \xrightarrow{\psi_1} & \mathcal{I}som_S(\underline{\zeta}_1, \underline{\zeta}_2), \\ & \searrow_{\phi_1} & & & \end{array}$$

where the square is cartesian, and ϕ_3 is given by the universal property of fiber product. Here ζ_i is the corresponding log source and target of ξ_i given by the natural map in Observation 2.1.8, and $\underline{\xi}_i$ is the underline map of ξ_i given by the natural map in Observation 2.1.9. The object $\underline{\zeta}_i$ can be obtained by removing log structures on ζ_i , or given by the source and target of ξ_i .

Note that any isomorphism of ξ_1 and ξ_2 induces trivial isomorphism of the underlying structure of the target $X_S \rightarrow S$. Thus, the sheaf $\mathcal{I}som_S(\underline{\zeta}_1, \underline{\zeta}_2)$ is the sheaf of isomorphisms of the underlying curves. Since $\mathcal{I}som_S(\xi_1, \xi_2)$, $\mathcal{I}som_S(\zeta_1, \zeta_2)$, and $\mathcal{I}som_S(\underline{\zeta}_1, \underline{\zeta}_2)$ are represented by algebraic spaces of finite type over S , it is enough to show that ϕ_3 is representable and of finite type.

Consider an S -scheme U , and an arrow $U \rightarrow I$ given by a pair (τ, λ) , where

$$\tau \in \mathcal{I}som_S(\zeta_1, \zeta_2)(U) \quad \text{and} \quad \lambda \in \mathcal{I}som_S(\underline{\xi}_1, \underline{\xi}_2)(U),$$

such that their induced elements in $\mathcal{I}som_S(\underline{\zeta}_1, \underline{\zeta}_2)(U)$ coincide. Now we have a cartesian diagram :

$$\begin{array}{ccc} I' & \longrightarrow & \mathcal{I}som_S(\xi_1, \xi_2) \\ \downarrow & & \downarrow \\ U & \xrightarrow{(\tau, \lambda)} & I. \end{array}$$

Here I' is the sheaf over U which for any $V \rightarrow U$ associated a unital set $\{*\}$ if $(\tau, \lambda)_V$ induces an isomorphism between $\xi_{1,V}$ and $\xi_{2,V}$, and the empty set otherwise. Next we will show that $I' \rightarrow U$ is a locally closed immersion of finite type.

For simplicity, we assume $U = S$, denote by $\tau = (\rho, \theta, \gamma)$ as in Remark 2.1.7. We need to show that the commutativity of the following diagram of log schemes is represented by a

locally closed immersion of finite type:

$$\begin{array}{ccc} (C_1, \mathcal{M}_{C_1}) & \xrightarrow{f_1} & (X, \mathcal{M}_{X,1}) \\ \rho \downarrow & & \gamma \downarrow \\ (C_2, \mathcal{M}_{C_2}) & \xrightarrow{f_2} & (X, \mathcal{M}_{X,2}). \end{array}$$

Since the map τ already gives an isomorphism of the underlying structure, we only need to consider the commutativity of

$$(2.2.2) \quad \begin{array}{ccc} \mathcal{M}_{C_1} & \xleftarrow{f_1^b} & f_1^* \mathcal{M}_{X,1} \\ \rho^b \uparrow & & \uparrow \gamma^b \\ \rho^* \mathcal{M}_{C_2} & \xleftarrow{\rho^* \circ f_2^b} & \rho^* \circ f_2^* \mathcal{M}_{X,2}. \end{array}$$

Our statement follows from the following lemma. □

Lemma 2.2.2. *The condition that (2.2.2) commutes, is represented by a quasi-compact locally closed immersion $Z \rightarrow S$.*

Proof. The commutativity of (2.2.2) is equivalent to the equality

$$(2.2.3) \quad \rho^b \circ (\rho^* \circ f_2^b) = f_1^b \circ \gamma^b.$$

By [Ols03a, 3.6], the condition that (2.2.3) holds on the level of characteristic is an open condition on C_1 . Since $C_1 \rightarrow S$ is flat and proper, by shrinking S , we can assume that the equality (2.2.3) on the level of characteristic holds.

With this assumption on the characteristic, the proof in [Ols03a, 3.6] shows that the (2.2.3) is represented by a closed subscheme $T \subset C_1$ on the fiber. Note that the statement is locally on S . Further shrinking S , we can assume that the family $C \rightarrow S$ is projective. Now what we want is the maximal closed subscheme $Z \subset S$ parameterizing fibers $C_1 \times_S Z \subset T$ as in [Abr94, Definition 3, 4]. Then the lemma can be deduced from the “essential free” case of [SGA64, VIII Theorem 6.5]. See [Abr94, Theorem 6(3)] for the reduction argument. □

Next, we check the Artin’s criteria [Art74, 5.1].

2.3. \mathcal{LM} is a stack in the étale topology. By [Art74, 1.1], or [LMB00, Definition 3.1], we need to prove the following:

- (1) the isomorphism functor is a sheaf in the étale topology;
- (2) any étale descent datum for an object of \mathcal{LM} is effective.

Since the isomorphism functor is shown to be representable, it is a sheaf in the étale topology. For the second condition, let $\{S_i \rightarrow S\}_i$ be an étale covering of S , and $\xi_i \in \mathcal{LM}(S_i)$ for each i . Assume that we have isomorphisms $\phi_{ij} : \xi_i|_{S_i \times_S S_j} \rightarrow \xi_j|_{S_i \times_S S_j}$ for each pair (i, j) , which satisfy the cocycle condition.

For any i , let ζ_i be the corresponding log curve and target as in Remark 2.1.7 for ξ_i . Since such ζ_i is parametrized by the algebraic stack \mathfrak{B} , we can glue them together to obtain ζ over S , whose restriction to each S_i is ζ_i . Then étale locally we have a log map from ζ given by ξ_i . Since log maps are defined in terms of homomorphisms of étale sheaves, they can be glued from étale local data. Therefore we can glue ξ_i to obtain the log map ξ with the source curve given by ζ .

2.4. \mathcal{LM} is limit preserving. Consider

$$R = \varinjlim R_i,$$

where $\{R_i\}$ is a filtering inductive system of noetherian rings. Denote by $S = \text{Spec } R$ and $S_i = \text{Spec } R_i$. By [Art74, Section 1], we need to show that the following map of groupoids is an equivalence of categories:

$$\varinjlim \mathcal{LM}(S_i) \rightarrow \mathcal{LM}(S)$$

Consider a log map $\xi = (C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S, f)$ in $\mathcal{LM}(S)$. Since the stack \mathfrak{B} is locally of finite type, we have the family $\zeta = (C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S)$ coming from $\zeta_i = (C_i \rightarrow S_i, X_{S_i} \rightarrow S_i, \mathcal{M}_{S_i})$ over S_i for some i . Also notice that we have an induced map $S \rightarrow \mathcal{K}$ given by the underlying map. Since \mathcal{K} is locally of finite type, the underlying map \underline{f} comes from $\underline{f}_{i'}$ over some $S_{i'}$. We choose an index i_0 such that $i_0 > i$ and $i_0 > i'$.

It remains to consider the map of log structures $f^b : f^* \mathcal{M}_X \rightarrow \mathcal{M}_C$. We first introduce two stacks \mathcal{L}^Δ and \mathcal{L}^Λ as in [Ols05, section 2].

Remark 2.4.1. Consider a scheme U over \mathbb{Z} . Objects in $\mathcal{L}^\Delta(U)$ are commutative diagrams of log structures on U of the following form

(2.4.1)

$$\begin{array}{ccc} & \mathcal{M}_1 & \\ \swarrow & & \searrow \\ \mathcal{M}_2 & \xrightarrow{\quad} & \mathcal{M}_3. \end{array}$$

Objects in \mathcal{L}^Λ are diagrams of log structures on U of the following form

(2.4.2)

$$\begin{array}{ccc} & \mathcal{M}_1 & \\ \swarrow & & \searrow \\ \mathcal{M}_2 & & \mathcal{M}_3. \end{array}$$

It was shown in [Ols05, 2.4] that those two stacks \mathcal{L}^Δ and \mathcal{L}^Λ are algebraic stacks locally of finite type. Note that there is a natural morphism $\mathcal{L}^\Delta \rightarrow \mathcal{L}^\Lambda$ by dropping the bottom arrow in (2.4.1) to obtain (2.4.2).

Observation 2.4.2. Consider $\zeta = (\pi_C : C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S)$ the family of log sources and targets constructed above. There is a natural diagram of log structures on C as follows

(2.4.3)

$$\begin{array}{ccc} & \pi_C^* \mathcal{M}_S & \\ \swarrow & & \searrow \\ f^* \mathcal{M}_X & & \mathcal{M}_C. \end{array}$$

This induces a natural map $C \rightarrow \mathcal{L}^\Lambda$. Consider the fiber product $\mathcal{L}^\Delta \times_{\mathcal{L}^\Lambda} C$. This gives an algebraic stack parameterizing the bottom arrows f^b that fits in the above commutative diagram.

The map f^b is equivalent to a map $C \rightarrow \mathcal{L}^\Delta \times_{\mathcal{L}^\Lambda} C$. Note that the algebraic stack $\mathcal{L}^\Delta \times_{\mathcal{L}^\Lambda} C$ is locally of finite presentation. By [LMB00, Proposition 4.18(i)], we have the map f^b coming

from some $f_{i_1}^b$ over S_{i_1} for some $i_1 > i_0$. This map is compatible with all the log structures coming from base and target. Indeed, consider the composition

$$p_j : C_j \rightarrow \mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_j \rightarrow C_j.$$

Applying [LMB00, Proposition 4.18(i)] again, we see that the identity $p = id_C : C \rightarrow C$ comes from p_j for some $i_2 > i_1$. Thus, the map f_{i_2} also compatible with the underlying map f . This proves the essential surjectivity.

The full faithfulness follows from [LMB00, Proposition 4.15(i)] and the fact that the diagonal $\mathcal{LM} \rightarrow \mathcal{LM} \times_B \mathcal{LM}$ is representable and locally of finite type.

2.5. Deformations and obstructions. By [Art74, Definition 5.1], it remains to find a smooth cover of \mathcal{LM} . As in Observation 2.1.8, we have a representable map of stack $\mathcal{LM} \rightarrow \mathfrak{B}$. Since \mathfrak{B} is an algebraic stack, it would be enough to produce a smooth cover for $\mathcal{LM}_U := \mathcal{LM} \times_{\mathfrak{B}} U$, where $U \rightarrow \mathfrak{B}$ is an arbitrary smooth map. This can be done by checking Artin's criteria [Art74, 5.2] for \mathcal{LM}_U relative to U . First we consider the deformations and obstructions.

Let A_0 be a reduced noetherian ring over U , and $A' \rightarrow A \rightarrow A_0$ be an infinitesimal extension of A_0 , where $A' \rightarrow A$ is surjective whose kernel I is a finite A_0 -module, hence is a square-zero ideal. Denote by $S = \text{Spec } A$ and $S' = \text{Spec } A'$. Consider a log map $\xi_A = (C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S, f) \in \mathcal{LM}_U$. Let $\xi_0 = (C_0 \rightarrow S_0, X_{S_0} \rightarrow S_0, \mathcal{M}_{S_0}, f_0)$ be the restriction of ξ_A over A_0 . Since we are over U , the log sources and targets $(C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S)$ comes from the structure morphism $S \rightarrow U$. Note that we have another family of log sources and targets $(C' \rightarrow S', X_{S'} \rightarrow S', \mathcal{M}_{S'})$, which are also from the structure map $S' \rightarrow U$. To obtain a deformation of ξ_A over S' , it is equivalent to producing a dotted arrow f' that fits in the following commutative diagram:

$$(2.5.1) \quad \begin{array}{ccccc} (C, \mathcal{M}_C) & \xrightarrow{k} & (C', \mathcal{M}_{C'}) & & \\ & \searrow f & & \swarrow f' & \\ & & (X_S, \mathcal{M}_{X_S}) & \xrightarrow{j} & (X_{S'}, \mathcal{M}_{X_{S'}}) \\ & & \downarrow & & \downarrow \\ & & (S, \mathcal{M}_S) & \xrightarrow{i} & (S', \mathcal{M}_{S'}) \end{array}$$

Note that the front and back squares in (2.5.1) are cartesian squares. Let $\mathbf{L}_{X_S/S}^{\log}$ be the logarithmic cotangent complex of the log map $(X_S, \mathcal{M}_{X_S}) \rightarrow (S, \mathcal{M}_S)$ as in [Ols05]. By [Ols05, 5.9], we have the following results:

- (1) there is a canonical class $o \in \text{Ext}^1(f^* \mathbf{L}_{X_S/S}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0})$, whose vanishing is necessary and sufficient for the existence of a morphism f' fitting in the above diagram.
- (2) if $o = 0$, then the set of such maps f' is a torsor under $\text{Ext}^0(f^* \mathbf{L}_{X_S/S}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0})$.

Since the family of targets $X^{\log} \rightarrow B^{\log}$ is log flat, by [Ols05, 1.1(iv)] we have

$$\text{Ext}^1(f^* \mathbf{L}_{X_S/S}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0}) \cong \text{Ext}^1(f_0^* \mathbf{L}_{X_{S_0}/S_0}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0}).$$

Thus we define $\mathcal{D}_{\xi_A}(I) = \text{Ext}^0(f^* \mathbf{L}_{X_S/S}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0})$ and $\mathcal{O}_{\xi_0}(I) = \text{Ext}^1(f_0^* \mathbf{L}_{X_{S_0}/S_0}^{\log}, I \otimes_{A_0} \mathcal{O}_{C_0})$ to be the modules of deformations and obstructions respectively. Note that the log cotangent complex $\mathbf{L}_{X_S/S}^{\log}$ is bounded above with coherent cohomologies. The conditions of deformation and obstruction modules in [Art74, 5.2(4)] follows from the standard property of cohomology, see for example [AV02, 5.3.4].

2.6. Schlessinger's conditions. By [Art74, 5.2(2)], we need to verify Schlessinger's conditions (S1) and (S2) as in [Art74, section 2]. The condition (S2) follows from the cohomological description of the module of deformation \mathcal{D} . Next we check the condition (S1') [Art74, 2.3], which is a stronger version of (S1).

Indeed, consider an infinitesimal extension $A' \rightarrow A \rightarrow A_0$ as in Section 2.5, and a U -algebra homomorphism $B \rightarrow A$ such that the composition $B \rightarrow A_0$ is surjective. Consider $\xi_A \in \mathcal{LM}_U(A)$. For any surjection $R \rightarrow A$, denote by $\mathcal{LM}_{\xi_A}(R)$ the category of log maps over $\text{Spec } R$ whose restriction to $\text{Spec } A$ is ξ_A . Then we need to show that

$$\mathcal{LM}_{\xi_A}(A' \times_A B) \rightarrow \mathcal{LM}_{\xi_A}(A') \times \mathcal{LM}_{\xi_A}(B)$$

is an equivalence of categories.

First, consider the essential surjectivity. Consider objects $\xi_{A'} \in \mathcal{LM}_{\xi_A}(A')$ and $\xi_B \in \mathcal{LM}_{\xi_A}(B)$. Denote by $\xi_{A'} = (\zeta_{A'}, f_{A'})$ and $\xi_B = (\zeta_B, f_B)$, where $\zeta_{A'}$ and ζ_B are the corresponding log sources and targets as in Remark 2.1.7. Note that the two families $\zeta_{A'}$ and ζ_B correspond to maps $\text{Spec } A' \rightarrow U$ and $\text{Spec } B \rightarrow U$, which induce the same map $\text{Spec } A \rightarrow U$ by restricting to $\text{Spec } A$. Then we can glue them together to obtain a map $\text{Spec } B \times_A A' \rightarrow U$. This induces a family $\zeta_{B \times_A A'}$ over $\text{Spec } B \times_A A'$, whose restrictions to $\text{Spec } A'$ and $\text{Spec } B$ are $\zeta_{A'}$ and ζ_B respectively. Since the stack \mathcal{K} parameterizing the underlying maps is algebraic, the same argument as above produces a gluing $\underline{f}_{A' \times_A B}$ of $\underline{f}_{A'}$ and \underline{f}_B .

It remains to produce a compatible morphism of log structures $f_{A' \times_A B}^b$. Next we choose an affine open cover $V_{B \times_A A'} = \bigcup_i V_i$ of the log source curve in $\zeta_{B \times_A A'}$, its restrictions to A' and B give the affine open covers $V_{A'}$ and V_B for curves of $\zeta_{A'}$ and ζ_B respectively. Consider the stack

$$\mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_{A'}$$

induced by $\zeta_{A'}$ and the map $\underline{f}_{A'}$ as in Observation 2.4.2. Similarly, we have stack

$$\mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_B$$

induced by ζ_B and \underline{f}_B . They can be glued to give $\mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_{A' \times_A B}$ which corresponds to $\zeta_{A' \times_A B}$. Consider the maps $V_{A'} \rightarrow \mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_{A'}$ and $V_B \rightarrow \mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_B$ induced by $f_{A'}$ and f_B respectively. Note that these maps can be glued together and descend to a map

$$C_{A' \times_A B} \rightarrow \mathcal{L}^\Delta \times_{\mathcal{L}^\Delta} C_{A' \times_A B}.$$

This induce a map of log structures

$$f_{A' \times_A B}^b : \underline{f}_{A' \times_A B}^* \mathcal{M}_{X_{A' \times_A B}} \rightarrow \mathcal{M}_{C_{A' \times_A B}}.$$

By construction $f_{A' \times_A B}^b$ is compatible with $\zeta_{A'} \times_A B$ and the underlying map $\underline{f}_{A' \times_A B}$.

The full faithfulness follows from the representability of isomorphism functor of log maps.

2.7. Compatibility with formal completions. Let \hat{A} be a complete local ring, and m be the maximal ideal of \hat{A} . Denote by $A_n = \hat{A}/m^n$, $S = \text{Spec } \hat{A}$, and $S_n = \text{Spec } A_n$. Since we work over a fixed chart $U \rightarrow \mathcal{LM}$, it is enough to consider a family of log maps $\{\xi_n = (C_n \rightarrow S_n, X_{S_n} \rightarrow S_n, \mathcal{M}_S, f_n)\}_n$ such that $\xi_n \in \mathcal{LM}_U(S_n)$, and $\xi_n|_{S_k} = \xi_k$ for any $n \geq k$. According to [Art74, 5.2(3)], we need to show that there exists an element $\xi \in \mathcal{LM}_U(S)$, such that $\xi|_{S_n} = \xi_n$ for any n .

Denote by $\zeta_n = (C_n \rightarrow S_n, X_{S_n} \rightarrow S_n, \mathcal{M}_{S_n})$ the family of log sources and targets of ξ_n . For each n , there is a map $S_n \rightarrow U$ induced by ζ_n , such that they fit in the following commutative diagrams for any $k \leq n$:

$$\begin{array}{ccc} S_n & \longrightarrow & U \\ & \searrow & \nearrow \\ & S_k & \end{array}$$

Thus the above diagram induces a map $S \rightarrow U$, whose restriction to S_n is the map given by ζ_n as above. By pulling back the universal family over U , we obtain a family of log sources and targets $\zeta = (C \rightarrow S, X_S \rightarrow S, \mathcal{M}_S)$. Note that $\zeta|_{S_n} = \zeta_n$ for any n .

Denote by $\underline{\xi}_n$ the usual pre-stable map over S_n . Consider the family of compatible underlying maps $\{\underline{\xi}_n\}$. By [Gro61, 5.4.1], there exists a unique underlying map (up to a unique isomorphism) $\underline{f} : C \rightarrow X_S$ such that $\underline{f}|_{S_n} = \underline{f}_n$.

Now to construct ξ , we need to construct a log map $f : (C, \mathcal{M}_C) \rightarrow (X_S, \mathcal{M}_{X_S})$, which is compatible with the underlying map \underline{f} and f_n for all n . By definition of log maps, this is equivalent to constructing a map of log structures $f^\flat : \underline{f}^* \mathcal{M}_{X_S} \rightarrow \mathcal{M}_C$, which is compatible with f_n^\flat and the log structure \mathcal{M}_S on the base. For simplicity, denote by $\mathcal{M} = \underline{f}^* \mathcal{M}_{X_S}$.

Choose an affine étale cover of C , such that over each affine chart $V \rightarrow C$ the log structures $\mathcal{M}|_V$, $\mathcal{M}_C|_V$, and $\pi_1^* \mathcal{M}_S$ can be obtained by taking the log structure associated to $\Gamma(\mathcal{M}, V) \rightarrow \mathcal{O}_V$, $\Gamma(\mathcal{M}_C, V) \rightarrow \mathcal{O}_V$, and $\pi_1^* \mathcal{M}_S \rightarrow \mathcal{O}_V$ respectively. Since the charts of fine log structures always exist étale locally, the above choice of cover exists. We first construct f^\flat on such chart V .

Denote by $V_n = V \times_S S_n$. Then the canonical map $V_n \rightarrow C_n$ gives an affine étale chart. Consider the compatible families of monoids $\{\Gamma(\mathcal{M}_n, V_n)\}_n$ and $\{\Gamma(\mathcal{M}_{C_n}, V_n)\}_n$. For simplicity, let \mathcal{N} to be one of the monoids $\Gamma(\mathcal{M}, V)$, $\Gamma(\mathcal{M}_C, V)$, or $\Gamma(\pi_1^* \mathcal{M}_S, V)$, and \mathcal{N}_n to be the one of the corresponding reductions $\Gamma(\mathcal{M}_n, V_n)$, $\Gamma(\mathcal{M}_{C_n}, V_n)$, or $\Gamma(\pi_1^* \mathcal{M}_{S_n}, V_n)$. Denote by $q_n : \mathcal{N} \rightarrow \mathcal{N}_n$ the restriction map, and by $p_n : \varprojlim \mathcal{N}_n \rightarrow \mathcal{N}_n$ the canonical map. Let $p_{nk} : \mathcal{N}_n \rightarrow \mathcal{N}_k$ be the restriction map for all $k \leq n$. Assume that $V_n = \text{Spec } R_n$, and $R = \varprojlim R_n$. Thus, we write $V = \text{Spec } R$.

Note that inverse limit exists in the category of monoids, and their formation commutes with the forgetful functor to the category of sets ([Ogu06, Chapter I, 1.1]). Furthermore, the inverse limit of a family of integral monoids is again integral ([Ogu06, Chapter I, 1.2]). Consider an element $e \in \mathcal{N}$. This induces a family of compatible elements $\{q_n(e)\}_n \in \varprojlim \mathcal{N}_n$. In this way, we obtain a canonical map of integral monoids:

$$p : \mathcal{N} \rightarrow \varprojlim \mathcal{N}_n.$$

Lemma 2.7.1. (1) *Consider an element $e \in \mathcal{N}_n$. Then $p_{nk}(e) \in R_k^*$ for some $k \leq n$ if and only if $e \in R_n^*$. Furthermore, the map p_{nk} induces a natural isomorphism $\bar{p}_{nk} : \mathcal{N}_n/R_n^* \rightarrow \mathcal{N}_k/R_k^*$.*

- (2) Consider an element $e \in \mathcal{N}$. Then $q_n(e) \in R_n^*$ for some n if and only if $e \in R^*$. Furthermore, the map q_n induces a natural isomorphism of monoids $\bar{q}_n : \mathcal{N}/R^* \rightarrow \mathcal{N}_n/R_n^*$.
- (3) There is a natural inclusion $R^* \hookrightarrow \varprojlim \mathcal{N}_n$, which fits in the following commutative diagram

(2.7.1)

$$\begin{array}{ccc} & R^* & \\ \swarrow & & \searrow \\ \mathcal{N} & \xrightarrow{p} & \varprojlim \mathcal{N}_n \end{array}$$

where the left side arrow is the natural inclusion of units given by the corresponding log structures.

- (4) The canonical projection $p_n : \varprojlim \mathcal{N}_n \rightarrow \mathcal{N}_n$ induces an isomorphism of monoids

$$\bar{p}_n : (\varprojlim \mathcal{N}_n)/R^* \rightarrow \mathcal{N}_n/R_n^*.$$

- (5) The canonical map $p : \mathcal{N} \rightarrow \varprojlim \mathcal{N}_n$ induces an isomorphism of monoids

$$\bar{p} : \mathcal{N}/R^* \rightarrow (\varprojlim \mathcal{N}_n)/R^*.$$

Proof. The first part of Statement (1) follows from the following commutative diagram

$$\begin{array}{ccc} \mathcal{N}_n & \xrightarrow{p_{nk}} & \mathcal{N}_k \\ \downarrow & & \downarrow \\ R_n & \longrightarrow & R_k, \end{array}$$

where the two vertical maps are given by the structure morphism of the corresponding log structures. This immediately implies the existence of \bar{p}_{nk} . The surjectivity of p_{nk} for any $k \leq n$ implies that \bar{p}_{nk} is also surjective. To see the injectivity, consider two elements $a, b \in \mathcal{N}_n$ such that $p_{nk}(a) = p_{nk}(b) + \log u$ for some $u \in R_k^*$. Without loss of generality, we can assume that $a + c' = b + c$ in \mathcal{N}_n . Thus $p_{nk}(c') = p_{nk}(c) + \log u$, which implies $p_{nk}(c' - c) \in R_k^*$, hence $c' - c \in R_n^*$. This proves the second part of Statement (1).

Statement (2) can be proved similarly as the first one.

To prove (3), consider $e \in \varprojlim \mathcal{N}_n$, which can be represented by a family of compatible elements $\{e_n \in \mathcal{N}_n\}_n$. Assume that $e_{n'} \in R_{n'}^*$ for some n' . Then the first statement implies that $e_n \in R_n^*$ for all n . Thus we have a unique element $e \in R^* \subset R$ such that $e|_{V_n} = e_n$. This induces a canonical map $R^* \hookrightarrow \varprojlim \mathcal{N}_n$. Now the commutativity of (2.7.1) can be checked directly.

In fact the above argument proves that $p_n(e) \in R_n^*$ for some $e \in \varprojlim \mathcal{N}_n$, if and only if $e \in R^*$. Thus we obtain a canonical map $\bar{p}_n : (\varprojlim \mathcal{N}_n)/R^* \rightarrow \mathcal{N}_n/R_n^*$. Note that we have the following commutative diagram

(2.7.2)

$$\begin{array}{ccc} \mathcal{N} & & \\ \downarrow p & \searrow q_n & \\ \varprojlim \mathcal{N}_n & \xrightarrow{p_n} & \mathcal{N}_n \end{array}$$

The surjectivity of q_n implies that \bar{p}_n is also surjective. The injectivity of \bar{p}_n can be proved similarly as for the first statement. This proves (4).

Finally, note that (2.7.2) induces a commutative diagram

$$(2.7.3) \quad \begin{array}{ccc} \mathcal{N}/R^* & & \\ \bar{p} \downarrow & \searrow \bar{q}_n & \\ (\varprojlim \mathcal{N}_n)/R^* & \xrightarrow{\bar{p}_n} & \mathcal{N}_n/R_n^*. \end{array}$$

Since both \bar{q}_n and \bar{p}_n are isomorphisms of monoids, we conclude that \bar{p} is also an isomorphism. This proves (5). \square

Proposition 2.7.2. *The map of monoids $p : \mathcal{N} \rightarrow \varprojlim \mathcal{N}_n$ is an isomorphism.*

Proof. By Lemma 2.7.1(3) and (5), we have a commutative diagram

$$\begin{array}{ccc} \mathcal{N} & \xrightarrow{p} & \varprojlim \mathcal{N}_n \\ \downarrow & & \downarrow \\ \mathcal{N}/R^* & \xrightarrow{\bar{p}} & \varprojlim \mathcal{N}_n/R_n^*. \end{array}$$

Pick two sections $e, e' \in \mathcal{N}$ such that $p(e) = p(e')$. Denote by \bar{e} and \bar{e}' the corresponding images in \mathcal{N}/R^* . It follows from Lemma 2.7.1(5) that $\bar{e} = \bar{e}'$. Thus, we have $e = e' + \log u$ for some $u \in R^*$. But the assumption $p(e) = p(e')$ implies that $p(u) = 1 \in \varprojlim \mathcal{N}_n$. By Lemma 2.7.1(3), we have $u = 1$. This proves that p is also injective.

To prove the surjectivity, consider an element $a \in \varprojlim \mathcal{N}_n$. Since \bar{p} is an isomorphism, denote by \bar{a} the image of a in \mathcal{N}/R^* . Let a' be a lifting of \bar{a} in \mathcal{N} . Then there exists an element $u \in R^*$ such that $a = p(a') + \log u$. Thus $p(a' + \log u) = a$. \square

Pick an element $\{e_n \in \Gamma(\mathcal{M}_n, V_n)\}_n \in \varprojlim \Gamma(\mathcal{M}_n, V_n)$. We obtain a compatible family $\{f_n(e_n)\}_n \in \varprojlim \Gamma(\mathcal{M}_{C_n}, V_n)$. Thus the compatible morphism of log structures $\{f_n\}$ induces a natural map of monoids

$$\varprojlim \Gamma(\mathcal{M}_n, V_n) \rightarrow \varprojlim \Gamma(\mathcal{M}_{C_n}, V_n).$$

By Proposition 2.7.2, we have a natural map of monoids:

$$\Gamma(f^b, V) : \Gamma(\mathcal{M}, V) \rightarrow \Gamma(\mathcal{M}_C, V).$$

Next we show that $\Gamma(f^b, V)$ induces a map of log structures $f_V^b : \mathcal{M}|_V \rightarrow \mathcal{M}_C|_V$. Since the two log structures $\mathcal{M}|_V$ and $\mathcal{M}_C|_V$ can be obtained from $\Gamma(\mathcal{M}, V)$ and $\Gamma(\mathcal{M}_C, V)$ respectively, it is enough to show that the following diagram is commutative:

$$\begin{array}{ccc} \Gamma(\mathcal{M}, V) & \xrightarrow{\Gamma(f^b, V)} & \Gamma(\mathcal{M}_C, V) \\ \searrow \exp_1 & & \swarrow \exp_2 \\ & R & \end{array}$$

where \exp_1 and \exp_2 are the structure morphism of the corresponding log structures. To see this, consider any section $s \in \Gamma(\mathcal{M}, V)$. Since $\exp_1(s)|_{S_n} = \exp_1 \circ \Gamma(f^b, V)(s)|_{S_n}$ for any n , we have $\exp_1(s) = \exp_1 \circ \Gamma(f^b, V)(s)$. This proves the commutativity.

We claim that f_V^b is compatible with the log structure on the base. This is equivalent to showing the commutativity of the following diagram of log structures on V :

$$(2.7.4) \quad \begin{array}{ccc} & \pi_1^* \mathcal{M}_S|_V & \\ \pi_2^b \swarrow & & \searrow \pi_1^b \\ \mathcal{M}|_V & \xrightarrow{f_V^b} & \mathcal{M}_C|_V, \end{array}$$

where $\pi_1 : C \rightarrow S$ and $\pi_2 : X_S \rightarrow S$ are the projections. Since $\pi_1^* \mathcal{M}_S|_V$ can be obtained by taking the log structure associated to $\Gamma(\pi_1^* \mathcal{M}_S, V) \rightarrow \mathcal{O}_V$. Hence to verify the commutativity of (2.7.4) it is enough to show that the following diagram is commutative:

$$\begin{array}{ccc} & \Gamma(\pi_1^* \mathcal{M}_S, V) & \\ \pi_2^b \swarrow & & \searrow \pi_1^b \\ \Gamma(\mathcal{M}, V) & \xrightarrow{f_V^b} & \Gamma(\mathcal{M}_C, V). \end{array}$$

But This follows from the definition of f_V^b , and the following commutative diagram for each n :

$$\begin{array}{ccc} & \pi_{1,n}^* \mathcal{M}_{S_n} & \\ \swarrow & & \searrow \\ f^* \mathcal{M}_n & \xrightarrow{f_n^b} & \mathcal{M}_{C_n}. \end{array}$$

Thus we obtain the desired map f_V^b over each affine chart V .

Finally, notice that the construction of f_V^b is functorial. Hence, we are able to obtain a global map f^b by gluing f_V^b on each affine chart. This finishes the proof of compatibility with formal completions.

3. MINIMAL LOGARITHMIC MAPS TO RANK ONE DELIGNE-FALTINGS LOG PAIRS

3.1. Basic definitions and notations.

Definition 3.1.1. We call the log scheme $X^{\log} = (X, \mathcal{M}_X)$ a *rank one Deligne-Faltings pair*, if

- (1) X is a projective variety over \mathbb{C} ;
- (2) \mathcal{M}_X is a DF log structure on X as in Definition A.2.1, with a global presentation $\mathbb{N} \rightarrow \overline{\mathcal{M}}_X$.

Remark 3.1.2. The results in Section 3 and 4 still hold if we assume X to be only separated of finite type over \mathbb{C} . However, the projectivity is essential for the properness of the stack $\mathcal{K}_\Gamma(X)$ as in Definition 3.6.5.

Conventions 3.1.3. In the rest of this paper, we fix a Deligne-Faltings pair (X, \mathcal{M}_X) as our target of log maps, with a global presentation $\mathbb{N} \rightarrow \overline{\mathcal{M}}_X$. Denote by (L, s) the pair consisting of a line bundle L , and a morphism of sheaves $s : L \rightarrow \mathcal{O}_X$ corresponding to \mathcal{M}_X . Let D be the vanishing locus of the section $s \in H^0(L^\vee)$. Denote by δ the standard generator of \mathbb{N} . For convenience, locally we identify δ with its image in $\overline{\mathcal{M}}_X$.

Remark 3.1.4. Note that if $s = 0$, then $D = X$. If s is not a zero section, then D is a divisor in X . Thus, we have $L = \mathcal{O}_X(-D)$, with the natural inclusion $s : \mathcal{O}_X(-D) \hookrightarrow \mathcal{O}_X$. The section δ locally lifts to a section in \mathcal{O}_X , whose vanishing locus gives the divisor D .

Remark 3.1.5. The target X^{\log} should be viewed as a log scheme over a point with trivial log structures. Thus, we can simplify the notations in Section 2.1 as follows. A log map over a usual scheme S is given by the triple $(C \rightarrow S, \mathcal{M}_S, f)$, where $(C \rightarrow S, \mathcal{M}_S)$ is a log curve in Definition B.2.2, and $f : (C, \mathcal{M}_C) \rightarrow (X, \mathcal{M}_X)$ is a log map.

Consider two log maps $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ and $\xi' = (C' \rightarrow S, \mathcal{M}'_S, f')$ over a scheme S . An arrow $\xi \rightarrow \xi'$ over S is a pair (ρ, θ) as in Definition B.3.1 such that the following diagram commutes:

$$\begin{array}{ccc}
 & & (X, \mathcal{M}_X) \\
 & \nearrow f & \\
 (C, \mathcal{M}_C) & \xrightarrow{\rho} & (C', \mathcal{M}_{C'}) & \nearrow f' \\
 \downarrow & & \downarrow & \\
 (S, \mathcal{M}_S) & \xrightarrow{\theta} & (S, \mathcal{M}'_S) &
 \end{array}$$

where the square is a cartesian square of log schemes. This is compatible with Definition 2.1.3 and 2.1.4.

3.2. Log maps on the level of characteristic. Consider a log map $\xi = (\pi : C \rightarrow S, \mathcal{M}_S, f)$ as in Remark 3.1.5, where $S = \text{Spec } k$ is a geometric point and $(C \rightarrow S, \mathcal{M}_S)$ is a log pre-stable curve. Pick a point $p \in C$, which sits in an irreducible component Z . We have a map of characteristic monoids:

$$(3.2.1) \quad \bar{f}^b : f^*(\overline{\mathcal{M}}_X)_p \rightarrow \overline{\mathcal{M}}_{C,p}.$$

First consider the case p is a smooth non-marked point. By the description in Definition B.2.1, we have $\bar{f}^b(\delta) = e \in \overline{\mathcal{M}}_S$ at p . By [Ols03a, 3.5(i),(iii)], the equality $\bar{f}^b(\delta) = e$ lifts to an étale neighborhood of p .

Definition 3.2.1. We call e the *degeneracy* of Z . Note that if $p \notin D$ for some $p \in Z$, then the image e vanishes in $\overline{\mathcal{M}}_S$. A component Z is called *degenerate* if its degeneracy is not zero. This is equivalent to saying that Z maps to D via f .

Next, we consider the case where p is a marked point. Locally at p , we have $\mathcal{M}_C \cong \pi^* \mathcal{M}_S \oplus_{\mathcal{O}_C^*} \mathcal{N}$, where \mathcal{N} is the canonical log structure associated to the marked point p . Then on the level of characteristic, we have

$$(3.2.2) \quad \bar{f}^b(\delta) = e + c_p \cdot \sigma_p,$$

where $e \in \overline{\mathcal{M}}_S$, the element σ_p is the generator of $\overline{\mathcal{N}}_p$, and c_p is a non-negative integer.

Observation 3.2.2. When we generalize (3.2.2) to nearby smooth points, any lifting of σ_p in the structure sheaf becomes invertible. Thus, the element e is the degeneracy of the component Z containing p .

Definition 3.2.3. We call c_p the *contact order* of f at p .

Lemma 3.2.4. Consider a log map $\xi = (C' \rightarrow S', \mathcal{M}_{S'}, g)$ over a scheme S' , and a marking Σ_i on C' . There is an open subset in S' , such that the contact order along the fixed marking Σ_i is constant.

Proof. Consider the relative characteristic $\overline{\mathcal{M}}_{C'/S'}$. This is a locally constant sheaf along Σ_i , with stalks given by \mathbb{N} . Thus along Σ_i there is a map of locally constant sheaves $g^*\overline{\mathcal{M}}_X \rightarrow \overline{\mathcal{M}}_{C'/S'}$, which locally at $p \in \Sigma_i$ is given by $\mathbb{N} \rightarrow \mathbb{N}$ by $1 \mapsto c$, for some positive integer c . Note that the correspondence $1 \mapsto c$ can be generalize to the nearby points of p . Therefore it forms an open condition on the base. \square

Remark 3.2.5. When D is a divisor, the contact order of a marked point Σ in a non-degenerate component can be identified with the local intersection multiplicity $(C \cdot D)_\Sigma$.

Finally, let us consider the case where p is a node joining two irreducible components Z and Z' . Let e_p be the element in $\overline{\mathcal{M}}_S$ smoothing the node p . Denote by $\log x_p$ and $\log y_p$ the elements in $\overline{\mathcal{M}}_C$ given by the local coordinates of the two components Z and Z' at p respectively as in Section B.4. Then locally at p we have the equation in $\overline{\mathcal{M}}_C$:

$$(3.2.3) \quad e_p = \log x_p + \log y_p.$$

Thus, without loss of generality we can assume that

$$(3.2.4) \quad \bar{f}^b(\delta) = e + c_p \cdot \log x_p,$$

where c_p is a positive integer.

Definition 3.2.6. The integer c_p is called *the contact order of f at the node p* . If $c_p \neq 0$, then p is called a *distinguished node*. A point $p \in C$ is called a *distinguished point*, if it is a marked point or node with non-trivial contact order. Otherwise, it is called *non-distinguished point*.

Lemma 3.2.7. *Using notations as above, the degeneracy of Z is e , and the degeneracy of Z' is $e + c_p \cdot e_p$.*

Proof. When we generalize (3.2.4) to a smooth point in Z' , the section y becomes invertible. Then the statement for Z' follows from the definition of degeneracy of a component. For Z , the proof is similar. \square

Lemma 3.2.7 gives a way to put a partial order on the set of irreducible components as follows:

Definition 3.2.8. Using the notations as above, we call Z the *lower component* of p , and Z' the *upper component* of p .

Lemma 3.2.9. *Consider a log map $\xi = (C' \rightarrow S', \mathcal{M}_{S'}, g)$, and a connected singularity $p \subset C'$. There is an open subset in S' , such that over each fiber we have that either the node p is smoothed out, or its contact order remains the same.*

Proof. The proof is similar to the one for Lemma 3.2.4. \square

3.3. Marked graph. We next introduce the marked graph, which will be used later to describe the combinatorial data associated to log maps.

Definition 3.3.1. A *weighted graph* G is a connected graph with the following data:

- (1) A subset $V_n(G) \subset V(G)$ of the set of vertices of G , which is called the *set of nondegenerate vertices*.
- (2) For each edge $l \in E(G)$, we associate a non-negative integer weight c_l called the *contact order* of l .

These data are subject to the only condition that if the edge l is a loop, then $c_l = 0$.

Note that the set $V_n(G)$ can be empty. If the contact order of an edge l is zero, then l is called the *non-distinguished edge*, otherwise is called a *distinguished edge*. Two vertices are called *adjacent* if they are connected by an edge. Denote by \underline{G} the underlying graph of G , obtained by removing all weights.

Definition 3.3.2. Consider a weighted graph G as in the above definition. An *orientation* on G is an orientation on the underlying graph \underline{G} , except that we allow some edges to be *non-oriented*, i.e. an edge with two directions. Consider an edge l from v_1 to v_2 under the orientation. Then v_1 is called the *initial vertex of l* , and v_2 is called the *end vertex of l* . We denote this by $v_1 \leq v_2$. If l is orientated, then we write $v_1 < v_2$.

An orientation on G is called *compatible* if

- (1) An edge $l \in E(G)$ is non-oriented if and only if $c_l = 0$.
- (2) If $v \in V_n(G)$, then for any other adjacent vertex v' of v we have $v \leq v'$.

Note that if $v, v' \in V_n(G)$, then any edges between them is non-oriented. The graph G is called a *marked graph*, if it is a weighted graph with a compatible orientation.

A *path* is a non-repeated sequence of edges (l_1, l_2, \dots, l_m) , such that the end vertex of l_j is the initial vertex of l_{j+1} . Such a path is called a *cycle* if the initial vertex of l_1 is the end vertex of l_m . A cycle is called *strict* if it contains at least one oriented edges. A vertex $v \in V(G)$ is called *minimal* (respectively *maximal*) if it is not the end (respectively initial) vertex of any oriented edge. Thus by condition (2) above, any vertex $v \in V_n(G)$ is minimal.

Construction 3.3.3. Consider a marked graph G as in Definition 3.3.2. For each edge $l \in E(G)$ (respectively each vertex $v \in V(G)$), we introduce a variable e_l (resp. e_v), which is called the *element associated to l* (resp. v). For any $v \in V_n(G)$, we set

$$(3.3.1) \quad h_v : \quad e_v = 0.$$

Consider an edge $l \in E(G)$ with initial vertex v_1 and end vertex v_2 . We associate an equation

$$(3.3.2) \quad h_l : \quad e_{v_2} = e_{v_1} + c_l \cdot e_l.$$

Consider the monoid

$$(3.3.3) \quad M(G) = \left\langle e_v, e_l \mid v \in V(G), l \in E(G) \right\rangle \Big/ \left\langle h_l, h_v \mid l \in E(G), v \in V_n(G) \right\rangle$$

Denote by $T(G)$ the torsion part of $M(G)^{gp}$. Then we have the following composition

$$M(G) \rightarrow M(G)^{gp} \rightarrow M(G)^{gp}/T(G).$$

Denote by $N(G)$ the image of $M(G)$ in $M(G)^{gp}/T(G)$, and $\overline{M}(G)$ the saturation of $N(G)$ in $M(G)^{gp}/T(G)$.

Definition 3.3.4. The monoid $\overline{M}(G)$ constructed above is called the *associated monoid* of the marked graph G .

Note that $N(G)$ is the image of $M(G)$ in $\overline{M}(G)$. By the definition of $\overline{M}(G)$ and Proposition A.1.1, we have the following:

Lemma 3.3.5. *By viewing $N(G)$ and $\overline{\mathcal{M}}(G)$ as sub-monoids of $\overline{\mathcal{M}}(G)^{gp} = M(G)^{gp}/T(G)$, we have that for any $a \in \overline{\mathcal{M}}(G)$ there exists $b \in N(G)$ and a positive integer m such that $b = m \cdot a$.*

Definition 3.3.6. The marked graph G is called *admissible* if $\overline{\mathcal{M}}(G)$ is a sharp monoid, and the image of e_l in $\overline{\mathcal{M}}(G)$ is non-trivial for all $l \in E(G)$.

Corollary 3.3.7. *If G is admissible, then there is no strict cycle in G .*

Proof. If there is a strict cycle (l_1, \dots, l_k) , then we have $\sum_{i=1}^k c_{l_i} e_{l_i} = 0$. The strictness implies that at least one of the c_{l_i} is non-zero. Thus, the monoid $\overline{\mathcal{M}}(G)$ fails to be sharp, which contradicts the admissibility assumption. \square

Note that when G is admissible, the monoid $\overline{\mathcal{M}}(G)$ generates a strongly convex rational cone $C(\overline{\mathcal{M}}(G))$ in the vector space $\overline{\mathcal{M}}(G)^{gp} \otimes \mathbb{Q}$ (see [Ful93, Page 4]).

Lemma 3.3.8. *Consider an irreducible element $a \in \overline{\mathcal{M}}(G)$, where G is admissible. Then at least one of the following holds:*

- (1) *There is a positive integer n and a minimal vertex v , such that $n \cdot a = e_v$.*
- (2) *There is a positive integer n and an edge l , such that $n \cdot a = e_l$.*

Proof. Let n be the minimal positive integer such that $n \cdot a \in N(G)$. Assume that $n \cdot a = b + c$ with $b, c \in N(G)$. Since a is an irreducible element of $\overline{\mathcal{M}}(G)$, and G is admissible, then a generates a ray, which is extremal in the strongly convex rational cone $C(\overline{\mathcal{M}}(G))$. Thus we have positive numbers n_1 and n_2 such that $b = n_1 \cdot a$ and $c = n_2 \cdot a$. The minimality of n implies that either $b = 0$ or $c = 0$. Since b and c in $N(G)$ are elements associated to edges or vertices, the element $n \cdot a$ must satisfy one of the two possibilities above. \square

3.4. Marked graphs associated to log maps. Consider a log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a geometric point S , such that the log structure \mathcal{M}_S is fs.

Construction 3.4.1. We construct a weighted graph G_ξ of ξ with an orientation as in Definition 3.3.2:

- (1) The underlying graph \underline{G}_ξ is given by the dual graph of the curve C .
- (2) The subset $V_n(G)$ consists of the vertices corresponding to non-degenerate components.
- (3) For each edge $l \in E(G_\xi)$, we associate a non-negative integers c_l , where c_l is the contact order of the node l as in Definition 3.2.6.
- (4) Let $l \in E(G_\xi)$ be a node joining two irreducible components $v_1, v_2 \in V(G_\xi)$. Then we define an orientation by putting $v_1 \leq v_2$ if v_1 is the lower component, and v_2 is the upper component of l as in Definition 3.2.8.

Consider a node $l \in E(G_\xi)$. Denote by e'_l the element in $\overline{\mathcal{M}}_S$ smoothing l , and by e_l the element associated to l in $\overline{\mathcal{M}}(G_\xi)$. Then consider an irreducible component $v \in V(G_\xi)$. Denote by e'_v the degeneracy of v in ξ , and by $e_v \in \overline{\mathcal{M}}(G_\xi)$ the element associated to v . We define a correspondence

$$(3.4.1) \quad e_l \mapsto e'_l \quad \text{and} \quad e_v \mapsto e'_v$$

Proposition 3.4.2. *Assuming that \mathcal{M}_S is fs, the correspondence (3.4.1) induces a canonical morphism of monoids*

$$\phi : \overline{\mathcal{M}}(G_\xi) \rightarrow \overline{\mathcal{M}}_S.$$

Proof. Note that (3.4.1) induces a map $M(G_\xi) \rightarrow \overline{\mathcal{M}}_S$. By Proposition A.1.1, this induces a unique map $\phi' : M(G_\xi)^{Sat} \rightarrow \overline{\mathcal{M}}_S$. Since the monoid $\overline{\mathcal{M}}_S$ is sharp, if $e \in M(G_\xi)^{Sat}$ is torsion, then $\phi'(e) = 0$. Thus, there is a unique map $\phi : \overline{\mathcal{M}}(G_\xi) \rightarrow \overline{\mathcal{M}}_S$ induced by ϕ' . \square

Corollary 3.4.3. *The graph G_ξ is an admissible marked graph.*

Proof. The compatibility of the orientation follows from Lemma 3.2.7. Let us consider the admissibility. First note that e_l is non-trivial for any $l \in E(G)$, since its image in $\overline{\mathcal{M}}_S$ is the element smoothing the node l , which is non-trivial. For any element $a \in \overline{\mathcal{M}}(G_\xi)$, if a is invertible, then by Lemma 3.3.5, there exists some positive integer m such that $m \cdot a = \sum_i d_i e_i$, where e_i are elements associated to some edges or vertices, and d_i are non-negative integers. Since the monoid $\overline{\mathcal{M}}_S$ is sharp, we have $\phi(a) = \sum_i d_i \phi(e_i) = 0$ in $\overline{\mathcal{M}}_S$. If $d_i \neq 0$, then $\phi(e_i) = 0$, which implies that e_i is the element associated to a non-degenerate component. Thus we have $e_i = 0$ in $\overline{\mathcal{M}}(G_\xi)$. This implies that $a = 0$ in $\overline{\mathcal{M}}(G_\xi)$, which proves the statement. \square

Definition 3.4.4. We call G_ξ the *marked graph of ξ* .

3.5. Minimal logarithmic maps. We still consider a log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a geometric point S with fs log structure \mathcal{M}_S .

Definition 3.5.1. The log map ξ over S is called *minimal* if the induced canonical map ϕ in Proposition 3.4.2 is an isomorphism of monoids. A family of log maps ξ_T over a scheme T is called *minimal* if each geometric fiber is minimal.

Proposition 3.5.2 (Openness of minimal log maps). *Let $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ be a family of log maps over a scheme S , and assume that $\xi_{\bar{s}}$ is minimal for some point $s \in S$. Then there exists an étale neighborhood of s with all geometric fibers minimal.*

Proof. Shrinking S , we may assume that S is connected, and we have a chart $\beta : \overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \mathcal{M}_S$ by Proposition A.1.3. We next show that for any $\bar{t} \in S$, the fiber $\xi_{\bar{t}}$ is minimal.

Denote by

$$K_{\bar{t}} = \{ a \in \overline{\mathcal{M}}_{S, \bar{s}} \mid \beta(a) \text{ is invertible at } \bar{t} \}.$$

Note that $K_{\bar{t}}$ is a submonoid of $\overline{\mathcal{M}}_{S, \bar{s}}$. Consider the following composition

$$\overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \overline{\mathcal{M}}_{S, \bar{s}}^{gp} \rightarrow \overline{\mathcal{M}}_{S, \bar{s}}^{gp} / K_{\bar{t}}^{gp}.$$

By [Ols03a, 3.5], we have $\overline{\mathcal{M}}_{S, \bar{s}}^{gp} / K_{\bar{t}}^{gp} \cong \overline{\mathcal{M}}_{S, \bar{t}}^{gp}$. The above composition induces a map $q : \overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \overline{\mathcal{M}}_{S, \bar{t}}$, which is exactly the specialization map as in [Ols03a, 3.5(iii)]. We construct a new graph from the marked graph $G_{\xi_{\bar{s}}}$ as follows.

- (1) For an edge $l \in E(G_{\xi_{\bar{s}}})$, if $q(e_l) = 0$, then we contract l , and identify the two end vertices of l and the corresponding associated elements.
- (2) For a vertex $v \in V(G_{\xi_{\bar{s}}})$, if $q(e_v) = 0$, then we put $e_v = 0$ in G' .

Other vertices and edges in $G_{\xi_{\bar{s}}}$, and their associated elements and contact orders remain the same. We denote by G' the resulting graph.

First note that the underlying graph \underline{G}' is the dual graph of $C_{\bar{t}}$, since an edge $l \in E(G_{\xi_{\bar{s}}})$ gets contracted if and only if the corresponding node is smoothed out over \bar{t} . Furthermore, the orientation on $G_{\xi_{\bar{s}}}$ induces a natural orientation on G' . Since all contact orders remain the same, the graph G' is in fact the marked graph $G_{\xi_{\bar{t}}}$ of $\xi_{\bar{t}}$.

The construction of G' gives a map of monoids:

$$(3.5.1) \quad M(G_{\xi_{\bar{s}}}) \rightarrow M(G_{\xi_{\bar{t}}}) \rightarrow \overline{\mathcal{M}}_{G_{\xi_{\bar{t}}}}.$$

By the same argument in Proposition 3.4.2, we obtain a canonical map of monoids:

$$(3.5.2) \quad q' : \overline{\mathcal{M}}_{S, \bar{s}} \cong \overline{\mathcal{M}}(G_{\xi_{\bar{s}}}) \rightarrow \overline{\mathcal{M}}(G_{\xi_{\bar{t}}}),$$

which gives the following commutative diagram:

$$(3.5.3) \quad \begin{array}{ccc} & \overline{\mathcal{M}}_{S, \bar{s}} & \\ q' \swarrow & & \searrow q \\ \overline{\mathcal{M}}(G_{\xi_{\bar{t}}}) & \xrightarrow{\quad} & \overline{\mathcal{M}}_{S, \bar{t}}, \end{array}$$

where the bottom map is the canonical map as in Proposition 3.4.2. Consider the induced commutative diagram:

$$(3.5.4) \quad \begin{array}{ccc} & \overline{\mathcal{M}}_{S, \bar{s}}^{gp} & \\ (q')^{gp} \swarrow & & \searrow q^{gp} \\ \overline{\mathcal{M}}(G_{\xi_{\bar{t}}})^{gp} & \xrightarrow{\quad} & \overline{\mathcal{M}}_{S, \bar{t}}^{gp}, \end{array}$$

Note that both q^{gp} and $(q')^{gp}$ are surjective maps. By the construction of q , the group $K_{\bar{t}}^{gp}$ is the kernel of q^{gp} . On the other hand, the construction of G' and the fact that $G' = G_{\xi_{\bar{t}}}$ implies that $K_{\bar{t}}^{gp}$ is also the kernel of $(q')^{gp}$. Since the monoids in (3.5.3) are fine and saturated, the map $\overline{\mathcal{M}}(G_{\xi_{\bar{t}}}) \rightarrow \overline{\mathcal{M}}_{S, \bar{t}}$ is an isomorphism. This proves the statement. \square

Definition 3.5.3. Denote by $\mathcal{K}_{g,n}^{min}(X^{log})$ the stack parameterizing minimal log maps to X^{log} , with the fixed genus g , and n -markings.

Corollary 3.5.4. $\mathcal{K}_{g,n}^{min}(X^{log})$ is an open substack of the stack $\mathcal{LM}_{g,n}(X^{log})$ of log maps, hence is an algebraic stack.

Proof. This follows from Theorem 2.1.10 and Proposition 3.5.2. \square

3.6. Log stable maps.

Definition 3.6.1. A log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a geometric point S is called *log stable* if its underlying map is stable in the usual sense, and \mathcal{M}_S is fs. A family of log maps ξ_T over a scheme T is called log stable if its geometric fibers are log stable. A log stable map is called *minimal log stable* if it satisfies the minimality condition as in Definition 3.5.1.

Similarly, we can work over log schemes rather than the usual schemes. Then we have the following:

Definition 3.6.2. A log map ξ^{log} over a fs log scheme (S, \mathcal{M}_S) as in Definition 2.1.2 is called stable, if its underlying map is stable in the usual sense.

Conventions 3.6.3. In this paper, we fix the discrete data $\Gamma = (\beta, g, n, \mathbf{c})$ where

- (1) $\beta \in H^2(X, \mathbb{Z})$ is a curve class in X ;
- (2) n and g are two non-negative integers;

(3) $\mathbf{c} = (c_i)_{i=1}^n$ is a set of non-negative integers such that

$$(3.6.1) \quad \sum_{i=1}^n c_i = c_1(L^\vee) \cap \beta$$

where $c_1(L^\vee)$ is the first chern class of the line bundle L^\vee as in Conventions 3.1.3.

Definition 3.6.4. A minimal log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a geometric point S is called a Γ -minimal log stable if

- (1) The source curve $(C \rightarrow S, \mathcal{M}_S)$ is a log pre-stable curve of genus g with n marked points.
- (2) $f_*[C] = \beta$.
- (3) For any $i \in \{1, 2, \dots, n\}$, the contact order along section Σ_i is given by c_i .

A log map ξ' over an arbitrary scheme T is called Γ -minimal log stable if its geometric fibers are all Γ -minimal log stable. The arrows between log stable maps are the same as the arrow of log maps in Definition 2.1.4.

Definition 3.6.5. Denote by $\mathcal{K}_{g,n}^{mst}(X^{log}, \beta)$ the stack parameterizing minimal log stable maps with genus g , n marked points, and curve class β . Let $\mathcal{K}_\Gamma(X^{log})$ be the stack parameterizing Γ -minimal log stable maps. These are substacks of $\mathcal{LM}_{g,n}(X^{log})$ as in Theorem 2.1.10.

Theorem 3.6.6. *The stack $\mathcal{K}_{g,n}^{mst}(X^{log}, \beta)$ is an open substack of $\mathcal{K}_{n,g}^{min}(X^{log})$, hence is algebraic.*

Proof. Note that the stability condition is a condition on the underlying map, which is well known to be open. \square

Remark 3.6.7. Denote by Λ the set of discrete data Γ as in Convention 3.6.3 with fixed g , n and β . Note that Λ is a finite set. By Lemma 3.2.4, we have the disjoint union of open and closed substacks

$$\mathcal{K}_{g,n}^{mst}(X^{log}, \beta) = \coprod_{\Gamma \in \Lambda} \mathcal{K}_\Gamma(X^{log}).$$

3.7. A quasi-finiteness result. We fix a minimal log stable map $\xi_1 = (\pi : C \rightarrow S, \mathcal{M}_1, f_1)$ over a geometric point S . Denote by G the marked graph of ξ_1 . Choose a chart $\beta : \overline{\mathcal{M}}(G) \rightarrow \mathcal{M}_S$. Let $\mathcal{N}_1 \subset \mathcal{M}_S$ be the sub-log structure generated by $\beta(N(G))$. Since different choice of β only differ by invertible elements, the log structure \mathcal{N}_1 does not depend on the choice of β .

Consider the fine log scheme (S, \mathcal{N}_1) with the sub-log structure $\mathcal{N}_1 \subset \mathcal{M}_1$ induced by $N(G)$. Since the map of characteristics $\overline{\mathcal{M}}_S^{C/S} \rightarrow \overline{\mathcal{M}}(G)$ factors through $N(G)$, the structure map $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S$ factors through \mathcal{N}_1 . This induces a log curve $(C \rightarrow S, \mathcal{N}_1)$. Denote by $\mathcal{N}_{C,1}$ the log structure of $(C \rightarrow S, \mathcal{N}_1)$ on C , and by $\mathcal{M}_{C,1}$ the log structure of ξ_1 on C . Then $\mathcal{N}_{C,1}$ is a sub-log structure of $\mathcal{M}_{C,1}$. Again by considering the map of characteristics, we see that the log map $f_1^\flat : f_1^* \mathcal{M}_X \rightarrow \mathcal{M}_{C,1}$ factors through $\mathcal{N}_{C,1}$. Then we obtain a log map $g_1 : (C, \mathcal{N}_{C,1}) \rightarrow X^{log}$. Denote by $\xi'_1 = (C \rightarrow S, \mathcal{N}_1, g_1)$ the log map over S .

Definition 3.7.1. The log map ξ'_1 is called the *coarse log map* of ξ_1 .

Remark 3.7.2. The log structures of coarse log maps are in general not saturated. The above construction yields a natural arrow $\xi_1 \rightarrow \xi'_1$.

Corollary 3.7.3. *The pair $(\xi_1, \xi_1 \rightarrow \xi'_1)$ is unique up to a unique isomorphism*

Proof. This follows from the uniqueness of the log structure \mathcal{N}_1 . \square

The following result reveals the importance of the notion of coarse log maps.

Lemma 3.7.4. *Consider another minimal log stable map $\xi_2 = (C \rightarrow S, \mathcal{M}_2, f_2)$ whose underlying structure and marked graph are identical to those of ξ_1 . Then there exists a canonical isomorphism of coarse log maps $\xi'_2 \cong \xi'_1$.*

Proof. Let $\xi'_2 = (C \rightarrow S, \mathcal{N}_2, g_2)$ be the coarse log map of ξ_2 . Denote by $\mathcal{N}_{C,i}$ the log structures on C corresponding to ξ'_i for $i = 1, 2$. Consider the solid diagram of log structures on C :

$$(3.7.1) \quad \begin{array}{ccc} & \mathcal{N}_{C,1} & \\ g_1^\flat \nearrow & \vdots & \nwarrow \phi_1 \\ \underline{f}^* \mathcal{M}_X & & \mathcal{M}_C^{C/S} \\ g_2^\flat \searrow & \downarrow \psi_{\mathcal{N}} & \swarrow \phi_2 \\ & \mathcal{N}_{C,2} & \end{array}$$

where $\mathcal{N}_{C,i}$ is the associated log structure on C with respect to \mathcal{N}_i . We will first construct the dashed arrow $\psi_{\mathcal{N}}$, which makes (3.7.1) commutative.

Since the underlying structures of ξ_1 and ξ_2 are identical, it would be enough to construct $\psi_{\mathcal{N}} : \pi^* \mathcal{N}_1 \rightarrow \pi^* \mathcal{N}_2$. We fix a chart $\beta_i : N(G) \rightarrow \mathcal{N}_i$. Consider a section $e \in \pi^* \mathcal{N}_i$ for $i = 1, 2$. We want to define the image $\psi_{\mathcal{N}}(e)$. For this, it is enough to assume that $e = \pi^*(\beta_1(\bar{e}))$ for some element $\bar{e} \in N(G)$ associated to a vertex or an edge.

We first assume that \bar{e} is an element associated to an edge $l \in E(G)$. Then there exists a section $e_l \in \mathcal{M}_S^{C/S}$, such that $\phi_1(e_l) = e$. Thus, to make (3.7.1) commutative one has to define

$$(3.7.2) \quad \psi_{\mathcal{N}}(e) = \phi_2(e_l)$$

We then consider the case that \bar{e} is an element associated to a vertex $v \in V(G)$. We can assume that v is degenerate, otherwise \bar{e} is trivial in $N(G)$. We may restrict (3.7.1) to a small neighborhood of a non-distinguished point as in Definition 3.2.6 on the component corresponding to v . Denote by $\delta \in \underline{f}^* \mathcal{M}_X$ a local generator. Since on the level of characteristic we have $\bar{\phi}_1(\delta) = \bar{e}$ in $\overline{\mathcal{N}}_{C,1}$, we may assume $f_1^\flat(\delta) = e$. Thus, one has to define

$$(3.7.3) \quad \psi_{\mathcal{N}}(e) = f_2^\flat(\delta)$$

It is clear that (3.7.2) and (3.7.3) make (3.7.1) commutative. Now we need to show that the map $\psi_{\mathcal{N}}$ is well-defined. The only issue here is to check the left triangle of (3.7.1) at a distinguished node.

We may assume that p is a distinguished node joining two components Z_1 and Z_2 with contact order c . We need to check that the map $\psi_{\mathcal{N}}$ defined at the generic points of the two components can be extended to p . Let x_j be the local coordinate of Z_j at p . Denote by $\log x_j$ the corresponding section in both $\mathcal{N}_{C,1}$ and $\mathcal{N}_{C,2}$. Then we automatically have

$$\psi_{\mathcal{N}}(\log x_i) = \log x_i.$$

Without loss of generality, assume that the orientation of the node is given by $Z_1 > Z_2$. Then locally at p , we have

$$(3.7.4) \quad f_1^b(\delta) = e_{Z_2} + c \cdot \log x_2$$

and

$$(3.7.5) \quad f_2^b(\delta) = \psi_{\mathcal{N}}(e_{Z_2}) + c \cdot \log x_2$$

where $\psi_{\mathcal{N}}(e_{Z_2})$ is defined by (3.7.3) at some smooth non-marked point of Z_2 .

On the other hand, we have a section $e_l \in \mathcal{M}_S^{C/S}$ such that

$$e_l = \log x_1 + \log x_2.$$

We identify e_l with the corresponding sections in $\mathcal{N}_{S,i}$ and $\mathcal{N}_{C,i}$ via ϕ_i . Now combining this with (3.7.4) and (3.7.5), and generalizing to a smooth non-marked point of Z_1 , we get

$$f_1^b(\delta) + c \cdot \log x_1 = e_{Z_2} + c \cdot e_l$$

and

$$f_2^b(\delta) + c \cdot \log x_1 = \psi_{\mathcal{N}}(e_{Z_2}) + c \cdot e_l.$$

Since x_1 is invertible away from p , it make sense to write “ $-\log x_1$ ” as above. Hence

$$\psi_{\mathcal{N}}(e_{Z_2} + c \cdot e_l) = \psi_{\mathcal{N}}(e_{Z_2}) + c \cdot e_l.$$

This proves that the definitions of $\psi_{\mathcal{N}}$ on the two components meeting at p are compatible. Therefore the map $\psi_{\mathcal{N}}$ is well-defined. In particular, the above construction gives a canonical isomorphism $\pi^* \mathcal{N}_1 \cong \pi^* \mathcal{N}_2$.

Note that $\pi : C \rightarrow S$ forms a flat cover. Since log structures can be glued under fppf topology [Ols03a], the map $\psi_{\mathcal{N}}$ descends to a well-defined isomorphism of log structures $\mathcal{N}_1 \rightarrow \mathcal{N}_2$, which induces an isomorphism $\xi'_1 \cong \xi'_2$. The uniqueness follows from that of $\psi_{\mathcal{N}}$ in the above construction. \square

Proposition 3.7.5. *There are at most finitely many minimal log stable maps over a geometric point with fixed underlying map and marked graph.*

Proof. Fixing a discrete data Γ , the number of possible choices of contact orders along marked points is finite. It is enough to show that the number of Γ -minimal log stable maps with fixed underlying structure and marked graph is at most finite.

Consider a Γ -minimal log stable map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a geometric point S with the fixed underlying structure $\underline{\xi}$ and marked graph G . Denote by $\xi' = (C \rightarrow S, \mathcal{N}_S, g)$ the coarse log map of ξ over S . Then we have the natural arrow $\xi \rightarrow \xi'$.

On the other hand, consider the saturation map $\mathcal{S} : (S^S, \mathcal{M}) = (S, \mathcal{N}_S)^{\text{Sat}} \rightarrow (S, \mathcal{N}_S)$. Denote by ξ^S the log stable map over S^S obtained by pulling back ξ' via \mathcal{S} . It is easy to check that ξ^S is minimal. By [Ogu06, Chapter II,2.4.5], we have a canonical map $h' : (S, \mathcal{M}_S) \rightarrow (S^S, \mathcal{M})$, such that $h = \mathcal{S} \circ h'$. This induces an arrow of minimal log stable maps $\xi \rightarrow \xi^S$. By comparing the characteristic, it is easy to see that h' is a strict closed immersion.

Since the underlying map of \mathcal{S} is finite, the statement follows from Lemma 3.7.4. \square

3.8. Finiteness of automorphisms. Let $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ be a minimal log stable map over a geometric point S . We fix a chart $\overline{\mathcal{M}}_S \rightarrow \mathcal{M}_S$, and identify the elements e_v and e_l for $v \in V(G_\xi)$ and $l \in E(G_\xi)$ with their images in \mathcal{M}_S .

Proposition 3.8.1. *Notations as above, the set $\mathcal{A}ut_S(\xi)(S)$ is finite.*

Proof. Note that the underlying automorphism of f is finite. Fixing an underlying automorphism (ρ, id_S) , it is enough to show that there are finitely many automorphisms of ξ whose underlying structure are given by (ρ, id_S) . For simplicity, we assume that $\underline{\rho} = id_C$, and other cases can be proved similarly.

Let (ρ, θ) be an automorphism with the underlying structure given by (id_C, id_S) . First we consider a node $l \in E(G_\xi)$. Denote by x and y the local coordinates of l . We can choose x and y so that $e_l = \log x + \log y$. Note that we have

$$(3.8.1) \quad \rho^b(e_l) = \rho^b(\log x) + \rho^b(\log y) = \log \rho^*(x) + \log \rho^*(y) = \log x + \log y.$$

Since $\underline{\rho} = id_C$, the element e_l is fixed by ρ for any l . The same argument shows that the log structure from the marked points are also fixed by ρ .

Now consider a minimal vertex $v \in V(G_\xi)$. Locally on the component of v , we have

$$f^b(\delta) = e_v + \log h,$$

where h is a local invertible section. Note that we have

$$(3.8.2) \quad \rho^b(e_v + \log h) = \rho^b(e_v) + \log \rho^*(h) = \rho^b(e_v) + \log h.$$

Since ρ fixes the section $f^b(\delta)$, the map ρ^b also fixes the element e_v . Thus, the automorphism (ρ, θ) acts trivially on all elements associated to vertices and edges of G_ξ . By Lemma 3.3.5 and 3.3.8, the number of choices of (ρ, θ) is finite. \square

Denote by $\xi' = (C \rightarrow S, \mathcal{N}_S, g)$ the coarse log map of ξ . Then any automorphism $(\rho, \theta) \in \mathcal{A}ut_S(\xi)(S)$ induces a unique automorphism of ξ' . Indeed, the above proof of Proposition 3.8.1 implies the following:

Corollary 3.8.2. *Consider an automorphism $(\rho, \theta) \in \mathcal{A}ut_S(\xi)(S)$. The induced map of sub-log structures $\theta^b : \mathcal{N}_S \rightarrow \mathcal{N}_S$ is uniquely determined by the underlying automorphism (ρ, id_S) . In particular, the automorphism of ξ' induced by (ρ, θ) is uniquely determined by the underlying automorphism (ρ, id_S) .*

Denote by $\underline{\xi}$ the usual stable maps obtained by removing log structures on ξ . We can strengthen the result of Proposition 3.8.1 as follows:

Lemma 3.8.3. *The map of groups $\mathcal{A}ut_S(\xi)(S) \rightarrow \mathcal{A}ut_S(\underline{\xi})(S)$ is an isomorphism.*

Proof. Consider an element $(\rho, id_S) \in \mathcal{A}ut_S(\underline{\xi})(S)$. It is enough to show that there is exactly one element $(\rho, \theta) \in \mathcal{A}ut_S(\xi)(S)$, which is the pre-image of (ρ, id_S) . Consider the following diagram:

$$(3.8.3) \quad \begin{array}{ccc} \mathcal{N}_S & \hookrightarrow & \mathcal{M}_S \\ \downarrow & & \downarrow \theta^b \\ \mathcal{N}_S & \hookrightarrow & \mathcal{M}_S, \end{array}$$

where the left vertical arrow can be constructed similarly by (3.8.1) and (3.8.2), which is uniquely determined by the underlying map (ρ, id_S) . Corollary 3.8.2 implies that any (ρ, θ)

over (ρ, id_S) induces the same map $\mathcal{N}_S \rightarrow \mathcal{N}_S$ as in (3.8.3). Hence to find (ρ, θ) , it is equivalent to find the dashed arrow θ^b , which makes the above diagram (3.8.3) commutative. By the adjointness of saturation and inclusion functors of log structures as in [Ogu06, Chapter II,2.4.5], we have the following commutative diagram:

$$(3.8.4) \quad \begin{array}{ccccc} (S, \mathcal{M}_S) & \xhookrightarrow{i} & (S, \mathcal{N}_S)^{Sat} & \longrightarrow & (S, \mathcal{N}_S) \\ & \searrow \theta' & \downarrow \exists! \cong & & \downarrow \\ & & (S, \mathcal{N}_S)^{Sat} & \longrightarrow & (S, \mathcal{N}_S) \end{array}$$

where $(S, \mathcal{N}_S)^{Sat}$ is the saturation of (S, \mathcal{N}_S) .

Denote by ξ' the coarse log curve of ξ over S . Then by taking the saturation, we obtain a minimal log stable map $(\xi')^S$ over $(S, \mathcal{N}_S)^{Sat}$. Note that the left triangle of (3.8.4) induces a commutative diagram of minimal log stable maps:

$$(3.8.5) \quad \begin{array}{ccc} \xi & \longrightarrow & (\xi')^S \\ & \searrow & \downarrow \\ & & (\xi')^S \end{array}$$

This gives a unique θ^b as in (3.8.3), hence a unique isomorphism of ξ . \square

Proposition 3.8.4. *The natural map $\mathcal{K}_\Gamma(X^{log}) \rightarrow \mathcal{K}_{g,n}(X, \beta)$ by removing log structures from minimal log stable maps is representable.*

Proof. This follows from Lemma 3.8.3 and [LMB00, 8.1.1]. \square

4. THE STACK OF MINIMAL LOG MAPS AS CATEGORY FIBERED OVER \mathbf{LogSch}^{fs}

By the construction in last section, the stacks $\mathcal{K}_{g,n}^{min}(X^{log})$, $\mathcal{K}_{g,n}^{mst}(X^{log}, \beta)$, and $\mathcal{K}_\Gamma(X^{log})$ as open sub-stacks of $\mathcal{LM}_{g,n}(X^{log})$, are fibered categories over \mathbf{Sch} , parameterizing minimal log maps over usual schemes with various numerical conditions. In this section, we give a different categorical explanation as categories fibered over \mathbf{LogSch}^{fs} .

4.1. The universal property of minimal log maps. In this subsection, we fix a log map $\xi = (C \rightarrow S, \mathcal{M}_S, f : (C, \mathcal{M}_C) \rightarrow (X, \mathcal{M}_X))$ such that the log structure \mathcal{M}_S is fs.

Proposition 4.1.1. *There exists a minimal log map over S*

$$\xi_{min} = (C \rightarrow S, \mathcal{M}_S^{min}, f_{min} : (C, \mathcal{M}_C^{min}) \rightarrow (X, \mathcal{M}_X)),$$

and a map of fs log schemes $\Phi : (S, \mathcal{M}_S) \rightarrow (S, \mathcal{M}_S^{min})$, which fits in the following commutative diagram

$$(4.1.1) \quad \begin{array}{ccccc} & & & & (X, \mathcal{M}_X) \\ & & & \nearrow f & \\ (C, \mathcal{M}_C) & \xrightarrow{\Phi_C} & (C, \mathcal{M}_C^{min}) & \xrightarrow{f_{min}} & \\ \downarrow & & \downarrow & & \\ (S, \mathcal{M}_S) & \xrightarrow{\Phi} & (S, \mathcal{M}_S^{min}) & & \end{array}$$

where the square is a ceterian square of log schemes. Furthermore, the datum (Φ, ξ_{min}) is unique up to a unique isomorphism.

Proof. Note that the statement is local on S . Then the proposition follows from Lemmas 4.1.2, 4.1.3, 4.1.4, and 4.1.5 below. \square

By Construction 3.4.1, for each geometric point $\bar{t} \in S$ we can associate a marked graph $G_{\xi_{\bar{t}}}$ to the fiber $\xi_{\bar{t}}$. It was shown in Lemma 3.4.3 that $G_{\xi_{\bar{t}}}$ is admissible. By Proposition 3.4.2, we have a canonical morphism of monoids

$$(4.1.2) \quad \phi_{\bar{t}} : \overline{\mathcal{M}}(G_{\xi_{\bar{t}}}) \rightarrow \overline{\mathcal{M}}_{S, \bar{t}}$$

Lemma 4.1.2. *Assume that we have a log pre-stable curve $(C \rightarrow S, \mathcal{M}_S^{min})$ and a morphism $\Phi : (S, \mathcal{M}_S) \rightarrow (S, \mathcal{M}_S^{min})$ such that*

- (1) *For each point $s \in S$, we have a fixed isomorphism $\overline{\mathcal{M}}_{S, \bar{s}}^{min} \cong \overline{\mathcal{M}}(G_{\xi_{\bar{s}}})$.*
- (2) *The induced map $\bar{\Phi}_{\bar{s}}^b : \overline{\mathcal{M}}(G_{\xi_{\bar{s}}}) \cong \overline{\mathcal{M}}_{S, \bar{s}}^{min} \rightarrow \overline{\mathcal{M}}_{S, \bar{s}}$ on the level of characteristic is identical to canonical map $\phi_{\bar{s}}$ as (4.1.2).*
- (3) *The log pre-stable curve $(C \rightarrow S, \mathcal{M}_S)$ is the pull-back of $(C \rightarrow S, \mathcal{M}_S^{min})$ via Φ .*

Then we have a unique log map $f_{min} : (C, \mathcal{M}_C^{min}) \rightarrow (X, \mathcal{M}_X)$, which fits in diagram 4.1.1. Note that $(C \rightarrow S, \mathcal{M}_S^{min}, f_{min})$ forms a minimal log map over the scheme S .

Proof. Since all the underlying maps are fixed, it is enough to construct a map of log structures $f_{min}^b : f^*(\mathcal{M}_X) \rightarrow \mathcal{M}_C^{min}$, which fits in the following commutative diagram

$$\begin{array}{ccc} & f^*(\mathcal{M}_X) & \\ f_{min}^b \swarrow & & \searrow f^b \\ \mathcal{M}_C^{min} & \xrightarrow{\Phi_C^b} & \mathcal{M}_C \end{array}$$

Consider an arbitrary closed point $p \in C$, which lies in an irreducible component corresponding to the vertex $v \in V(G_{\xi_{\bar{s}}})$. Then locally at p , we have

$$(4.1.3) \quad f^b(\delta) = e_{v,0} + \log h,$$

where $e_{v,0} \in \mathcal{M}_S$ near \bar{s} , and h is a non-zero regular section locally near p . Note that there are two possible cases: if p is a smooth non-marked point, then h is a locally invertible section; if p is a special point with contact order c , then $h = u \cdot \sigma^c$, where u is a locally invertible section, and σ is a local coordinate function vanishing at p . Note that the underlying map Φ_C is an identity. Thus, to define $f_{min}^b(\delta)$ locally at p , it is enough to find a lifting $\tilde{e}_v \in \mathcal{M}_S^{min}$ of $e_{v,0}$, such that the image of \tilde{e}_v in $\overline{\mathcal{M}}_{S, \bar{s}}^{min}$ is the element associated to the vertex v .

We first consider the uniqueness. Assume that we have two lifting \tilde{e}_v and \tilde{e}'_v , such that their images in $\overline{\mathcal{M}}_{S, \bar{s}}^{min}$ are given by the element associated to v . Then, we have $\tilde{e}_v = \log u + \tilde{e}'_v$ for some locally invertible function u . This implies that

$$\Phi_C^b(\tilde{e}_v) = \Phi_C^b(\log u) + \Phi_C^b(\tilde{e}'_v).$$

Since \tilde{e}_v and \tilde{e}'_v are two lifting of $e_{v,0}$, we have $\Phi_C^b(\log u) = 1$. Note that the underlying map $\Phi_C = id_C$. It follows that $u = 1$. This shows that the lifting is unique.

Now we consider the existence of the lifting. Denote by $\bar{e}_{v,0}$ the image of $e_{v,0}$ in the characteristic $\overline{\mathcal{M}}_{S, \bar{s}}$. By (2) the map of monoids $\bar{\Phi}_{\bar{s}}^b$ is identical to $\phi_{\bar{s}}$. Then we have a unique element $\bar{e}_v \in \overline{\mathcal{M}}_{S, \bar{s}}^{min}$, which corresponds to the element associated to v in the graph G , and

$\bar{\Phi}_{\bar{s}}^b(\bar{e}_v) = \bar{e}_{v,0}$. Thus, locally we can lift \bar{e}_v to an element $\tilde{e}_v \in \mathcal{M}_S^{min}$ such that $\Phi_{\bar{s}}^b(\tilde{e}_v) = e_v$. Then we define

$$(4.1.4) \quad f_{min}^b(\delta) = \tilde{e}_v + \log h.$$

The uniqueness of the lifting shows that the construction in (4.1.4) can be glued globally to obtain a unique map f_{min}^b . We can check locally that the map of monoids f_{min}^b is compatible with the structure morphisms of the corresponding log structures. This finishes the proof of the statement. \square

In fact, in the above proof we constructed a log map f_{min} , which is minimal at \bar{s} , hence minimal in a neighborhood of \bar{s} , by the openness of minimality. We next construct a unique log prestable curve $(C \rightarrow S, \mathcal{M}_S^{min})$ satisfying the three conditions in the above lemma. Note that the question is local on S . Pick a point $\bar{s} \in S$. Shrinking S , we can assume that there is a global chart $\beta : \overline{\mathcal{M}}_{S,\bar{s}} \rightarrow \mathcal{M}_S$. We have the canonical map $\phi_{\bar{s}} : \overline{\mathcal{M}}(G_{\xi_{\bar{s}}}) \rightarrow \overline{\mathcal{M}}_{S,\bar{s}}$. Consider the pre-log structure given by the following composition:

$$\overline{\mathcal{M}}(G_{\xi_{\bar{s}}}) \xrightarrow{\phi_{\bar{s}}} \overline{\mathcal{M}}_{S,\bar{s}} \xrightarrow{\beta} \mathcal{M}_S \xrightarrow{\exp} \mathcal{O}_S.$$

Denote by \mathcal{M}_S^{min} the log structure associated to the above pre-log structure. Thus, the construction above gives a global chart $\beta_{min} : \overline{\mathcal{M}}(G_{\xi_{\bar{s}}}) \rightarrow \mathcal{M}_S^{min}$ and a natural map $\Phi^b : \mathcal{M}_S^{min} \rightarrow \mathcal{M}_S$.

Note that the construction of \mathcal{M}_S^{min} depends on the choice of the chart β . Assume that we have another log structure \mathcal{M}_1^{min} and a map $\Phi_1^b : \mathcal{M}_1^{min} \rightarrow \mathcal{M}_S$ over S , which comes from another chart $\beta_1 : \overline{\mathcal{M}}_{S,\bar{s}} \rightarrow \mathcal{M}_S$. Then we have:

Lemma 4.1.3. *There is a unique isomorphism of log structures $\mathcal{M}_1^{min} \rightarrow \mathcal{M}_S^{min}$ fitting in the following commutative diagram:*

$$\begin{array}{ccc} \mathcal{M}_1^{min} & \xrightarrow{\quad} & \mathcal{M}_S^{min} \\ & \searrow \Phi_1^b & \swarrow \Phi^b \\ & \mathcal{M}_S & \end{array}$$

Proof. Consider an irreducible element $a \in \overline{\mathcal{M}}(G)$. Then the construction of \mathcal{M}_1^{min} and \mathcal{M}_S^{min} implies that

$$\Phi_1^b \circ \beta_1(a) + \log u = \Phi^b \circ \beta_{min}(a),$$

where u is a unique invertible section. We define

$$\beta_1(a) \mapsto \beta_{min}(a) + \log u^{-1}.$$

This induces a unique map $\mathcal{M}_1^{min} \rightarrow \mathcal{M}_S^{min}$, which satisfies the condition of the lemma. \square

Lemma 4.1.4. *There exists a unique dashed arrow which fits in the following commutative diagram:*

$$(4.1.5) \quad \begin{array}{ccc} \mathcal{M}_S^{min} & \xrightarrow{\Phi^b} & \mathcal{M}_S \\ & \swarrow \phi_{min} \quad \searrow \phi & \\ & \mathcal{M}_S^{C/S} & \end{array} .$$

where ϕ is the structure arrow defining the log pre-stable curve $(C \rightarrow S, \mathcal{M}_S)$, see Definition B.2.2.

Proof. Further shrinking S , we can choose a global chart $\mathbb{N}^m \rightarrow \mathcal{M}_S^{C/S}$. Let e be a generator of \mathbb{N}^m which corresponds to an edge $l \in V(G_{\xi_{\bar{s}}})$. For convenience, we will identify e with its image in $\mathcal{M}_S^{C/S}$. Consider $\phi(e) \in \mathcal{M}_S$, and its image $\bar{\phi}(e) \in \overline{\mathcal{M}}_S$. Now on the level of characteristic, there is a unique element $\bar{e}' \in \overline{\mathcal{M}}_S^{min}$, which corresponds to the element associated to l , such that $\bar{\Phi}^b(\bar{e}') = \bar{\phi}(e)$. A similar argument as in the proof of Lemma 4.1.2 shows that there is a unique section $e' \in \mathcal{M}_S^{min}$ such that $\Phi^b(e') = \phi(e)$. Then we can define $\phi_{min}(e) = e'$ for every generator e . This gives the map $\phi_{min} : \mathcal{M}_S^{min} \rightarrow \mathcal{M}_S$.

Note that our construction depends on a fixed chart $\mathbb{N}^m \rightarrow \mathcal{M}_S^{C/S}$. However, a similar argument as in the proof of Lemma 4.1.3 shows that different choice of the global chart will induce the same map ϕ_{min} . This finishes the proof. \square

Note that the arrow ϕ_{min} induces a log pre-stable curve $(C \rightarrow S, \mathcal{M}_S^{min})$. Denote by \mathcal{M}_C^{min} the corresponding log structure on C associated to the log curve. By Lemma 3.2.9, we can further shrink S , and assume that the contact order of the nodes on each geometric fiber is given by the weights of the edges of $G_{\xi_{\bar{s}}}$. Now we have:

Lemma 4.1.5. *The log pre-stable curve $(C \rightarrow S, \mathcal{M}_S^{min})$ and the log map $(S, \mathcal{M}_S)^{min} \rightarrow (S, \mathcal{M}_S)$ induced by Φ^b satisfy the condition (1), (2) and (3) in Lemma 4.1.2.*

Proof. Note that (3) follows from the commutativity of (4.1.5) and Definition B.2.2. For (1) and (2), we can repeat the argument in Lemma 4.1.2. Indeed, the same construction there yields a log map $(C, \mathcal{M}_C^{min}) \rightarrow (X, \mathcal{M}_X)$, which is minimal at \bar{s} . Now the openness of minimality implies that all points in a neighborhood of \bar{s} are minimal. Therefore, the property (1) and (2) in Lemma 4.1.2 follows. \square

4.2. Proof of Theorem 1.2.3. By the definition of log stack in Section A.3, the stack $\mathcal{K}_\Gamma(X^{log})$ carries a natural log structure $\mathcal{M}_{\mathcal{K}_\Gamma(X^{log})}$ as follows. For any $g : S \rightarrow \mathcal{K}_\Gamma(X^{log})$, the log structure $g^*\mathcal{M}_{\mathcal{K}_\Gamma(X^{log})}$ is exactly the log structure on S given by the minimal log map ξ over S induced by g . Now we have a universal diagram of log stacks:

$$\begin{array}{ccc} (\mathfrak{C}_\Gamma, \mathcal{M}_{\mathfrak{C}_\Gamma}) & \longrightarrow & X^{log} \\ \downarrow & & \\ (\mathcal{K}_\Gamma(X^{log}), \mathcal{M}_{\mathcal{K}_\Gamma(X^{log})}) & & \end{array}$$

where the pair $(\mathfrak{C}_\Gamma, \mathcal{M}_{\mathfrak{C}_\Gamma})$ is the universal curve over $\mathcal{K}_\Gamma(X^{log})$ with universal log structures $\mathcal{M}_{\mathfrak{C}_\Gamma}$. This diagram gives a log stable map $\xi_{\mathcal{K}_\Gamma(X^{log})}$ over $(\mathcal{K}_\Gamma(X^{log}), \mathcal{M}_{\mathcal{K}_\Gamma(X^{log})})$.

Now consider a log stable map ξ^{log} over (S, \mathcal{M}_S) . Then the tuple $(\xi^{log}, S, \mathcal{M}_S)$ gives a log stable map over S . The universal property of minimal log map implies that there is a unique minimal log map $\xi_{min} = (\xi_{min}^{log}, S, \mathcal{M}_S^{min})$ over S , and a map of log schemes $g : (S, \mathcal{M}_S) \rightarrow (S, \mathcal{M}_S^{min})$, such that $\xi^{log} = g^*\xi_{min}^{log}$ as in Definition 2.1.2. This induces a unique log map $f : (S, \mathcal{M}_S) \rightarrow (\mathcal{K}_\Gamma(X^{log}), \mathcal{M}_{\mathcal{K}_\Gamma(X^{log})})$ such that $\xi^{log} = f^*\xi_{\mathcal{K}_\Gamma(X^{log})}$.

Theorem 1.2.3 follows.

Remark 4.2.1. Using the same argument as above, we can show that the two stacks $\mathcal{K}_{n,g}^{mst}(X^{log}, \beta)$ and $\mathcal{K}_{g,n}^{min}(X^{log})$ with their universal minimal log structures can be viewed as categories fibered over \mathbf{LogSCh}^{fs} , parameterizing log maps over fs log schemes with corresponding numerical conditions.

Remark 4.2.2. If the log structure \mathcal{M}_X on the target X is trivial, the stack $\mathcal{K}_{g,n}^{mst}(X^{log}, \beta)$ is isomorphic to the stack $\mathcal{K}_{g,n}(X, \beta)$ of usual stable maps with the minimal log structure coming from the canonical log structure of its universal curve.

5. THE BOUNDEDNESS THEOREM FOR MINIMAL LOG STABLE MAPS

5.1. Statement of the boundedness theorem. In this section, we fix the target $X^{log} = (X, \mathcal{M}_X)$ as in Convention 3.1.3. The main result of this section is the following:

Theorem 5.1.1. *There exists a scheme T of finite type, and a map $g : T \rightarrow \mathcal{K}_\Gamma(X^{log})$, which exhausts all geometric point of $\mathcal{K}_\Gamma(X^{log})$. Namely, for any point $\xi \in \mathcal{K}_\Gamma(X^{log})(\mathbb{C})$, there exists a lifting $\text{Spec } \mathbb{C} \rightarrow T$, such that its composition with g gives ξ .*

Proof. The proof of this theorem will occupy the whole section. Indeed, we will prove that the map $\mathcal{K}_\Gamma(X^{log}) \rightarrow \mathcal{K}_{g,n}(X, \beta)$, obtained by removing all log structures, is of finite type. In Section 5.3, we will bound the choices of marked graph by stratifying $\mathcal{K}_{g,n}(X, \beta)$. In Section 5.4, we will construct a family of minimal log stable maps, which exhausts all possible minimal log structures with fixed underlying map and marked graph. This will be achieved by considering isomorphisms of corresponding line bundles. The result from Section 5.2 will be used in the above argument. \square

5.2. Isomorphisms of line bundles induced by log stable maps. Consider a log stable map (not necessarily minimal) $\xi = (C \rightarrow S, \mathcal{M}_S, f)$ over a scheme S . In this subsection, we put the following assumption

(5.2.1) The characteristic $\overline{\mathcal{M}}_S$ is a constant sheaf of monoids on S .

Lemma 5.2.1. *With the assumptions as above, the marked graphs of all geometric fibers of ξ are isomorphic.*

Proof. Note that the elements smoothing the distinguished nodes are in $\overline{\mathcal{M}}_S$. Then the statement follows from the assumption (5.2.1). \square

Given the log stable map ξ over S as above, let us consider the following commutative diagram:

$$\begin{array}{ccc} f^*(\mathcal{M}_X) & \xrightarrow{f^\flat} & \mathcal{M}_C \\ \downarrow p_1 & & \downarrow p_2 \\ \mathbb{N} \longrightarrow f^*(\overline{\mathcal{M}}_X) & \xrightarrow{\bar{f}^\flat} & \overline{\mathcal{M}}_C. \end{array}$$

The composition of the bottom arrow $\mathbb{N} \rightarrow \overline{\mathcal{M}}_C$ locally lifts to a chart of a sub-log structure of \mathcal{M}_C . Denote by \mathcal{M} the resulting sub-log structure. Note that this is also a DF-log structure. The map f^\flat induces an isomorphism of log structures $f^*(\mathcal{M}_X) \rightarrow \mathcal{M}$. By the argument in Section A.2, this gives an isomorphism of the corresponding line bundles and sections. Next, we will describe this isomorphism on each irreducible component of C .

Pick a point $\bar{s} \in S$. Shrinking S , we may choose a lifting of global chart $\beta : \overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \mathcal{M}_S$. Consider the induced map $\hat{\beta} : \overline{\mathcal{M}}_S \rightarrow \mathcal{M}_C$. Denote by $\widehat{\mathcal{M}}_C = \mathcal{M}_C^{gp} / (\overline{\mathcal{M}}_S)^{gp}$ the quotient

given by the map $\hat{\beta}$. Consider the following commutative diagram:

$$(5.2.2) \quad \begin{array}{ccccc} 0 & \longrightarrow & (\overline{\mathcal{M}}_S)^{gp} & \xrightarrow{\hat{\beta}^{gp}} & \mathcal{M}_C^{gp} & \longrightarrow & \widehat{\mathcal{M}}_C \\ & & & & \uparrow & \nearrow f^{\flat} & \\ & & & & f^*(\mathcal{M}_X) & & \end{array}$$

where the map f^{\flat} is given by the composition $f^*(\mathcal{M}_X) \rightarrow \mathcal{M}_C \rightarrow \widehat{\mathcal{M}}_C$.

Remark 5.2.2. Note that the morphism f^{\flat} depends on the choice of a lifting $\beta : \overline{\mathcal{M}}_{S,\bar{s}} \rightarrow \mathcal{M}_S$. This will be important when we discuss the valuative criterion.

Conventions 5.2.3. Consider the irreducible component C_v of C corresponding to a vertex $v \in G_\xi$. Note that C_v is connected. Denote by $\{p_l\}_{l \in \Lambda_v^{low}}$ the set of splitting nodes, joining v with v' for some $v' \leq v$. Let $\{p_l\}_{l \in \Lambda_v^{up}}$ be the set consisting of the following special points in C_v :

- (1) the set of splitting nodes, joining v with v'' for some $v \leq v''$;
- (2) the marked points with non-trivial contact orders.

Denote by c_l the contact order at p_l for $l \in \Lambda_v^{low} \cup \Lambda_v^{up}$. Consider the line bundle

$$L_v = \prod_{l \in \Lambda_v^{low}} \mathcal{O}_{C_v}(c_l \cdot p_l) \otimes \prod_{l \in \Lambda_v^{up}} \mathcal{O}_{C_v}(-c_l \cdot p_l).$$

Note that the line bundle L_v only depends on the graph G_ξ .

Proposition 5.2.4. *Assume that the element associated to a vertex $v \in G_\xi$ is not zero. Then the map f^{\flat} induces a natural isomorphism of line bundles*

$$\hat{f}_v^{\flat} : f^*(L)_v \rightarrow L_v.$$

Proof. We first construct \hat{f}_v^{\flat} locally. There are three cases.

Case 1: Consider a closed point p of p_l for $l \in \Lambda_v^{up}$. Locally at p we have

$$f^{\flat}(\delta) = e_v + c_l \log \sigma_l,$$

where σ_l is the local coordinate of p in C_v defining the marking p_l , and e_v is contained in the image of $\hat{\beta}$. Thus, we have $\hat{f}_v^{\flat}(\delta) = c_l \log \sigma_l$. Then locally near p we define

$$(5.2.3) \quad \hat{f}_v^{\flat}(\delta) = \sigma_l^{c_l},$$

Note that $\sigma_l^{c_l}$ is the local section of L_v at p .

Case 2: Consider a closed point p of the splitting node p_l for $l \in \Lambda_v^{low}$. Assume that p_l joining vertices v' and v such that $v' \leq v$. Locally at p we have

$$(5.2.4) \quad f^{\flat}(\delta) = e_{v'} + c_l \log \sigma'_l,$$

where $e_{v'}$ is in the image of $\hat{\beta}$. By a nice choice of coordinates we have

$$(5.2.5) \quad c_l \cdot e_l = c_l \log \sigma_l + c_l \log \sigma'_l, \quad \text{in } \mathcal{M}_C$$

where σ'_l is the local coordinate of p_l in C'_v , and e_l is the element smoothing node, and contained in the image of $\hat{\beta}$. Then we have

$$1 = c_l \log \sigma_l + c_l \log \sigma'_l, \quad \text{in } \widehat{\mathcal{M}}_C.$$

This induces

$$\hat{f}^{\flat}(\delta) = c_l \log \sigma_l = 1 - c_l \log \sigma_l.$$

Then locally at the node p we define

$$(5.2.6) \quad \hat{f}_v^{\flat}(\delta) = \left(\frac{1}{\sigma_l}\right)^{c_l}.$$

Note that this is a local generator of L_v at p .

Case 3: Locally at a point p which is not contained in one of the p_l for $l \in \Lambda_v^{up} \cup \Lambda_v^{low}$, we have

$$f^{\flat}(\delta) = e_v + \log h,$$

where h is an invertible function at p and e_v is contained in the image of $\hat{\beta}$. Then the map $\hat{f}^{\flat}(\delta) = \log h$ induces

$$(5.2.7) \quad \hat{f}_v^{\flat}(\delta_\lambda) = h.$$

Note that the local construction of \hat{f}_v^{\flat} is uniquely determined by \hat{f}^{\flat} , which is a map of sheaves of monoids. Thus these local definitions can be glued to obtain a global map. We also notice that δ lifts to a the local generator of L . Therefore we construct an isomorphism of line bundles \hat{f}_v^{\flat} as required. \square

Remark 5.2.5. The local calculation shows that the isomorphism \hat{f}_v^{\flat} in Proposition 5.2.4 depends on the choice of the chart β .

5.3. Finiteness of the discrete data.

Proposition 5.3.1. *The following set is finite:*

$$\{G \mid G \text{ is the marked graph of some } \xi \in \mathcal{K}_\Gamma(X^{log})(\mathbb{C})\}.$$

Proof. Step 1: Bounding the choices of underlying dual graph. Denote by $\mathcal{K}_{g,n}(X, \beta)$ the Kontsevich moduli space of stable maps, with n -marked points, genus g , and curve class β in X . Note that we have a morphism

$$\mathcal{K}_\Gamma(X^{log}) \rightarrow \mathcal{K}_{g,n}(X, \beta),$$

by removing all log structures. Let $U \rightarrow \mathcal{K}_{g,n}(X, \beta)$ be an affine étale chart. Consider the following cartesian diagram without log structures:

$$\begin{array}{ccc} \mathcal{K}_U & \longrightarrow & \mathcal{K}_\Gamma(X^{log}) \\ \downarrow & & \downarrow \\ U & \longrightarrow & \mathcal{K}_{g,n}(X, \beta). \end{array}$$

Since the stack $\mathcal{K}_{g,n}(X, \beta)$ is of finite type, it is enough to prove that the set of dual graphs corresponding to the geometric point of \mathcal{K}_U is finite. Denote by $C_U \rightarrow U$ the universal curve and $\underline{f}_U : C_U \rightarrow X$ the universal map over U .

Since U is of finite type, it is covered by finitely many strata, where the family of curves over each stratum have a fixed dual graph. We put the reduced scheme structure on each stratum. Let S be the stratum corresponding to a graph G . Denote by $\underline{f} : C \rightarrow X$ the universal map over S .

Step 2: Bounding the choices of distinguished nodes and orientations. Since G is a finite graph, the number of choices of distinguished nodes is finite. We first fix a choice of distinguished nodes on G . So we fix an orientation on G such that:

- (1) If C_v does not degenerate to D , then $v \in V_n(G)$
- (2) The non-oriented edges are in one-to-one correspondence to the non-distinguished nodes.
- (3) No cycles contain distinguished edges.

Step 3: Bounding the choices of contact orders. Since we fixed the orientation and distinguished edges on G , we can use the notations $\{p_l\}_{l \in \Lambda_v^{low}}$ and $\{p_l\}_{l \in \Lambda_v^{up}}$ for the two sets of distinguished points on the subcurve C_v as in Conventions 5.2.3. Denote by c_l the possible contact order at the distinguished point p_l . Since the dual graph of the underlying curve is fixed, the multi-degree of $\underline{f}^*(L)$ on C_v is fixed for any $v \in V(G)$. By Proposition 5.2.4, we have

$$(5.3.1) \quad \deg \underline{f}^*(L)|_{C_v} = \sum_{l \in \Lambda_v^{low}} c_l - \sum_{l' \in \Lambda_v^{up}} c_{l'}.$$

First, consider a maximal vertex $v \in V(G)$. Then the set $\{p_l\}_{l \in \Lambda_v^{up}}$ is given by the discrete data Γ . Since the contact orders are all positive, the choices of c_l for $l \in \Lambda_v^{low}$ is finite by (5.3.1).

Consider an arbitrary vertex $v' \in V(G)$. We assume that for any adjacent vertex v of v' such that $v' \leq v$, the number of choices of the contact orders along the splitting nodes joining v' and v is finite. Then by taking into account all contact orders from adjacent vertices and those from marked points of C , a similar argument shows that the possible choices of contact order c_l for $l \in \Lambda_{v'}^{low}$ are also finite in number. Since G is a finite graph, this proves that the choice of contact orders on G is finite.

This finishes the proof of the proposition. \square

5.4. Proof of Theorem 5.1.1. Consider the family of usual stable maps $\underline{f} : C \rightarrow X$ over S as in Step 1 of the above proof. Fix a possible marked graph G_0 with $\overline{G}_0 = G$ the dual graph of C . We use the notations as in Step 3 of the above proof, and assume that (5.3.1) holds for any $v \in V(G_0)$. Since the stack $\mathcal{K}_{g,n}(X, \beta)$ is of finite type, to prove Theorem 5.1.1, it is enough to prove the following:

Proposition 5.4.1. *Notations and assumptions as above, there exists a scheme T of finite type over S , and a family of minimal log stable maps ξ over T , which satisfies the following conditions: for any minimal log stable map ξ' over \bar{s} , with marked graph given by G_0 , and underlying map $\underline{\xi}'$ given by the pull-back of \underline{f} via $\bar{s} \rightarrow S$, there exists a lifting $\bar{s} \rightarrow T$, such that ξ' is isomorphic to the pull-back $\xi_{\bar{s}}$.*

Proof. By shrinking S , we can assume that S is affine, and the canonical log structure $\mathcal{M}_S^{C/S}$ on S coming from the family $C \rightarrow S$ has a global chart $\mathbb{N}^n \rightarrow \overline{\mathcal{M}}_{S, \bar{s}}^{C/S}$ for some geometric point $\bar{s} \in S$. Consider the pre-log structure $\overline{\mathcal{M}}(G_0) \rightarrow \mathcal{O}_S$, given by $e \mapsto 0$ for any non-trivial element $e \in \overline{\mathcal{M}}(G_0)$. Denote by \mathcal{M}_S the new log structure associated to the pre-log structure. Note that there is a map $\mathbb{N}^n \rightarrow \overline{\mathcal{M}}(G_0)$ given by the corresponding nodes. This induces a map $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S$, hence a log pre-stable curve $\zeta = (C \rightarrow S, \mathcal{M}_S)$ over S . Note that any minimal log map ξ' over $\bar{s} \in S$ as in the statement has the source log curve isomorphic to $\zeta_{\bar{s}}$.

Denote by \mathcal{M}_C the log structure on C corresponding to the log pre-stable curve ζ . Note that over C we have another log structure $\underline{f}^*(\mathcal{M}_X)$. Since the dual graph G_0 is fixed, we have a morphism of sheaves of monoids on \overline{C} :

$$\overline{f}^b : \underline{f}^*(\overline{\mathcal{M}}_X) \rightarrow \overline{\mathcal{M}}_C,$$

which is locally described as in Section 3.2. To define a log map $f : (C, \mathcal{M}_C) \rightarrow X^{\log}$, it is enough to define a map of log structures $f^\flat : \underline{f}^*(\mathcal{M}_X) \rightarrow \mathcal{M}_C$ fitting in the following commutative diagram:

$$(5.4.1) \quad \begin{array}{ccc} & & \mathcal{O}_C \\ & \nearrow & \uparrow \\ \underline{f}^*(\mathcal{M}_X) & \overset{f^\flat}{\dashrightarrow} & \mathcal{M}_C \\ \downarrow p_1 & & \downarrow p_2 \\ \mathbb{N} \longrightarrow \underline{f}^*(\overline{\mathcal{M}}_X) & \xrightarrow{\bar{f}^\flat} & \overline{\mathcal{M}}_C \end{array} ,$$

where the two vertical arrows are the canonical projection, and the arrow $\mathbb{N} \rightarrow \underline{f}^*(\overline{\mathcal{M}}_X)$ is the pull-back of the global presentation. Note that the arrow \bar{f}^\flat is injective. Denote by δ_X and δ_C the image of δ in $\underline{f}^*(\overline{\mathcal{M}}_X)$ and $\overline{\mathcal{M}}_C$ respectively. The inverse images $p_1^{-1}(\delta_X)$ and $p_2^{-1}(\delta_C)$ form two \mathcal{O}_C^* -torsors. Denote by $\mathcal{I}som_C(p_1^{-1}(\delta_X), p_2^{-1}(\delta_C))$ the presheaf of isomorphisms of the two torsors over C . To find a dashed arrow as in (5.4.1) is equivalent to have a global section of $\mathcal{I}som_C(p_1^{-1}(\delta_X), p_2^{-1}(\delta_C))$. Note that the torsor $p_1^{-1}(\delta_X)$ corresponds to the line bundle \underline{f}^*L . Denote by L_C the corresponding line bundle of $p_2^{-1}(\delta_C)$. Then we have

$$\mathcal{I}som_C(p_1^{-1}(\delta_X), p_2^{-1}(\delta_C)) \cong \mathcal{I}som_C(\underline{f}^*L, L_C) \cong \mathcal{I}som_C(\underline{f}^*L \otimes L_C^{-1}, \mathcal{O}_C).$$

Denote by I the above presheaf. It is well-known that line bundles are parametrized by the algebraic stack $\mathcal{B}\mathbb{G}_m$. Thus, I is represented by a \mathbb{G}_m -torsor, which is a separated algebraic space of finite type. Let $\pi : C \rightarrow S$ be the projection. By [Ols06, Theorem 1.5], there is an algebraic space π_*I locally of finite type over S , which for any $Y \rightarrow S$ associates the groupoid of isomorphisms $(\underline{f}^*L \otimes L_C^{-1})_Y \rightarrow \mathcal{O}_{C_Y}$. We have the following lemma for the boundedness of π_*I . \square

Lemma 5.4.2. *The algebraic space π_*I is of finite type over S .*

Proof. By our assumption on G_0 , the two line bundles L_C and \underline{f}^*L have the same degree when restrict to each irreducible component over $\bar{s} \in S$. Since S is affine, by [FP97, Proposition 1], there is a unique closed subscheme $T \subset S$ which represents the condition that the line bundle $\underline{f}^*L \otimes L_C^{-1}$ is trivial. The same proof shows that over that locus the line bundle is pulled back from the base. Its sheaf of trivializations is again represented by a \mathbb{G}_m -torsor $U \rightarrow \pi_*I$, which is of finite type. \square

By pulling back via $\pi_*I \rightarrow S$, we have a family of log pre-stable curves $\zeta_{\pi_*I} = (C_I \rightarrow \pi_*I, \mathcal{M}_{\pi_*I})$, a usual stable map $f_{\pi_*I} : C_I \rightarrow X$, and a morphism of sheaves of monoids $f_{\pi_*I}^\flat : \underline{f}_{\pi_*I}^*\mathcal{M}_X \rightarrow \mathcal{M}_{C_I}$, where \mathcal{M}_{C_I} is the log structure on C_I given by the log curve ζ_{π_*I} .

Lemma 5.4.3. *The set of points $t \in \pi_*I$, whose fiber $f_{\pi_*I, \bar{t}}^\flat$ gives a morphism of log structures, forms a closed subset of π_*I .*

Proof. The condition that $f_{\pi_* I}^b$ is a morphism of log structures is equivalent to having the following commutative diagram:

$$(5.4.2) \quad \begin{array}{ccc} \underline{f}_{\pi_* I}^* \mathcal{M}_X & \xrightarrow{f_{\pi_* I}^b} & \mathcal{M}_{C_I} \\ \searrow \text{exp}_X & & \swarrow \text{exp}_C \\ & \mathcal{O}_{C_I} & \end{array}$$

where the two arrows exp_X and exp_C are the structure maps of the corresponding log structures $\underline{f}_{\pi_* I}^* \mathcal{M}_X$ and \mathcal{M}_{C_I} . Locally on C_I , we choose a generator $\delta \in \underline{f}_{\pi_* I}^* \mathcal{M}_X$, then the commutativity of the diagram is equivalent to the following equality of sections of \mathcal{O}_{C_I} :

$$\text{exp}_X(\delta) = \text{exp}_C \circ f_{\pi_* I}^b(\delta),$$

which is a closed condition. Let $V \subset C_I$ be the closed sub-scheme representing the commutativity of (5.4.2) over C_I , and V^c the complement of V in C_I . Denote by W the image of V^c in $\pi_* I$ via the projection $C_I \rightarrow \pi_* I$. Since the family of curves is flat, the image W is open in $\pi_* I$. Thus, the complement W^c of W is closed in $\pi_* I$. This proves the lemma. \square

We take $T = W^c$ as in the above proof with the reduced scheme structure. Then T is a closed subspace of $\pi_* I$. By pulling back families from $\pi_* I$, we obtain a family of minimal log stable maps ξ over T . According to our construction, this family ξ over T satisfies the lifting property in Proposition 5.4.1.

Theorem 5.1.1 follows from the above arguments.

6. THE WEAK VALUATIVE CRITERION FOR MINIMAL LOG STABLE MAPS

6.1. Statement of the weak valuative criterion. Let R be a discrete valuation ring, and K the fraction field of R . Denote by π the uniformizer of R , and write $S = \text{Spec } R$. Let s and η be the closed and generic point of S respectively. If R' is another discrete valuation ring, we will write π' for its uniformizer. Denote by s' and η' the closed and generic point of $S' = \text{Spec } R'$ respectively.

Theorem 6.1.1. *With the notations above, consider a minimal log stable map ξ_η over η . Possibly after a base change given by an injection $R \hookrightarrow R'$ of DVR, which induce a finite extension of fraction fields, we have an extension of minimal log stable maps given by the following cartesian diagram:*

$$\begin{array}{ccc} \xi_{\eta'} & \longrightarrow & \xi_{S'} \\ \downarrow & & \downarrow \\ \eta' & \longrightarrow & S', \end{array}$$

where $\xi_{\eta'}$ is the pull-back of ξ_η via $\eta' \rightarrow \eta$, and $\xi_{S'}$ is a minimal log stable map over S' . Furthermore, the extension $\xi_{S'}$ is unique up to a unique isomorphism and its formation commutes with further injections of discrete valuation rings.

Proof. We first assume that ξ_η is a minimal log map over η , which is not necessarily stable. Possibly after base change, we fix an extension of the underlying pre-stable map $\underline{f} : C \rightarrow S$, such that its restriction to the generic fiber is given by the pull-back of \underline{f}_η . Denote by $\underline{\xi}$ the extended underlying map. Here for simplicity, we still use S to denote the new base. The

existence of compatible minimal log structures on $\underline{\xi}$ will be proved in Section 6.3. This will be achieved by constructing an extension of certain simplified log maps, and using the universal property of minimal log maps. The uniqueness of the extended minimal log structure on $\underline{\xi}$ will be proved in Section 6.5.

In case of stable maps, the extended underlying map $\underline{\xi}$ is unique up to a unique isomorphism. Hence the theorem will be proved by the above argument. \square

Remark 6.1.2. By Observation 2.1.9, there is a map $\mathcal{K}_{g,n}^{\min}(X^{\log}, \beta) \rightarrow \mathcal{K}_{g,n}^{\text{pre}}(X, \beta)$, where $\mathcal{K}_{g,n}^{\text{pre}}(X, \beta)$ is the stack of usual pre-stable maps, and $\mathcal{K}_{g,n}^{\min}(X^{\log}, \beta)$ is as in Definition 3.5.3. Then our proof implies that this map of stacks satisfies the weak valuative criterion.

6.2. Local analysis of the extended underlying map. Let $\xi_\eta = (C_\eta \rightarrow \eta, \mathcal{M}_\eta, f_\eta)$. We first consider the case ξ_η is a log map (not necessarily minimal and stable). We still use $\underline{f} : C \rightarrow X$ to denote the extended underlying map over S . Possibly after a base change, we fix a chart $\beta_\eta : \overline{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$. Denote by G_η the marked graph of ξ_η . If a node of C is smoothed out over η , then we call it a *special node*, otherwise we call it a *generic node*.

Consider a point $p \in C_{\bar{s}}$, and choose an étale neighborhood U of p . Write $U_\eta := U \times_S \eta$. Shrinking U , we have

$$(6.2.1) \quad f_\eta^\flat(\delta) = \beta_\eta(e_v) + \log u_p, \text{ over } U_\eta$$

where $e_v \in \overline{\mathcal{M}}_\eta$ is the degeneracy of some vertex $v \in G_\eta$, and $u_p \in \mathcal{O}_{U_\eta}$.

Assume that p is not a generic node. Shrinking U further, we can assume that U is connected, and does not contain a generic node. We also assume that U does not contain points on other components which are not containing p . Note that in this case U is normal, and u_p extends to a rational function on U . Denote by ν_π the evaluation of the divisor given by the uniformizer π . Let $n_p = \nu_\pi(u_p)$, then we have the following result.

Lemma 6.2.1. *With the above assumption, the point p satisfies one of the following possibilities:*

- (1) *If p is a non-distinguished point, then there is a neighborhood of p , which contains only non-distinguished points over η , and we have $u_p = \pi^{n_p} \cdot h_p$, where $h_p \in \mathcal{O}_U^*$.*
- (2) *If p is a marked point with contact order c over η , then $u_p = \pi^{n_p} \cdot x^c \cdot h_p$, where $h_p \in \mathcal{O}_U^*$, and the section containing p is given by the vanishing of $x \in \mathcal{O}_U$.*
- (3) *If p is a special node, then $u_p = \pi^{n_p} \cdot x^c \cdot h_p$, where $h_p \in \mathcal{O}_U^*$, the section $x \in \mathcal{O}_U$ is a local coordinate of one component at p , and c is a non-negative integer.*

Note that if in (3) we have $c = 0$, then this is compatible with the case described in (1).

Proof. Since $n_p = \nu_\pi(u_p)$, and u_p is well-defined over the generic fiber, we have $u_p = \pi^{n_p} \cdot h'_p$ for some $h'_p \in \mathcal{O}_U$. Since p is non-distinguished, there is a neighborhood of p , which contains only non-distinguished points over η . It follows that $h'_p \in \mathcal{O}_{U_\eta}^*$. Since $\nu_\pi(h'_p) = 0$, then $h'_p \in \mathcal{O}_U^*$. This proves (1).

For (2), we have $\nu_x(h'_p) = c$, where ν_x is the evaluation map given by the divisor corresponding to the vanishing of x . Then we have $h'_p = x^c \cdot h_p$, such that the restriction $h_p|_{U_\eta}$ is invertible. The same argument as for (1) shows that $h_p \in \mathcal{O}_U^*$.

Consider the case where p is a special node. Let x and y be local coordinates of the two components meeting at p . Choosing the coordinates appropriately, we may assume that $x \cdot y = \pi^n$ for some positive integer n . Without loss of generality, we can assume

that $\nu_y(h'_p) = 0$ and $\nu_x(h'_p) = c$ for some non-negative integer c . Thus, as in (2), we have $h'_p = x^c \cdot h_p$ for some $h_p \in \mathcal{O}_U^*$. This proves (3). \square

Observation 6.2.2. For a smooth point p , the integer n_p and the rational section u_p in (6.2.1) depend on the choice of the chart β_η . We call the integer n_p *the special degeneracy of p with respect to the chart β_η* . Let Z be the irreducible component of the fiber containing p . Then it is not hard to see that generic points on Z also have n_p as the special degeneracy under β_η . Thus, we call n_p *the special degeneracy of Z under β_η* .

Remark 6.2.3. Consider a node p joining two irreducible components Z_1 and Z_2 over \bar{s} . First we assume that p is a special node. Let x and y be local coordinates on Z_1 and Z_2 at p respectively, such that $x \cdot y = \pi^n$. By Lemma 6.2.1(3), we can assume that $u_p = \pi^{n_p} \cdot x^c \cdot h_p$. Thus, we can check that the special degeneracy of Z_1 is n_p , and the special degeneracy of Z_2 is $n_p + c \cdot n$. In this case, we write $Z_1 \leq Z_2$. Note that if $c = 0$, we have both $Z_1 \leq Z_2$, and $Z_2 \leq Z_1$.

Consider the case where p is a generic node. We take the normalization of C along all the generic node. Then we obtain a set of usual pre-stable curves $\{C_v\}_{v \in V(G_\eta)}$ over S . If $Z_1 \subset C_{v_1}$ and $Z_2 \subset C_{v_2}$, and $v_1 \leq v_2$, then we define $Z_1 \leq Z_2$. We thus define an orientation on the dual graph G of the curve $C_{\bar{s}}$.

The following result, which gives a way of comparing sections in the base log structure, is crucial in the proof of the uniqueness of the extension.

Lemma 6.2.4. *Using the notations as above, consider another chart $\beta'_\eta : \overline{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$, and a generic point $p \in C_{\bar{s}}$ lies in the component corresponding to $v \in V(G_\eta)$. The two special degeneracy of p given by β_η and β'_η are the same, if and only if $\beta'_\eta(e_v) = \log u + \beta_\eta(e_v)$ for some $u \in R^*$.*

Proof. The “if” part is obvious. Consider the other direction. As in (6.2.1), locally at a non-marked point p we have

$$f_\eta^\flat(\delta) = \beta'_\eta(e_v) + \log u'_p = \beta_\eta(e_v) + \log u_p.$$

Then the assumption implies that $u \cdot u'_p = u_p$, for some $u \in R^*$. Thus, we have

$$\beta'_\eta(e_v) = \beta_\eta(e_v) + \log u,$$

which proves the statement. \square

Lemma 6.2.5. *With the notations as above, the integer c as in (2) and (3) of Lemma 6.2.1 does not depend on the choice of chart β_η . Therefore the orientation on G defined in Remark 6.2.3 does not depend on the choice of the chart β_η .*

Proof. Consider another chart $\beta_\eta : \overline{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$, and

$$f_\eta^\flat(\delta) = \beta'_\eta(e_v) + \log u'_p,$$

for some $u'_p \in \mathcal{O}_{U_\eta}^*$. Then we have

$$\beta'_\eta(e_v) = \beta_\eta(e_v) + \log a,$$

for some element $a \in K$. Comparing with (6.2.1), we have

$$u_p = a \cdot u'_p.$$

Since c is given by the evaluation ν_x , this implies the statement of the lemma. \square

We take the normalization of C along all generic nodes. For each $v \in V(G_\eta)$, denote by C_v the corresponding connected component. Now we consider the case where p is a generic node. Let $l \in E(G)$ be the edge corresponding to the generic node p , and assume that l connects two vertices v_1 and v_2 . Denote by p_η the corresponding node over the generic point. Again we have the section u_p over U_η as in (6.2.1). By shrinking U , we can choose two regular sections x and y on U_η , which correspond to the coordinates of the two components meeting at p_η in C_η . Choosing the coordinates appropriately, we may assume

$$(6.2.2) \quad \beta_\eta(e_l) = \log x + \log y, \quad \text{in } U_\eta.$$

Without loss of generality, we can assume that $u_p = x^c$, where c is the contact order at p_η . Then u_p vanishes along the component with coordinate y .

By taking the normalization of U along the generic node given by l , we obtain two subschemes U_1 and U_2 of U . By shrinking U , we can assume that $U_i \subset C_{v_i}$ for $i = 1, 2$. We still use x and y to denote restriction of x and y to U_1 and U_2 respectively, and p_i the pre-image of p in U_i for $i = 1, 2$. Then x and y can be viewed as a rational function on U_1 and U_2 respectively. Let Σ_1 and Σ_2 be the two sections in U_1 and U_2 respectively, coming from the splitting node l . Let σ_i be the regular functions on U_i , whose vanishing gives the section Σ_i for $i = 1, 2$.

Lemma 6.2.6. *With the notations as above, locally at p_i , we have*

- (1) $x = \pi^{n_1} \cdot \sigma_1 \cdot h_1$, where $n_1 = \nu_\pi(x)$, and $h_1 \in \mathcal{O}_{U_1}^*$.
- (2) $y = \pi^{n_2} \cdot \sigma_2 \cdot h_2$, where $n_2 = \nu_\pi(y)$, and $h_2 \in \mathcal{O}_{U_2}^*$.

Proof. The proof of this is similar to that for Lemma 6.2.1. □

Remark 6.2.7. Note that σ_1 and σ_2 form the coordinates at p . Choosing coordinates appropriately, we may assume that $h_1 = h_2 = 1$ in Lemma 6.2.6. Thus, we have $u_p = \pi^{c \cdot n_1} \cdot \sigma_1^c$.

6.3. Existence of the extension. Now we consider the minimal log map ξ_η and the extended underlying map ξ . Denote by $C(\overline{\mathcal{M}}_\eta)$ the convex rational polyhedral cone of $\overline{\mathcal{M}}_\eta$ in $\overline{\mathcal{M}}_\eta^{gp} \otimes_{\mathbb{Z}} \mathbb{Q}$. Since $\overline{\mathcal{M}}_\eta$ is sharp, the cone $C(\overline{\mathcal{M}}_\eta)$ is strongly convex.

Lemma 6.3.1. *There is a lattice point $\tilde{v} \in \overline{\mathcal{M}}_\eta^{gp}$ such that $(u, \tilde{v}) > 0$ for any non-zero element $u \in C(\overline{\mathcal{M}}_\eta)$, where (\cdot, \cdot) is the standard pairing in the Euclidean space $\overline{\mathcal{M}}_\eta^{gp} \otimes_{\mathbb{Z}} \mathbb{Q}$.*

Proof. This follows from [Ful93, Section 1.2(iv)]. □

We fix a lattice point \tilde{v} satisfies the condition in the above lemma. The set

$$\{(u, \tilde{v}) \mid u \in C(\overline{\mathcal{M}}_\eta)\} \subset \mathbb{Q}$$

forms a monoid, whose saturation is the rank one free monoid \mathbb{N} . Thus, we have a map of saturated monoids $l_{\tilde{v}} : \overline{\mathcal{M}}_\eta \rightarrow \mathbb{N}$. Consider the log structure \mathcal{M}'_η , associated to the pre-log structure $\mathbb{N} \rightarrow K$, $e \mapsto 0$ over η . We fix a chart $\beta_\eta : \overline{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$ and $\beta'_\eta : \overline{\mathcal{M}}'_\eta \cong \mathbb{N} \rightarrow \mathcal{M}'_\eta$. Then we have a morphism of log structures $\mathcal{M}_\eta \rightarrow \mathcal{M}'_\eta$ given by

$$\beta_\eta(e) \mapsto \beta'_\eta \circ l_{\tilde{v}}(e).$$

Denote by $\xi'_\eta = (C \rightarrow S, \mathcal{M}'_\eta, f'_\eta)$ the log map obtained by pulling back ξ_η via the map $(\eta, \mathcal{M}_\eta) \rightarrow (\eta, \mathcal{M}'_\eta)$. By Proposition 4.1.1, it is enough to construct a log map (not necessarily minimal) ξ' , such that its generic fiber is given by ξ'_η as above.

Lemma 6.3.2. *Using the notations as above, there exists a chart $\beta'_\eta : \overline{\mathcal{M}}'_\eta \rightarrow \mathcal{M}'_\eta$, such that no components of C over \bar{s} have negative special degeneracy under β'_η as in Observation 6.2.2.*

Proof. We fix an arbitrary chart β'_η as above. Consider an irreducible component Z over the closed point \bar{s} . Let $p \in Z$ be a smooth non-distinguished point $p \in Z$. Consider the nearby points of p over η . By Lemma 6.2.1, we have

$$(6.3.1) \quad (f'_\eta)^\flat(\delta) = \beta'_\eta(e) + \log \pi^n \cdot u,$$

where u is a locally invertible section near p , and $e \in \overline{\mathcal{M}}'_\eta$. If $n \geq 0$, then there is nothing to prove. Consider the case $n < 0$. Since the number of irreducible components over \bar{s} is finite, we can assume that n is minimal among the special degeneracy of all irreducible components of the closed fiber under β'_η . Consider the new chart given by

$$(6.3.2) \quad \beta''_\eta : \overline{\mathcal{M}}'_\eta \rightarrow \mathcal{M}_\eta, \quad e \mapsto \beta'_\eta(e) - n \cdot \log \pi.$$

It is not hard to check that (6.3.1) becomes

$$(f'_\eta)^\flat(\delta) = \beta''_\eta(e) + \log u.$$

Since n is minimal, by applying (6.3.1) and (6.3.2) to other components, it follows that no irreducible component of C over \bar{s} has negative special degeneracy under β''_η . \square

We fix a chart $\beta'_\eta : \overline{\mathcal{M}}'_\eta \rightarrow \mathcal{M}_\eta$, which satisfies the condition in Lemma 6.3.2. Consider the log structure \mathcal{M}'_S associated to the following pre-log structure on S :

$$\mathbb{N}^2 \rightarrow R, \quad e_\eta \mapsto 0, \text{ and } e_s \mapsto \pi,$$

where e_η and e_s form the basis of \mathbb{N}^2 . Now we identify $\mathcal{M}'_{S,\eta}$ with \mathcal{M}'_η , and the element e_η corresponds to the chart $\beta'_\eta : \overline{\mathcal{M}}'_\eta \rightarrow \mathcal{M}_\eta$.

Lemma 6.3.3. *With the notations as above, there is a morphism of log structures $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}'_S$, whose restriction to the generic point η is identical to the morphism of log structures $\mathcal{M}_\eta^{C_\eta/\eta} \rightarrow \mathcal{M}'_\eta$ given by ξ'_η .*

Proof. Possibly after a base change, we can choose a global chart $\beta^{C/S} : \overline{\mathcal{M}}_{S,\bar{s}}^{C/S} \rightarrow \mathcal{M}_S^{C/S}$. Denote by G the dual graph of $C_{\bar{s}}$, and $\{e_l\}_{l \in E(G)}$ the set of generators of $\overline{\mathcal{M}}_S$, such that $\beta^{C/S}(e_l)$ is an element in $\mathcal{M}_S^{C/S}$ smoothing the node corresponding to l in the closed fiber. Assume that l is smoothed out over η , then $\exp \circ \beta^{C/S}(e_l) = \pi^n \cdot h$, where n is a positive integer, and h is an invertible element in R . Thus, we define

$$e_l \mapsto n \cdot e_s + \log h.$$

If the node corresponding to l persists over η , then we have

$$e_l \mapsto n_\eta \cdot e_\eta + \log \pi^{n_s} + \log h, \quad \text{over } \eta$$

where n_η and n_s are two integers, and h is an invertible element in R . Note that n_η is positive, and we may assume that n_s is also positive by choosing a sufficiently large n in (6.3.2). Thus, we define

$$e_l \mapsto n_\eta \cdot e_\eta + n_s \cdot e_s + \log h.$$

This induces a map $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}'_S$, whose restriction to the generic point coincides with $\mathcal{M}_\eta^{C_\eta/\eta} \rightarrow \mathcal{M}'_\eta$. \square

Note that the map $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}'_S$ in the above lemma gives a log pre-stable curves $(C \rightarrow S, \mathcal{M}'_S)$, whose restriction to η is given by the log-prestable curve $(C_\eta \rightarrow \eta, \mathcal{M}'_\eta)$ of ξ'_η .

Proposition 6.3.4. *There is a log map ξ' over (S, \mathcal{M}'_S) with the log curve $(C \rightarrow S, \mathcal{M}'_S)$ and underlying map $\underline{\xi}$, whose restriction to η is identical to ξ'_η .*

Proof. It is enough to define the morphism of log structures $f^\flat : \underline{f}^* \mathcal{M}_X \rightarrow \mathcal{M}'_C$, where \mathcal{M}'_C is the log structure on C corresponding to the log curve $(C \rightarrow S, \mathcal{M}'_S)$. Pick a point $p \in C$ over \bar{s} , and an étale neighborhood U of p . By shrinking U , we can assume that over the generic point, we have

$$f^\flat_\eta(\delta) = n \cdot e_\eta + \log u_p, \quad \text{in } U_\eta,$$

where $u_p \in \mathcal{O}_{U_\eta}$.

We first assume that p is not a generic node. By Lemma 6.2.1, further shrinking U if necessary, the section u_p extend to U of the following form:

$$u_p = \pi^{n_1} \cdot h',$$

where $n_1 = \nu_\pi(u_p)$, and $h' \in \mathcal{O}_U$. Note that the Lemma 6.3.2 implies that the integer n_1 is non-negative. Thus, the only possible way to define f^\flat near p is given by

$$f^\flat(\delta) = n \cdot e_\eta + n_1 \cdot e_s + \log h'.$$

Next we consider the case p is a generic node. With the notations in Remark 6.2.7, we have

$$u_p = \pi^{c \cdot n} \cdot \sigma_1^c.$$

Thus, we define

$$f^\flat(\delta) = n \cdot e_\eta + n_1 \cdot e_s + c \cdot \log \sigma_1.$$

Note that our local construction is obtained by specializing the section u_p to the closed fiber. Since the underlying structure is fixed, such specialization is unique. Thus, the above construction can be glued together to obtain a global map f^\flat as we want. \square

6.4. Specializing the dual graph. Consider the dual graph \underline{G} of the underlying curve $C_{\bar{s}}$ of the fixed extension $\underline{\xi}$. For each edge $l \in E(\underline{G})$, if l corresponds to a special node, then we can associate to l a non-negative integer c given by Lemma 6.2.1; if l corresponds to a generic node, then we associate to l the contact order given by ξ_η . Denote by $V_n(\underline{G})$ the set of non-degenerate components of $C_{\bar{s}}$. Note that Remark 6.2.3 gives an orientation on \underline{G} , which is compatible with the contact orders defined on each edge and the subset $V_n(\underline{G}) \subset V(\underline{G})$. Thus, we obtain a marked graph. We use G to denote this graph with the discrete data.

Proposition 6.4.1. *Consider any minimal log map ξ over S with the fixed underlying map $\underline{\xi}$, which is an extension of ξ_η . Then the marked graph $G_{\xi_{\bar{s}}}$ is identical to the graph G with the orientation and contact orders as above.*

Proof. First note that the underlying graph of $G_{\xi_{\bar{s}}}$ and G are both given by the dual graph \underline{G} of the underlying curve, and their sets of non-degenerate vertices are the same. It is enough to check that the two graphs have the same contact orders and orientations. We denote the underlying graph to be \underline{G} . Consider an edge $l \in E(\underline{G})$. If l corresponds to a generic node, then by Lemma 3.2.9, the orientation and contact order of l is uniquely determined by the generic fiber ξ_η . Hence the two graphs $G_{\xi_{\bar{s}}}$ and G has the same orientation and contact order along l .

Next, consider the case where l corresponds to a special node p . Assume that the contact order of ξ at p is c . Note that the log structure is compatible with the underlying structure. Hence the two graphs have the same weight c in Lemma 6.2.1(3) associated to l , which only depends on f_η . The orientation of l in $G_{\xi_{\bar{s}}}$ is given by the one described in Remark 6.2.3. This implies that the two graphs $G_{\xi_{\bar{s}}}$ and G have the same orientation and contact order along l .

This finishes the proof of the statement. \square

Corollary 6.4.2. *The graph G is admissible.*

Proof. This follows from the existence of the extension of ξ_η and the above proposition. \square

Consider a minimal log map $\xi = (C \rightarrow S, \mathcal{M}_S, f)$, which is an extension of ξ_η over S with underlying map $\underline{\xi}$. Consider the natural map $q^{gen} : \overline{\mathcal{M}}(G)^{gp} \cong \overline{\mathcal{M}}_{S, \bar{s}}^{gp} \rightarrow \overline{\mathcal{M}}_\eta^{gp}$. This is a surjection. Denote by K_{sp} the kernel of q^{gen} . Then we have the following exact sequence:

$$(6.4.1) \quad 0 \longrightarrow K_{sp} \longrightarrow \overline{\mathcal{M}}(G)^{gp} \xrightarrow{q^{gen}} \overline{\mathcal{M}}_\eta^{gp} \longrightarrow 0.$$

Note that all groups involved in the exact sequence (6.4.1) are free abelian groups. We fix a decomposition, which is compatible with (6.4.1):

$$(6.4.2) \quad \overline{\mathcal{M}}(G)^{gp} = K_{sp} \oplus \overline{\mathcal{M}}_\eta^{gp}.$$

Denote by $q^{sp} : \overline{\mathcal{M}}(G)^{gp} \rightarrow K_{sp}$ the natural projection. Then for any element $e \in \overline{\mathcal{M}}(G)^{gp}$, we write $e = q^{gen}(e) + q^{sp}(e)$.

Possibly after a base change, we fix a global chart $\beta : \overline{\mathcal{M}}(G) \cong \overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \mathcal{M}_S$. Thus, we have a map of groups $\beta^{gp} : \overline{\mathcal{M}}_{S, \bar{s}}^{gp} \rightarrow \mathcal{M}_S^{gp}$. By [Ols03a, 3.5(i)], the group K_{sp} is generated by elements in $\overline{\mathcal{M}}(G)$, whose images in R is not zero. Consider the composition

$$(6.4.3) \quad \bar{\beta} := \nu_\pi \circ \exp \circ \beta^{gp} : K_{sp} \rightarrow \mathbb{Z},$$

where ν_π is the evaluation of of the fraction field K .

Lemma 6.4.3. *The map $\bar{\beta}$ only depends on the base S , and the fixed underlying extension $\underline{\xi}$.*

Proof. Consider an irreducible element $e \in \overline{\mathcal{M}}(G)$, whose image in R via β is $a \neq 0$. Without loss of generality, we can assume that $a = \pi^n$. By Lemma 3.3.8, there is a minimal positive integer n' , such that $n' \cdot e$ is the element associated to some minimal vertex or some special node l in G . In the first case, the the minimal vertex is specialized from a non-degenerate component over η . Hence, Lemma 6.2.1(1) implies that the degeneracy $n \cdot n'$ is uniquely determined by the generic fiber. If $n' \cdot e$ is the element associated to a special node, then this is determined by the generic fiber. Since the extended underlying map is fixed, this proves the statement. \square

Consider the map β'_η given by the composition

$$\overline{\mathcal{M}}_\eta \longrightarrow \overline{\mathcal{M}}_\eta^{gp} \longrightarrow \overline{\mathcal{M}}_{S, \bar{s}}^{gp} \xrightarrow{\beta^{gp}} \mathcal{M}_S^{gp},$$

where the middle arrow is the natural inclusion given by (6.4.2). Note that for any $e \in \overline{\mathcal{M}}_\eta$, the element $\beta'_\eta(e)$ generalize to a unique element in \mathcal{M}_η . Thus we obtain a chart for \mathcal{M}_η .

Definition 6.4.4. A chart $\beta'_\eta : \overline{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$ is called specializable, if it comes from a global chart $\beta : \overline{\mathcal{M}}_{S, \bar{s}} \rightarrow \mathcal{M}_S$ as above.

Remark 6.4.5. Note that the specializable chart depends on the choice of decomposition (6.4.2).

For any element $e \in \overline{\mathcal{M}}_{S, \bar{s}}$, consider the decomposition

$$e = q^{sp}(e) + q^{gen}(e) = e^{sp} + e^{gen}.$$

By the construction of β'_η , we have

$$(6.4.4) \quad \beta(e)_\eta = \beta^{gp}(e^{sp})_\eta + \beta'_\eta(e^{gen}), \quad \text{in } \mathcal{M}_\eta.$$

Note that $\exp \circ \beta^{gp}(e^{sp}) \in K$, hence is an invertible element in \mathcal{M}_η .

Lemma 6.4.6. *The special degeneracy with respect to β'_η only depends on $\bar{\beta}$ and q^{sp} .*

Proof. Let $Z \subset C_{\bar{s}}$ be an irreducible component, which corresponds to $v \in V(G)$. Then (6.4.4) shows that the special degeneracy with respect to the specializable chart β'_η is given by $\bar{\beta} \circ q^{sp}(e_v)$, where $e_v \in \overline{\mathcal{M}}(G)$ is the element associated to v . Now the statement follows from Observation 6.2.2. \square

6.5. Uniqueness of the extension. Assume that we have two minimal extensions $\xi_1 = (C \rightarrow S, \mathcal{M}_1, f_1)$ and $\xi_2 = (C \rightarrow S, \mathcal{M}_2, f_2)$ of ξ_η , with the same underlying $\underline{\xi}$. After a base change, we can assume that we have two global charts

$$(6.5.1) \quad \beta_1 : \overline{\mathcal{M}}(G) \rightarrow \mathcal{M}_1 \quad \text{and} \quad \beta_2 : \overline{\mathcal{M}}(G) \rightarrow \mathcal{M}_2.$$

for ξ_1 and ξ_2 respectively.

Lemma 6.5.1. *For any element $e \in \overline{\mathcal{M}}(G)$, we have a unique element $u \in R^*$ such that*

$$\beta_1(e)_\eta = u \cdot \beta_2(e)_\eta \quad \text{in } \mathcal{M}_\eta.$$

Thus, we have a canonical isomorphism of log structures $\mathcal{M}_1 \cong \mathcal{M}_2$.

Proof. We only need to consider the irreducible elements of $\overline{\mathcal{M}}(G)$. Let e be an irreducible element of $\overline{\mathcal{M}}(G)$. By Lemma 3.3.8, We have a minimal positive integer n such that $n \cdot e \in N(G)$ is either the element associated to some edge, or the element associated to some minimal vertex.

Consider the case where $n \cdot e$ is the element associated to some edge l . We identify the element $e_l \in \mathcal{M}_S^{C/S}$ smoothing l with its image in \mathcal{M}_1 or \mathcal{M}_2 , then we have

$$n \cdot \beta_1(e) + \log u_1 = e_l \quad \text{in } \mathcal{M}_1, \quad \text{and} \quad n \cdot \beta_2(e) + \log u_2 = e_l \quad \text{in } \mathcal{M}_2,$$

where $u_1, u_2 \in R^*$. By restricting to the generic point η , we have

$$e_{l, \eta} = n \cdot \beta_1(e)_\eta + \log u_1 = n \cdot \beta_2(e)_\eta + \log u_2 \quad \text{in } \mathcal{M}_\eta.$$

This implies that

$$\beta_1(e)_\eta = \log u + \beta_2(e)_\eta \quad \text{in } \mathcal{M}_\eta,$$

where $u \in R^*$ such that $u^n = u_2/u_1$. Since we fixed the isomorphism $\xi_{1, \eta} \cong \xi_{2, \eta} \cong \xi_\eta$, the element u is unique.

Next, we consider the case where $n \cdot e$ is the element associated to some minimal vertex $v' \in V(G)$, and assume that v' is specialized from $v \in V(G_\eta)$. Denote by $e^{sp} = q^{sp}(e)$ and $e^{gen} = q^{gen}(e)$. Let $\beta_{i, \eta}$ be the specializable chart induced by β_i for $i = 1, 2$. Note that by Lemma 6.4.3 and 6.4.6, the special degeneracy of v' does not depend on the choice of β_i . We may assume

$$\exp \circ \beta_1^{gp}(e^{sp}) = \exp \circ \beta_2^{gp}(e^{sp}) = \pi^{n'},$$

where $n' = \bar{\beta}(e^{sp})$. By (6.4.4), we have

$$n \cdot \beta_1(e)_\eta = n \cdot n' \cdot \log \pi + n \cdot \beta'_{1,\eta}(e^{gen})$$

and

$$n \cdot \beta_2(e)_\eta = n \cdot n' \cdot \log \pi + n \cdot \beta'_{2,\eta}(e^{gen}).$$

Comparing the special degeneracy under $\beta'_{1,\eta}$ and $\beta'_{2,\eta}$, and using Lemma 6.4.6 and 6.2.4, we obtain a unique element $u \in R^*$, which satisfies $u^n = 1$, such that

$$\beta'_{2,\eta}(e^{gen}) = \log u + \beta'_{1,\eta}(e^{gen}).$$

Note that such u is unique, since we fixed the isomorphism $\xi_{1,\eta} \cong \xi_{2,\eta}$. This proves the statement. \square

Proposition 6.5.2. *Possibly after a base change, the isomorphism $\xi_{1,\eta} \cong \xi_{2,\eta}$ can be extended uniquely to an isomorphism of ξ_1 and ξ_2 .*

Proof. For simplicity, we assume that $\xi_{1,\eta} = \xi_{2,\eta} = \xi_\eta$. We fix two global chart β_1 and β_2 as in (6.5.1). Denote by $\beta_{i,\eta} : \bar{\mathcal{M}}_\eta \rightarrow \mathcal{M}_\eta$ the specializable chart induced by β_i for $i = 1, 2$. By Lemma 6.5.1, we can identify \mathcal{M}_1 and \mathcal{M}_2 . Thus the two chart $\beta_{1,\eta}$ and $\beta_{2,\eta}$ are identical.

We first show that the following diagram of log structures commutes:

$$\begin{array}{ccc} & \mathcal{M}_S^{C/S} & \\ \psi_1 \swarrow & & \searrow \psi_2 \\ \mathcal{M}_1 & \xlongequal{\quad} & \mathcal{M}_2, \end{array}$$

where ψ_i is the structure map defining the corresponding log curve of ξ_i . Since we put the standard log structure along non-distinguished nodes, we only need to consider a special distinguished node p over the closed point. Let $e_p \in \mathcal{M}_S$ be a section smoothing p . Then we have

$$\psi_1(e_p) = \psi_2(e_p) + \log u.$$

where u is a unit in R . Since $\xi_{1,\eta} = \xi_{2,\eta} = \xi_\eta$, by restricting the above equation to the generic point η , we obtain $u = 1$. This proves the commutativity. Thus, we can identify the two log curves of ξ_1 and ξ_2 .

It remains to show that the two morphisms of log structures $f_{\xi_i}^b$ for $i = 1, 2$ are identical. Pick a point $p \in C$ over \bar{s} , then we need to prove that locally at p we have

$$(6.5.2) \quad f_{\xi_1}^b(\delta) = f_{\xi_2}^b(\delta).$$

Since the two log maps ξ_1 and ξ_2 are minimal, locally at p we have

$$\bar{f}_{\xi_1}^b(\delta) = \bar{f}_{\xi_2}^b(\delta) \text{ in } \bar{\mathcal{M}}_S.$$

Thus, locally at p , there exists an invertible function u such that

$$f_{\xi_1}^b(\delta) = f_{\xi_2}^b(\delta) + \log u.$$

Since $\xi_{1,\eta} = \xi_{2,\eta}$, by restricting to the generic fiber, we obtained that $u = 1$. This proves the equality (6.5.2) at p . Therefore, the statement of the proposition holds. \square

6.6. Proof of Theorem 1.2.1 and finiteness. Now we can give the proof of the main Theorem 1.2.1:

Proof. The boundedness is proved in Section 5, and the weak valuative criterion is proved in Section 6. Since the stack has finite diagonal, it was shown in [EHKV01, Theorem 2.7] that $\mathcal{K}_\Gamma(X^{\text{log}})$ admits a finite surjective morphism from a scheme. With this property and the weak valuative criterion, by [LMB00, Proposition 7.12] the stack is proper. The Deligne–Mumford property follows from Proposition 3.8.1.

The representability and finiteness of the map $\mathcal{K}_\Gamma(X^{\text{log}}) \rightarrow \mathcal{K}_{g,n}(X, \beta)$ follow from Proposition 3.8.4, and 3.7.5. \square

Denote by $K_\Gamma(X^{\text{log}})$ and $K_{g,n}(X, \beta)$ the coarse moduli spaces of $\mathcal{K}_\Gamma(X^{\text{log}})$ and $\mathcal{K}_{g,n}(X, \beta)$ respectively. It follows from [KM97, 1.3 Corollary] that $K_\Gamma(X^{\text{log}})$ exists and is proper. By the universal property of coarse moduli spaces, we have a natural map

$$(6.6.1) \quad K_\Gamma(X^{\text{log}}) \rightarrow K_{g,n}(X, \beta)$$

Again, since this arrow is quasi-finite, we have

Corollary 6.6.1. *The natural map (6.6.1) is finite.*

APPENDIX A. PREREQUISITES ON LOGARITHMIC GEOMETRY

A.1. Basic definitions and properties. Following [Kat89] and [Ogu06], we recall some basic terminology from logarithmic geometry.

Monoids. A *monoid* is a commutative semi-group with a unit. We usually use “+” and “0” to denote the binary operation and the unit of a monoid. A *morphism between two monoids* is required to preserve the unit.

Let P be a monoid, we can associate a group

$$P^{gp} := \{(a, b) \mid (a, b) \sim (c, d) \text{ if } \exists s \in P \text{ such that } s + a + d = s + b + c\}.$$

We have the following terminologies:

- (1) P is called *integral* if the natural map $P \rightarrow P^{gp}$ is injective.
- (2) P is called *saturated* if it is integral and satisfies that for any $p \in P^{gp}$, if $n \cdot p \in P$ for some positive integer n then $p \in P$.
- (3) P is *coherent* if it is finitely generated.
- (4) P is *fine* if it is integral and coherent.
- (5) P is *fs* if it is fine and saturated.
- (6) P is *sharp* if there are no other units except 0. A nonzero element p in a sharp monoid P is called *irreducible* if $p = a + b$ implies either $a = 0$ or $b = 0$. Denote by $\text{Irr}(P)$ the set of irreducible elements in a sharp monoid P .
- (7) A fine monoid P is called *free* if $P \cong \mathbb{N}^n$ for some positive integer n .
- (8) A monoid P is called *torsion free* if the associated group P^{gp} is torsion free.
- (9) The monoid P is called *toric* if P is fine, saturated, and sharp. Note that in this case P is automatically torsion free.
- (10) A morphism $h : Q \rightarrow P$ of two integral monoids is called *integral* if for any $a_1, a_2 \in Q$, and $b_1, b_2 \in P$ which satisfy $h(a_1)b_1 = h(a_2)b_2$, there exist $a_2, a_4 \in Q$ and $b \in P$ such that $b_1 = h(a_3)b$ and $a_1a_3 = a_2a_4$.

Denote by Mon^{int} and Mon^{sat} the categories of integral and saturated monoids respectively. Then there is a natural inclusion

$$\iota : Mon^{sat} \rightarrow Mon^{int}.$$

On the other hand, given an integral monoid M , the elements $a \in M^{gp}$, such that $m \cdot a \in M$ for some positive integer m form a saturated submonoid $M^{sat} \subset M^{gp}$. This induces another functor

$$Sat : Mon^{int} \rightarrow Mon^{sat}.$$

Proposition A.1.1. [Ogu06, Chapter I, 1.2.3(3)] *The functor Sat is left adjoint to the functor ι .*

Logarithmic structures. Let X be a scheme. A *pre-log structure* on X is a pair (\mathcal{M}, \exp) , which consists of a sheaf of monoids \mathcal{M} on the étale site $X_{\acute{e}t}$ of X , and a morphism of sheaves of monoids $\exp : \mathcal{M} \rightarrow \mathcal{O}_X$, called the *structure morphism* of \mathcal{M} . Here we view \mathcal{O}_X as a sheaf of monoid under multiplication.

A pre-log structure \mathcal{M} on X is called a *log structure* if $\exp^{-1}(\mathcal{O}_X^*) \cong \mathcal{O}_X^*$ via \exp . We sometimes omit the morphism \exp , and only use \mathcal{M} to denote the log structure if no confusion could arise. We call the pair (X, \mathcal{M}) a *log scheme*.

Given two log structures \mathcal{M} and \mathcal{N} on X , a *morphism of the log structures* $h : \mathcal{M} \rightarrow \mathcal{N}$ is a morphism of sheaves of monoids which is compatible with the structure morphisms of \mathcal{M} and \mathcal{N} .

Given a pre-log structure \mathcal{M} on X , we can associate a log structure \mathcal{M}^a given by

$$\mathcal{M}^a := \mathcal{M} \oplus_{\exp^{-1}(\mathcal{O}_X^*)} \mathcal{O}_X^*.$$

Consider a morphism of schemes $f : X \rightarrow Y$, and a log structure \mathcal{M}_Y on Y . We can define the *pull-back log structure* $f^*(\mathcal{M}_Y)$ to be the log structure associated to the pre-log structure

$$f^{-1}(\mathcal{M}_Y) \rightarrow f^{-1}(\mathcal{O}_Y) \rightarrow \mathcal{O}_X.$$

Consider two log schemes (X, \mathcal{M}_X) and (Y, \mathcal{M}_Y) . A morphism of log schemes $(X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$ is a pair (f, f^b) , where $f : X \rightarrow Y$ is a morphism of the underlying schemes, and $f^b : f^*(\mathcal{M}_Y) \rightarrow \mathcal{M}_X$ is a morphism of log structures on X . The morphism (f, f^b) is called *strict* if f^b is an isomorphism of log structures. It is called *vertical* if $\mathcal{M}_X/f^*(\mathcal{M}_Y)$ is a sheaf of groups under the induced monoidal operation. A standard example of log structures is the following:

Example A.1.2. Consider a normal crossing divisor D on a smooth scheme X . Then the following

$$\mathcal{M}_X = \{ g \in \mathcal{O}_X \mid g \text{ is invertible outside } D \}$$

with the natural injection $\mathcal{M}_X \rightarrow \mathcal{O}_X$ forms a log structure on X .

A.1.1. *Charts of log structures.* Let (X, \mathcal{M}) be a log scheme, and P a monoid. Denote by P_X the constant sheaf of monoid P on X . A chart of \mathcal{M} is a morphism $P_X \rightarrow \mathcal{M}$ such that the associated log structure of the composition $P_X \rightarrow \mathcal{M} \rightarrow \mathcal{O}_X$ is \mathcal{M} . The log structure \mathcal{M} is called a *fine log structure* on X if P is fine. If the monoid P is fs, then \mathcal{M} is called a *fs log structure*. We denote by $\mathbf{LogS\acute{c}h}$ the category of fine log schemes, and $\mathbf{LogS\acute{c}h}^{fs}$ the category of fs log schemes.

Let $\overline{\mathcal{M}} = \mathcal{M}/\mathcal{O}_X^*$ be the quotient sheaf. We call it the *characteristic of the log structure* \mathcal{M} . It is useful to notice that $\overline{f^*(\mathcal{M})} = f^{-1}(\overline{\mathcal{M}})$ for any morphism of schemes $f : Y \rightarrow X$.

A fine log structure \mathcal{M} is called locally free if for any $\bar{x} \in X$, we have $\overline{\mathcal{M}}_{\bar{x}} \cong \mathbb{N}^n$ for some positive integer n . Let $\overline{\mathcal{M}}_{\bar{x}}^{gp,tor}$ be the torsion part of $\overline{\mathcal{M}}_{\bar{x}}^{gp}$. The following result is very useful for creating charts.

Proposition A.1.3. [Ols03a, 2.1] *Using the notation as above, there exist an fppf neighborhood $f : X' \rightarrow X$ of x , and a chart $\beta : P \rightarrow f^*(\mathcal{M})$ such that for some geometric point $\bar{x}' \rightarrow X'$ lying over x , the natural map $P \rightarrow f^{-1}\overline{\mathcal{M}}_{\bar{x}'}$ is bijective. If $\overline{\mathcal{M}}_{\bar{x}}^{gp,tor} \otimes k(x) = 0$, then such a chart exists in an étale neighborhood of x .*

Remark A.1.4. If \mathcal{M} is a fs log structure on X , then the above proposition implies that there exists a section $\overline{\mathcal{M}}_{\bar{x}} \rightarrow \mathcal{M}_{\bar{x}}$, which can be lifted to a chart étale locally near x .

Consider a morphism $f : (X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$ of fine log schemes. A *chart of f* is a triple $(P_X \rightarrow \mathcal{M}_X, Q_Y \rightarrow \mathcal{M}_Y, Q \rightarrow P)$ where $P_X \rightarrow \mathcal{M}_X$ and $Q_Y \rightarrow \mathcal{M}_Y$ are charts of \mathcal{M}_X and \mathcal{M}_Y respectively, and $Q \rightarrow P$ is a morphism of monoids such that the following diagram is commutative:

$$\begin{array}{ccc} Q_X & \longrightarrow & P_X \\ \downarrow & & \downarrow \\ f^*(\mathcal{M}_Y) & \longrightarrow & \mathcal{M}_X. \end{array}$$

Note that the charts of morphism of fine log schemes exist étale locally.

Consider a morphism of log schemes $f : (X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$. With the help of charts, we can describe the log smoothness properties of f due to K. Kato [Kat89, Theorem 3.5]. The log map f is called *log smooth* if étale locally, there is a chart $(P_X \rightarrow \mathcal{M}_X, Q_Y \rightarrow \mathcal{M}_Y, Q \rightarrow P)$ of f such that:

- (1) $\text{Ker}(Q^{gp} \rightarrow P^{gp})$ and the torsion part of $\text{Coker}(Q^{gp} \rightarrow P^{gp})$ are finite groups;
- (2) the induced map $X \rightarrow Y \times_{\text{Spec}(\mathbb{Z}[Q])} \text{Spec} \mathbb{Z}[P]$ is smooth in the usual sense.

The map f is called *integral* if for every $p \in X$, the induced map $\overline{\mathcal{M}}_{f(\bar{p})} \rightarrow \overline{\mathcal{M}}_{\bar{p}}$ is integral. In general, the underlying structure map of a log smooth morphism need not be flat. However, it is shown in [Kat89, 4.5] that the underlying map of a log smooth and integral morphism is flat.

A.2. Deligne-Faltings log structures.

Definition A.2.1. Consider a scheme X . A locally free log structure \mathcal{M}_X on X is called a *Deligne-Faltings (DF) log structures*, if there is a morphism of locally constant sheaves of monoids $\beta : \mathbb{N}^k \rightarrow \overline{\mathcal{M}}_X$, which locally lifts to a chart. We call the map β a *global presentation of \mathcal{M}_X* .

Remark A.2.2. Consider a DF log structure \mathcal{M}_X with a global presentation $\mathbb{N}^k \rightarrow \overline{\mathcal{M}}_X$. Denote by $\{\delta_i\}_{i=1}^k$ the standard generators of \mathbb{N}^k . Then locally we have a lifting $\tilde{\beta} : \mathbb{N}^k \rightarrow \mathcal{M}_X$. Note that the section $\beta(\delta_i)$ with its inverse image under the canonical map $\pi : \mathcal{M}_X \rightarrow \overline{\mathcal{M}}_X$ is a \mathcal{O}_X^* -torsor, which corresponds to a line bundle L_i . The composition

$$\pi^{-1}(\beta(\delta_i)) \subset \mathcal{M}_X \rightarrow \mathcal{O}_X$$

gives a morphism of line bundles $s_i : L_i \rightarrow \mathcal{O}_X$. In fact, it was shown in [Kat89, Complement 1] that a locally free DF log structure as above is equivalent to the data consisting of a k -tuple of line bundles $(L_i)_{i=1}^k$ and morphisms of line bundles $s_i : L_i \rightarrow \mathcal{O}_X$ for each i .

Note that $s_i \in H^0(L_i^\vee)$. Denote by $D_i \subset X$ the vanishing locus of s_i . Note that D_i consists of the points where the image of δ_i in $\overline{\mathcal{M}}_X$ is non-trivial. If s_i is not a zero section, then D_i is a Cartier divisor in X . If s_i is a zero section, then $D_i = X$. Consider the sub-log structure $\mathcal{M}_X^g \subset \mathcal{M}_X$, which is given by the set of zero sections and the corresponding line bundles. We call \mathcal{M}_X^g the *generic part of \mathcal{M}_X* . Note that if $D_i = \emptyset$, then the sub-log structure generated by δ_i is trivial.

Example A.2.3. Consider a smooth Cartier divisor $D \subset X$, and the log structure \mathcal{M}_X associated to D defined in example A.1.2. Then \mathcal{M}_X forms a DF log structure on X , which corresponds to the line bundle $\mathcal{O}_X(-D)$ and the natural inclusion $\mathcal{O}_X(-D) \hookrightarrow \mathcal{O}_X$.

A.3. Olsson's Log Stacks. We follow [Ols03a] to introduce the algebraic stack parameterizing log structures. Let us fix a base scheme S , and consider an algebraic stack \mathcal{X} in the sense of [Art74], which means that

- (1) the diagonal $\mathcal{X} \rightarrow \mathcal{X} \times_S \mathcal{X}$ is representable and of finite type;
- (2) there exists a surjective smooth morphism $X \rightarrow \mathcal{X}$ from a scheme.

Now we can define a fine log structure \mathcal{M}_X on \mathcal{X} by repeating the definition of log structure on schemes in A.1, but using lisse-étale site instead of the étale site. We refer to [Ols03a, Section 5] for details of log structure on Artin stacks.

For any S -scheme T , and an arrow $g : T \rightarrow \mathcal{X}$, we obtain a fine log structure $g^*(\mathcal{M}_X)$ on the lisse-étale site $T_{\text{lisse-ét}}$ of T . It is shown in [Ols03a, 5.3] that such $g^*(\mathcal{M}_X)$ is isomorphic to a unique fine log structure on the étale site $T_{\text{ét}}$ of T . By abuse of notations, we denote by $g^*(\mathcal{M}_X)$ the corresponding log structure on T . Thus, we define a functor from \mathcal{X} to \mathbf{LogSch}_S , by pulling back the log structure \mathcal{M}_X . The stack \mathcal{X} associated with this functor is called a *log stack* in [Kat00]. A fine log scheme (X, \mathcal{M}_X) can be naturally viewed as a log algebraic stack.

Consider the fibered category $\mathcal{L}og_{(\mathcal{X}, \mathcal{M}_X)}$ over \mathcal{X} . Its objects are pairs

$$(g : X \rightarrow \mathcal{X}, g^*(\mathcal{M}_X) \rightarrow \mathcal{M}_X),$$

where g is a map from scheme X to \mathcal{X} , and $g^*(\mathcal{M}_X) \rightarrow \mathcal{M}_X$ is a morphism of fine log structures on X . An arrow

$$(g : X \rightarrow \mathcal{X}, g^*(\mathcal{M}_X) \rightarrow \mathcal{M}_X) \longrightarrow (h : Y \rightarrow \mathcal{X}, h^*(\mathcal{M}_X) \rightarrow \mathcal{M}_Y)$$

is a strict morphism of log schemes $(X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$, such that

- (1) the underlying map $X \rightarrow Y$ is a morphism over \mathcal{X} ;
- (2) the following diagram of log schemes commutes:

$$\begin{array}{ccc} (X, \mathcal{M}_X) & \longrightarrow & (Y, \mathcal{M}_Y) \\ \downarrow & & \downarrow \\ (X, g^*(\mathcal{M}_X)) & \longrightarrow & (Y, h^*(\mathcal{M}_X)). \end{array}$$

Remark A.3.1. In fact, the stack $\mathcal{L}og_{(\mathcal{X}, \mathcal{M}_X)}$ parametrizes log structures over $(\mathcal{X}, \mathcal{M}_X)$. An object $(g : X \rightarrow \mathcal{X}, g^*(\mathcal{M}_X) \rightarrow \mathcal{M}_X)$ as above can be viewed as a morphism of log stacks $(X, \mathcal{M}_X) \rightarrow (\mathcal{X}, \mathcal{M}_X)$.

Theorem A.3.2. [Ols03a, 5.9] *The fibered category $\mathcal{L}og_{(\mathcal{X}, \mathcal{M}_X)}$ is an algebraic stack locally of finite presentation over \mathcal{X} .*

APPENDIX B. LOGARITHMIC CURVES AND THEIR STACKS

In this section, we introduce the notion of log pre-stable curves, which will be smooth log curves with fs log structures. We will prove that the stack $\mathfrak{M}_{g,n}^{pre}$ parameterizing log pre-stable curves of genus g and n marked points is an open substack of some Olsson’s log stack as above, hence is algebraic in the sense of [Art74, 5.1]. We refer to [Kat00], [Moc95], and [Ols07] for more details of log structures on curves.

B.1. The canonical log structure on pre-stable curves. Consider the stack $\mathfrak{M}_{g,n}$ parameterizing genus g pre-stable curves with n marked points, and let $\mathfrak{C}_{g,n}$ be the universal family over $\mathfrak{M}_{g,n}$. Denote by $\{\Sigma_i : \mathfrak{M}_{g,n} \rightarrow \mathfrak{C}_{g,n}\}_{i=1}^n$ the n sections. The boundary $\mathfrak{M}_{g,n}^{sing} \subset \mathfrak{M}_{g,n}$ which parametrizes singular curves is a divisor with normal crossings on $\mathfrak{M}_{g,n}$. Hence the boundary divisor induces a canonical log structure $\mathcal{M}_{\mathfrak{M}_{g,n}}$ on $\mathfrak{M}_{g,n}$, which is defined using the smooth topology as in [Ols03a].

Note that each section Σ_i corresponds to a smooth divisor on $\mathfrak{C}_{g,n}$. By Example A.1.2, we have a log structure \mathcal{M}^{Σ_i} associated to this section Σ_i . The pre-image of $\mathfrak{M}_{g,n}^{sing}$ also gives a normal crossing divisor in $\mathfrak{C}_{g,n}$, hence a log structure $\mathcal{M}_{\mathfrak{C}_{g,n}}^\#$ on $\mathfrak{C}_{g,n}$. Consider the log structure

$$\mathcal{M}_{\mathfrak{C}_{g,n}} := \mathcal{M}_{\mathfrak{C}_{g,n}}^\# \oplus_{\mathcal{O}_{\mathfrak{C}_{g,n}}} \sum_i \mathcal{M}^{\Sigma_i}.$$

We call it the canonical log structure on $\mathfrak{C}_{g,n}$. There is a natural log smooth, integral, vertical map

$$(B.1.1) \quad (\mathfrak{C}_{g,n}, \mathcal{M}_{\mathfrak{C}_{g,n}}^\#) \rightarrow (\mathfrak{M}_{g,n}, \mathcal{M}_{\mathfrak{M}_{g,n}}),$$

whose underlying map is given by the family $\mathfrak{C}_{g,n} \rightarrow \mathfrak{M}_{g,n}$. By adding the log structure from the marked points, we have an induced log smooth, integral map

$$(B.1.2) \quad (\mathfrak{C}_{g,n}, \mathcal{M}_{\mathfrak{C}_{g,n}}) \rightarrow (\mathfrak{M}_{g,n}, \mathcal{M}_{\mathfrak{M}_{g,n}}).$$

Given any family $C \rightarrow S$ of usual pre-stable curves of genus g , with n marked points, we have the following cartesian diagram:

$$\begin{array}{ccc} C & \longrightarrow & \mathfrak{C}_{g,n} \\ \pi \downarrow & & \downarrow \\ S & \longrightarrow & \mathfrak{M}_{g,n}. \end{array}$$

Denote by $\mathcal{M}_C^{\#C/S}$, $\mathcal{M}_C^{C/S}$ and $\mathcal{M}_C^{\Sigma_i}$ the log structures on C , obtained by pulling back $\mathcal{M}_{\mathfrak{C}_{g,n}}^\#$, $\mathcal{M}_{\mathfrak{C}_{g,n}}$ and \mathcal{M}^{Σ_i} respectively. Let $\mathcal{M}_S^{C/S}$ be the log structure on S obtained by pulling back $\mathcal{M}_{\mathfrak{M}_{g,n}}$. Note that $\mathcal{M}_C^{\Sigma_i}$ is the log structure given by the section Σ_i . Now we have two canonical log maps obtained by pulling back (B.1.1) and (B.1.2) respectively:

$$(B.1.3) \quad (C, \mathcal{M}_C^{\#C/S}) \rightarrow (S, \mathcal{M}_S^{C/S}),$$

and

$$(B.1.4) \quad (C, \mathcal{M}_C^{C/S}) \rightarrow (S, \mathcal{M}_S^{C/S}).$$

Lemma B.1.1. *For any pair of fine log structures $(\mathcal{M}'_C, \mathcal{M}_S)$ over the family of pre-stable curves $C \rightarrow S$, such that the log map $(C, \mathcal{M}'_C) \rightarrow (S, \mathcal{M}_S)$ is log smooth, proper, integral*

and vertical, we have a unique pair of maps $\mathcal{M}_C^{\sharp C/S} \rightarrow \mathcal{M}'_C$ and $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S$ fitting in the following cartesian diagram of fine log schemes:

$$\begin{array}{ccc} (C, \mathcal{M}'_C) & \longrightarrow & (C, \mathcal{M}_C^{\sharp C/S}) \\ \downarrow & & \downarrow \\ (S, \mathcal{M}_S) & \longrightarrow & (S, \mathcal{M}_S^{C/S}), \end{array}$$

Proof. See [Ols07], and [Ols03b, 2.7] for a proof. \square

B.2. Log curves. With the description above, we are able to introduce the log structure on curves that we are interested in.

Definition B.2.1. A map of fine log schemes $(C, \mathcal{M}_C) \rightarrow (S, \mathcal{M}_S)$ with sections $\{\Sigma_i\}_{i=1}^n$ is called a genus g log curve with n -markings if

- (1) the family $C \rightarrow S$ with $\{\Sigma_i\}$ is the usual prestable curve of genus g and n -markings;
- (2) the log structure \mathcal{M}_C is of the form $\mathcal{M}_C = \mathcal{M}'_C \oplus_{\mathcal{O}_C^*} (\sum_j \mathcal{M}_C^{\Sigma_j})$;
- (3) the log map $(C, \mathcal{M}_C) \rightarrow (S, \mathcal{M}_S)$ comes from a log smooth, integral vertical map $(C, \mathcal{M}'_C) \rightarrow (S, \mathcal{M}_S)$ plus the log structure \mathcal{M}^{Σ_i} given by the markings.

By Lemma B.1.1, we have an equivalent definition of log curves using the canonical log structure.

Definition B.2.2. A genus g , log curve with n -marked points over a scheme S is given by the following data $(C \rightarrow S, \{\Sigma\}_{i=1}^n, \mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S)$, where

- (1) $(C \rightarrow S, \{\Sigma\}_{i=1}^n)$ is a usual family of pre-stable curves of genus g with n -markings;
- (2) $\mathcal{M}_S^{C/S} \rightarrow \mathcal{M}_S$ is a morphism of fine log structures.

If no confusion could arise, we will use $(C \rightarrow S, \mathcal{M}_S)$ for the log curves in the definition for short, and denote by \mathcal{M}_C for the log structure on the curves in the above Definition B.2.1.

Definition B.2.3. A log curve $(C \rightarrow S, \mathcal{M}_S)$ is called log pre-stable if the log structure \mathcal{M}_S is fine and saturated.

Remark B.2.4. By [Ols03a, 5.26], the condition that the base log structure \mathcal{M}_S is fine and saturated is an open condition on S .

B.3. The stack of log curves.

Definition B.3.1. Given two log curves $(C \rightarrow S, \mathcal{M}_S)$ and $(C' \rightarrow S, \mathcal{M}'_S)$ over S . Denote by \mathcal{M}_C and $\mathcal{M}_{C'}$ the log structure on C and C' associated to the two log curves respectively. An isomorphism between the above two log curves is a pair (ρ, θ) such that

- (1) $\theta : (S, \mathcal{M}_S) \rightarrow (S, \mathcal{M}'_S)$ and $\rho : (C, \mathcal{M}_C) \rightarrow (C', \mathcal{M}_{C'})$ are isomorphisms of log schemes;
- (2) the underlying map $\theta : S \rightarrow S$ is the identity, and $\rho : C \rightarrow C'$ is an isomorphism of usual prestable curves over S ;
- (3) the pair (ρ, θ) fit in the following commutative diagram:

$$\begin{array}{ccc} (C, \mathcal{M}_C) & \xrightarrow{\rho} & (C', \mathcal{M}_{C'}) \\ \downarrow & & \downarrow \\ (S, \mathcal{M}_S) & \xrightarrow{\theta} & (S, \mathcal{M}'_S). \end{array}$$

Denote by $\mathfrak{M}_{g,n}^{\log}$ the fibered category over \mathbb{C} parametrizing log curves with the arrow defined above. In fact, we have

$$\mathfrak{M}_{g,n}^{\log} \cong \mathcal{L}og(\mathfrak{m}_{g,n}, \mathcal{M}_{\mathfrak{m}_{g,n}}).$$

Thus, the fibered category $\mathfrak{M}_{g,n}^{\log}$ forms an algebraic stack in the sense of [Art74]. Denote by $\mathfrak{M}_{g,n}^{\text{pre}}$ the substack of $\mathfrak{M}_{g,n}^{\log}$ parameterizing log prestable curves. Then by Remark B.2.4, we have the following:

Corollary B.3.2. *The fibered category $\mathfrak{M}_{g,n}^{\text{pre}}$ is an open substack in $\mathfrak{M}_{g,n}^{\log}$, hence is algebraic.*

B.4. The canonical log structure at nodes. Note that the log structure $\mathcal{M}_{\mathfrak{m}_{g,n}}$ is locally free. Consider a usual prestable curve $C \rightarrow S$. Then the canonical log structure $\mathcal{M}_S^{C/S}$ is also locally free. For any closed point $s \in S$, we have

$$\overline{\mathcal{M}}_{S,\bar{s}}^{C/S} \cong \mathbb{N}^m,$$

where m is a non-negative integer.

Shrinking S if necessary, by Proposition A.1.3 we can choose a global chart $\overline{\mathcal{M}}_{S,\bar{s}}^{C/S} \cong \mathbb{N}^m \rightarrow \mathcal{M}_S^{C/S}$. Denote by $\{e_i\}_{i=1}^m$ the standard generators of \mathbb{N}^m .

Consider a node point $p \in C_{\bar{s}}$ in the fiber. Then there is an étale neighborhood U of p , which contains no other nodes and marked points. We have a special element $e_j \in \{e_i\}_{i=1}^m$, with the following chart:

$$\begin{array}{ccc} \mathbb{N}^{m-1} \oplus \mathbb{N}^2 & \longrightarrow & \mathcal{M}_C^{C/S}|_U \\ \uparrow (id, \Delta) & & \uparrow \pi^b \\ \mathbb{N}^{m-1} \oplus \mathbb{N} & \longrightarrow & \pi^*(\mathcal{M}_S^{C/S})|_U. \end{array}$$

Here on the bottom, the monoids \mathbb{N}^{m-1} and \mathbb{N} are generated by $\{e_i\}_{i \neq j}$ and e_j respectively, and on the top we assume that a and b are the standard generators of the monoid \mathbb{N}^2 . The map (id, Δ) is given by the identity on \mathbb{N}^{m-1} and the diagonal map $\Delta : e_j \mapsto a + b$.

Conventions B.4.1. Consider a log curve $(C \rightarrow S, \mathcal{M}_S)$. For convenience, we identify e_j with its image in $\overline{\mathcal{M}}_S$, and call it *the element in $\overline{\mathcal{M}}_S$ smoothing the node p* , or simply *the element smoothing p* .

For each node p_i over s , we fix an element $e_i \in \overline{\mathcal{M}}_{S,\bar{s}}^{C/S}$ smoothing it. Let $Irr(\overline{\mathcal{M}}_{S,\bar{s}}^{C/S})$ be the set of irreducible elements in the monoid $\overline{\mathcal{M}}_{S,\bar{s}}^{C/S}$. In fact we have $\{e_i\}_{i=1}^m = Irr(\overline{\mathcal{M}}_{S,\bar{s}}^{C/S})$, and a natural map:

$$s_{C_{\bar{s}}} : \{\text{nodes in } C_{\bar{s}}\} \rightarrow Irr(\overline{\mathcal{M}}_{S,\bar{s}}^{C/S})$$

given by $p_i \mapsto$ (the element e_i smoothing p_i). It was shown in [Kat00] that this map is a one-to-one correspondence. This means that all nodes in the fiber are smoothed independently.

Remark B.4.2. The bijection $s_{C_{\bar{s}}}$ implies that the canonical log structures $(\mathcal{M}_S^{C/S}, \mathcal{M}_C^{C/S})$ is special in the sense of [Ols03b, 2.6].

We give a local description of the relation between canonical log structure and the underlying structure at the nodes as in [Kat00, Section 3]. Let A be a local noetherian henselian ring, and s an element in the maximal ideal m_A of A . Let R be the henselization of $A[x, y]/(xy - s)$

at the ideal generated by x, y and m_A . We still use x, y to denote the corresponding elements in R .

Lemma B.4.3. [Kat00, 2.1] *Given $x', y' \in R$ such that $x'y' \in A$ and $(x', y', m_A) = (x, y, m_A)$ (equality of ideals in R). Then there exist units $u_x, u_y \in R^*$ with $u_x u_y \in A$ such that $x' = u_x x$ and $y' = u_y y$ (or $y' = u_x x$ and $x' = u_y y$).*

Consider the local family $\text{Spec } R \rightarrow \text{Spec } A$, the canonical log structure $(\mathcal{M}_R, \mathcal{M}_A)$ is given by the following commutative diagram of pre-log structures.

$$\begin{array}{ccc} \mathbb{N}^2 & \xrightarrow{(e_1, e_2) \mapsto (x, y)} & R \\ \Delta \uparrow & & \uparrow \\ \mathbb{N} & \xrightarrow{e \mapsto s} & A \end{array}$$

where e_1, e_2 (resp. e) are the standard generators of \mathbb{N}^2 (resp. \mathbb{N}), and $\Delta : e \mapsto e_1 + e_2$ is the diagonal map. For convenience, we sometimes use $\log x, \log y$ and $\log s$ denote the image of e_1, e_2 and e in the corresponding log structures.

REFERENCES

- [Abr94] Dan Abramovich, *Subvarieties of semiabelian varieties*, *Compositio Math.* **90** (1994), no. 1, 37–52. MR MR1266493 (95c:14054)
- [AC] Dan Abramovich and Qile Chen, *Logarithmic stable maps to Deligne-Faltings pairs II*, In preparation.
- [AF] Dan Abramovich and Barbara Fantechi, *Orbifold techniques in degeneration formulas*, preprint.
- [Art74] M. Artin, *Versal deformations and algebraic stacks*, *Invent. Math.* **27** (1974), 165–189. MR MR0399094 (53 #2945)
- [AV02] Dan Abramovich and Angelo Vistoli, *Compactifying the space of stable maps*, *J. Amer. Math. Soc.* **15** (2002), no. 1, 27–75 (electronic). MR MR1862797 (2002i:14030)
- [BF97] K. Behrend and B. Fantechi, *The intrinsic normal cone*, *Invent. Math.* **128** (1997), no. 1, 45–88. MR MR1437495 (98e:14022)
- [Che] Qile Chen, *The degeneration formula for logarithmic expanded degenerations*, arXiv:1009.4378v1 [math.AG], preprint.
- [EHKV01] Dan Edidin, Brendan Hassett, Andrew Kresch, and Angelo Vistoli, *Brauer groups and quotient stacks*, *Amer. J. Math.* **123** (2001), no. 4, 761–777. MR MR1844577 (2002f:14002)
- [FP97] W. Fulton and R. Pandharipande, *Notes on stable maps and quantum cohomology*, *Algebraic geometry—Santa Cruz 1995*, *Proc. Sympos. Pure Math.*, vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 45–96. MR MR1492534 (98m:14025)
- [Ful93] William Fulton, *Introduction to toric varieties*, *Annals of Mathematics Studies*, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry. MR MR1234037 (94g:14028)
- [Gro61] A. Grothendieck, *Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I*, *Inst. Hautes Études Sci. Publ. Math.* (1961), no. 11, 167. MR MR0163910 (29 #1209)
- [GS] Mark Gross and Bernd Siebert, *Logarithmic Gromov-Witten invariants*, preprint.
- [HM82] Joe Harris and David Mumford, *On the Kodaira dimension of the moduli space of curves*, *Invent. Math.* **67** (1982), no. 1, 23–88, With an appendix by William Fulton. MR MR664324 (83i:14018)
- [IP03] Eleny-Nicoleta Ionel and Thomas H. Parker, *Relative Gromov-Witten invariants*, *Ann. of Math.* (2) **157** (2003), no. 1, 45–96. MR MR1954264 (2004a:53112)
- [Kat89] Kazuya Kato, *Logarithmic structures of Fontaine-Illusie*, *Algebraic analysis, geometry, and number theory* (Baltimore, MD, 1988), *Johns Hopkins Univ. Press*, Baltimore, MD, 1989, pp. 191–224. MR MR1463703 (99b:14020)

- [Kat00] Fumiharu Kato, *Log smooth deformation and moduli of log smooth curves*, Internat. J. Math. **11** (2000), no. 2, 215–232. MR MR1754621 (2001d:14016)
- [Kim09] Bumsig Kim, *Logarithmic stable maps*, arXiv:0807.3611v2 (2009), To appear in the proceedings volume of the conference “New developments in Algebraic Geometry, Integrable Systems and Mirror symmetry”, RIMS, Kyoto.
- [KM97] Seán Keel and Shigefumi Mori, *Quotients by groupoids*, Ann. of Math. (2) **145** (1997), no. 1, 193–213. MR 1432041 (97m:14014)
- [Li01] Jun Li, *Stable morphisms to singular schemes and relative stable morphisms*, J. Differential Geom. **57** (2001), no. 3, 509–578. MR MR1882667 (2003d:14066)
- [Li02] ———, *A degeneration formula of GW-invariants*, J. Differential Geom. **60** (2002), no. 2, 199–293. MR MR1938113 (2004k:14096)
- [LMB00] Gérard Laumon and Laurent Moret-Bailly, *Champs algébriques*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 39, Springer-Verlag, Berlin, 2000. MR MR1771927 (2001f:14006)
- [LR01] An-Min Li and Yongbin Ruan, *Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds*, Invent. Math. **145** (2001), no. 1, 151–218. MR MR1839289 (2002g:53158)
- [Moc95] Shinichi Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci. **31** (1995), no. 3, 355–441. MR MR1355945 (96j:14017)
- [Ogu06] Arthur Ogus, *Lectures on logarithmic algebraic geometry*, TeXed notes (2006).
- [Ols03a] Martin C. Olsson, *Logarithmic geometry and algebraic stacks*, Ann. Sci. École Norm. Sup. (4) **36** (2003), no. 5, 747–791. MR MR2032986 (2004k:14018)
- [Ols03b] ———, *Universal log structures on semi-stable varieties*, Tohoku Math. J. (2) **55** (2003), no. 3, 397–438. MR MR1993863 (2004f:14025)
- [Ols05] ———, *The logarithmic cotangent complex*, Math. Ann. **333** (2005), no. 4, 859–931. MR MR2195148 (2006j:14017)
- [Ols06] ———, *Hom-stacks and restriction of scalars*, Duke Math. J. **134** (2006), no. 1, 139–164. MR MR2239345 (2007f:14002)
- [Ols07] ———, *(Log) twisted curves*, Compos. Math. **143** (2007), no. 2, 476–494. MR MR2309994 (2008d:14021)
- [Par09a] Brett Parker, *Exploded manifolds*, arXiv:0910.4201v1 (2009).
- [Par09b] ———, *Holomorphic curves in exploded manifolds: Compactness*, arXiv:0911.2241v1 (2009).
- [Par09c] ———, *Holomorphic curves in exploded torus fibrations: Regularity*, arXiv:0902.0087v1 (2009).
- [Ran87] Ziv Ran, *The degree of a Severi variety*, Bull. Amer. Math. Soc. (N.S.) **17** (1987), no. 1, 125–128. MR MR888887 (88i:14019)
- [SGA64] *Schémas en groupes. II: Groupes de type multiplicatif, et structure des schémas en groupes généraux*, Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 152, Springer-Verlag, Berlin, 1962/1964. MR MR0274459 (43 #223b)
- [Sie01] Bernd Siebert, *Gromov-Witten invariants in relative and singular cases*, Lecture given in the workshop on algebraic aspects of mirror symmetry, Universität Kaiserslautern, Germany, Jun. 26 2001.

QILE CHEN, DEPARTMENT OF MATHEMATICS, BOX 1917, BROWN UNIVERSITY, PROVIDENCE, RI, 02912, U.S.A

E-mail address: q.chen@math.brown.edu