

# Resolving singularities of varieties and families

Dan Abramovich  
Brown University

Joint work with Michael Temkin and Jarosław Włodarczyk



Rio de Janeiro  
August 7, 2018

# Resolving singularities of **varieties** and families

Dan Abramovich  
Brown University

Joint work with Michael Temkin and Jarosław Włodarczyk



Rio de Janeiro  
August 7, 2018

# Resolving singularities of varieties and families


Dan Abramovich  
Brown University

Joint work with Michael Temkin and Jarosław Włodarczyk



Rio de Janeiro  
August 7, 2018


# On singularities




Zitrus:  $x^2 + z^2 = y^3(1 - y)^3$

Real figures by Herwig Hauser, <https://imaginary.org/gallery/herwig-hauser-classic>

# On singularities




Zitrus:  $x^2 + z^2 = y^3(1 - y)^3$




Kolibri:  $x^2 = y^2 z^2 + z^3$

Real figures by Herwig Hauser, <https://imaginary.org/gallery/herwig-hauser-classic>


# On singularities



Zitrus:  $x^2 + z^2 = y^3(1 - y)^3$



Kolibri:  $x^2 = y^2 z^2 + z^3$




Daisy:  $(x^2 - y^3)^2 = (z^2 - y^2)^3$


Real figures by Herwig Hauser, <https://imaginary.org/gallery/herwig-hauser-classic>

Singularities are beautiful.


# On singularities



Zitrus:  $x^2 + z^2 = y^3(1 - y)^3$



Kolibri:  $x^2 = y^2z^2 + z^3$




Daisy:  $(x^2 - y^3)^2 = (z^2 - y^2)^3$


Real figures by Herwig Hauser, <https://imaginary.org/gallery/herwig-hauser-classic>

Singularities are beautiful.  
Why should we “get rid of them”?


# On singularities



Zitrus:  $x^2 + z^2 = y^3(1 - y)^3$



Kolibri:  $x^2 = y^2z^2 + z^3$



Daisy:  $(x^2 - y^3)^2 = (z^2 - y^2)^3$

Real figures by Herwig Hauser, <https://imaginary.org/gallery/herwig-hauser-classic>

Singularities are beautiful.

Why should we “get rid of them”?

Answer 1: to study singularities.

Answer 2: to study the structure of varieties.



# Singular and smooth points

## Definition

$\{f(x_1, \dots, x_n) = 0\}$  is **singular** at  $p$  if  $\frac{\partial f}{\partial x_i}(p) = 0$  for all  $i$ .  
Otherwise **smooth**.

# Singular and smooth points

## Definition

$\{f(x_1, \dots, x_n) = 0\}$  is **singular** at  $p$  if  $\frac{\partial f}{\partial x_i}(p) = 0$  for all  $i$ .  
Otherwise **smooth**.

In other words, if smooth,  $\{f = 0\}$  defines a submanifold of complex codimension 1.

# Singular and smooth points

## Definition

$\{f(x_1, \dots, x_n) = 0\}$  is **singular** at  $p$  if  $\frac{\partial f}{\partial x_i}(p) = 0$  for all  $i$ .  
Otherwise **smooth**.

In other words, if smooth,  $\{f = 0\}$  defines a submanifold of complex codimension 1.

In codimension  $c$ , the set  $\{f_1 = \dots = f_k = 0\}$  is smooth when  $d(f_1, \dots, f_k)$  has constant rank  $c$ .

# What is resolution of singularities?

## Definition

A resolution of singularities  $X' \rightarrow X$  is a **modification**<sup>a</sup> with  $X'$  nonsingular inducing an isomorphism over the smooth locus of  $X$ .

---

<sup>a</sup>proper birational map. For instance, blowing up.

# What is resolution of singularities?

## Definition

A resolution of singularities  $X' \rightarrow X$  is a **modification**<sup>a</sup> with  $X'$  nonsingular inducing an isomorphism over the smooth locus of  $X$ .

<sup>a</sup>proper birational map. For instance, blowing up.

## Theorem (Hironaka 1964)

*A variety  $X$  over a field of characteristic 0 admits a resolution of singularities  $X' \rightarrow X$ , so that the critical locus  $E \subset X'$  is a simple normal crossings divisor.<sup>a</sup>*

<sup>a</sup>Codim. 1, smooth components meeting transversally - as simple as possible

Always characteristic 0 ...




## Answer 1: Example of invariant - Stepanov's theorem

If  $X' \rightarrow X$  a resolution with critical  $E \subset X'$  a simple normal crossings divisor, define  $\Delta(E)$  to be the dual complex of  $E$ .


## Answer 1: Example of invariant - Stepanov's theorem

If  $X' \rightarrow X$  a resolution with critical  $E \subset X'$  a simple normal crossings divisor, define  $\Delta(E)$  to be the dual complex of  $E$ .



## Answer 1: Example of invariant - Stepanov's theorem


If  $X' \rightarrow X$  a resolution with critical  $E \subset X'$  a simple normal crossings divisor, define  $\Delta(E)$  to be the dual complex of  $E$ .





## Answer 1: Example of invariant - Stepanov's theorem

If  $X' \rightarrow X$  a resolution with critical  $E \subset X'$  a simple normal crossings divisor, define  $\Delta(E)$  to be the dual complex of  $E$ .



### Theorem (Stepanov 2006)

*The simple homotopy type of  $\Delta(E)$  is independent of the resolution  $X' \rightarrow X$ .*

Also work by Danilov, Payne, Thuillier, Harper...

## Answer 2: Example of structure result: compactifications

“Working with noncompact spaces is like trying to keep change with holes in your pockets”

Angelo Vistoli

## Answer 2: Example of structure result: compactifications

“Working with noncompact spaces is like trying to keep change with holes in your pockets”

Angelo Vistoli

### Corollary (Hironaka)

*A smooth quasiprojective variety  $X^0$  has a smooth projective compactification  $X$  with  $D = X \setminus X^0$  a simple normal crossings divisor.*

## Answer 2: Example of structure result: compactifications

“Working with noncompact spaces is like trying to keep change with holes in your pockets”

Angelo Vistoli

### Corollary (Hironaka)

*A smooth quasiprojective variety  $X^0$  has a smooth projective compactification  $X$  with  $D = X \setminus X^0$  a simple normal crossings divisor.*



## Answer 2: Example of structure result: compactifications

“Working with noncompact spaces is like trying to keep change with holes in your pockets”

Angelo Vistoli

### Corollary (Hironaka)

*A smooth quasiprojective variety  $X^0$  has a smooth projective compactification  $X$  with  $D = X \setminus X^0$  a simple normal crossings divisor.*



# Resolution of families: $\dim B = 1$

## Key Question


When are the singularities of a morphism  $X \rightarrow B$  simple?

# Resolution of families: $\dim B = 1$

## Key Question

When are the singularities of a morphism  $X \rightarrow B$  simple?

- If  $\dim B = 1$  the simplest one can have by modifying  $X$  is  $t = \prod x_i^{a_i}$ ,
- and if one also allows base change  $t = s^k$ , can have  $s = \prod x_i$ .  
[Kempf–Knudsen–Mumford–Saint-Donat 1973]



## Question

What makes these special?

# Log smooth schemes and log smooth morphisms

- A **toric variety** is a normal variety on which  $T = (\mathbb{C}^*)^n$  acts algebraically with a dense free orbit.
- Zariski locally defined by equations between monomials.



# Log smooth schemes and log smooth morphisms

- A **toric variety** is a normal variety on which  $T = (\mathbb{C}^*)^n$  acts algebraically with a dense free orbit.
- Zariski locally defined by equations between monomials.
- A variety  $X$  with divisor  $D$  is **toroidal** or **log smooth** if étale locally it looks like a toric variety  $X_\sigma$  with its toric divisor  $X_\sigma \setminus T$ .
- Étale locally it is defined by equations between monomials.

# Log smooth schemes and log smooth morphisms

- A **toric variety** is a normal variety on which  $T = (\mathbb{C}^*)^n$  acts algebraically with a dense free orbit.
- Zariski locally defined by equations between monomials.
- A variety  $X$  with divisor  $D$  is **toroidal** or **log smooth** if étale locally it looks like a toric variety  $X_\sigma$  with its toric divisor  $X_\sigma \setminus T$ .
- Étale locally it is defined by equations between monomials.
- A morphism  $X \rightarrow Y$  is **toroidal** or **log smooth** if étale locally it looks like a torus equivariant morphism of toric varieties.
- The inverse image of a monomial <sup>1</sup> is a monomial.

---

<sup>1</sup>defining equation of part of  $D_Y$

# Resolution of families: higher dimensional base

## Question

When are the singularities of a morphism  $X \rightarrow B$  simple?

The best one can hope for, after base change, is a **semistable** morphism:

## Definition (N-Karu 2000)

A log smooth morphism, with  $B$  smooth, is *semistable* if locally

$$\begin{aligned}t_1 &= x_1 \cdots x_{l_1} \\ &\vdots \\ t_m &= x_{l_{m-1}+1} \cdots x_{l_m}\end{aligned}$$

**In particular log smooth.**

Similar definition by Berkovich, all inspired by de Jong.

# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.

# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.
- (Tight) If the geometric generic fiber  $X_{\bar{\eta}}$  is smooth, such  $X_1 \rightarrow B_1$  can be found with  $X_{\bar{\eta}}$  unchanged.

---

<sup>a</sup>Alteration: Proper, surjective, generically finite

# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.
- (Tight) If the geometric generic fiber  $X_{\bar{\eta}}$  is smooth, such  $X_1 \rightarrow B_1$  can be found with  $X_{\bar{\eta}}$  unchanged.

---

<sup>a</sup>Alteration: Proper, surjective, generically finite

- One wants the **tight** version in order to compactify smooth families.

# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.
- (Tight) If the geometric generic fiber  $X_{\bar{\eta}}$  is smooth, such  $X_1 \rightarrow B_1$  can be found with  $X_{\bar{\eta}}$  unchanged.

---

<sup>a</sup>Alteration: Proper, surjective, generically finite

- One wants the **tight** version in order to compactify smooth families.
- I'll describe progress towards that.

# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.
- (Tight) If the geometric generic fiber  $X_{\bar{\eta}}$  is smooth, such  $X_1 \rightarrow B_1$  can be found with  $X_{\bar{\eta}}$  unchanged.

---

<sup>a</sup>Alteration: Proper, surjective, generically finite

- One wants the **tight** version in order to compactify smooth families.
- I'll describe progress towards that.
- Major early results by [KKMS 1973], [de Jong 1997].



# Ultimate goal: the semistable reduction problem

## Conjecture [N-Karu]

Let  $X \rightarrow B$  be a dominant morphism of varieties.

- (Loose) There is a base change<sup>a</sup>  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is semistable.
- (Tight) If the geometric generic fiber  $X_{\bar{\eta}}$  is smooth, such  $X_1 \rightarrow B_1$  can be found with  $X_{\bar{\eta}}$  unchanged.

---

<sup>a</sup>Alteration: Proper, surjective, generically finite

- One wants the **tight** version in order to compactify smooth families.
- I'll describe progress towards that.
- Major early results by [KKMS 1973], [de Jong 1997].
- Wonderful results in positive and mixed characteristics by de Jong, Gabber, Illusie and Temkin.

# Toroidalization and weak semistable reduction

This is key to what's known:

Theorem (Toroidalization,  $\aleph$ -Karu 2000,  $\aleph$ -K-Denef 2013)

There is a *modification*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat.

Theorem (Weak semistable reduction,  $\aleph$ -Karu 2000)

There is a *base change*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth, flat, *with reduced fibers*.

# Toroidalization and weak semistable reduction

This is key to what's known:

Theorem (Toroidalization,  $\aleph$ -Karu 2000,  $\aleph$ -K-Denef 2013)

There is a *modification*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat.

Theorem (Weak semistable reduction,  $\aleph$ -Karu 2000)

There is a *base change*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth, flat, *with reduced fibers*.

- Passing from weak semistable reduction to semistable reduction is a purely combinatorial problem [ $\aleph$ -Karu 2000],

# Toroidalization and weak semistable reduction

This is key to what's known:

Theorem (Toroidalization,  $\aleph$ -Karu 2000,  $\aleph$ -K-Denef 2013)

There is a *modification*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat.

Theorem (Weak semistable reduction,  $\aleph$ -Karu 2000)

There is a *base change*  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth, flat, *with reduced fibers*.

- Passing from weak semistable reduction to semistable reduction is a purely combinatorial problem [ $\aleph$ -Karu 2000],
- proven by [Karu 2000] for families of surfaces and threefolds, and

# Toroidalization and weak semistable reduction

This is key to what's known:

Theorem (Toroidalization,  $\aleph$ -Karu 2000,  $\aleph$ -K-Denef 2013)

There is a **modification**  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat.

Theorem (Weak semistable reduction,  $\aleph$ -Karu 2000)

There is a **base change**  $B_1 \rightarrow B$  and a modification  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth, flat, **with reduced fibers**.

- Passing from weak semistable reduction to semistable reduction is a purely combinatorial problem [ $\aleph$ -Karu 2000],
- proven by [Karu 2000] for families of surfaces and threefolds, and
- whose restriction to rank-1 valuation rings is proven in a preprint by [Karim Adiprasito - Gaku Liu - Igor Pak - Michael Temkin].

# Applications of **weak** semistable reduction

This is already useful for studying families:

Theorem (Karu 2000; K-SB 97, Alexeev 94, BCHM 11)

*The moduli space of stable smoothable varieties is projective<sup>a</sup>.*

---

<sup>a</sup>in particular bounded and proper

# Applications of **weak** semistable reduction

This is already useful for studying families:

Theorem (Karu 2000; K-SB 97, Alexeev 94, BCHM 11)

*The moduli space of stable smoothable varieties is projective<sup>a</sup>.*

---

<sup>a</sup>in particular bounded and proper

Theorem (Viehweg-Zuo 2004)

*The moduli space of canonically polarized manifolds is Brody hyperbolic.*

# Applications of weak semistable reduction

This is already useful for studying families:

Theorem (Karu 2000; K-SB 97, Alexeev 94, BCHM 11)

*The moduli space of stable smoothable varieties is projective<sup>a</sup>.*

---

<sup>a</sup>in particular bounded and proper

Theorem (Viehweg-Zuo 2004)

*The moduli space of canonically polarized manifolds is Brody hyperbolic.*

Theorem (Fujino 2017)

*Nakayama's numerical logarithmic Kodaira dimension is subadditive in families  $X \rightarrow B$  with generic fiber  $F$ :*

$$\kappa_\sigma(X, D_X) \geq \kappa_\sigma(F, D_F) + \kappa_\sigma(B, D_B).$$



# Main result

The following result is work-in-progress.

## Main result (Functorial toroidalization, $\aleph$ -Temkin-Włodarczyk)

Let  $X \rightarrow B$  be a dominant morphism.

- There are modifications  $B_1 \rightarrow B$  and  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat;
- this is compatible with base change  $B' \rightarrow B$ ;
- this is functorial, up to base change, with **log** smooth  $X'' \rightarrow X$ .

This implies the **tight** version of the results of semistable reduction type.

# Main result

The following result is work-in-progress.

## Main result (Functorial toroidalization, $\aleph$ -Temkin-Włodarczyk)

Let  $X \rightarrow B$  be a dominant morphism.

- There are modifications  $B_1 \rightarrow B$  and  $X_1 \rightarrow (X \times_B B_1)_{\text{main}}$  such that  $X_1 \rightarrow B_1$  is log smooth and flat;
- this is compatible with base change  $B' \rightarrow B$ ;
- this is functorial, up to base change, with **log** smooth  $X'' \rightarrow X$ .

This implies the **tight** version of the results of semistable reduction type.  
Application:

## Theorem (Deng 2018)

*The moduli space of minimal complex projective manifolds of general type is Kobayashi hyperbolic.*

## $\dim B = 0$ : log resolution via principalization

- To resolve **log** singularities, one embeds  $X$  in a **log** smooth  $Y \dots$

## $\dim B = 0$ : log resolution via principalization

- To resolve **log** singularities, one embeds  $X$  in a **log** smooth  $Y \dots$
- $\dots$  which can be done locally.

## $\dim B = 0$ : log resolution via principalization

- To resolve **log** singularities, one embeds  $X$  in a **log** smooth  $Y \dots$
- $\dots$  which can be done locally.
- One reduces to **principalization** of  $\mathcal{I}_X$  (Hironaka, Villamayor, Bierstone–Milman).

### Theorem (Principalization ... N-T-W)

*Let  $\mathcal{I}$  be an ideal on a log smooth  $Y$ . There is a functorial logarithmic morphism  $Y' \rightarrow Y$ , with  $Y'$  logarithmically smooth, and  $\mathcal{I}\mathcal{O}_{Y'}$  an invertible monomial ideal.*

## $\dim B = 0$ : log resolution via principalization

- To resolve **log** singularities, one embeds  $X$  in a **log** smooth  $Y \dots$
- $\dots$  which can be done locally.
- One reduces to **principalization** of  $\mathcal{I}_X$  (Hironaka, Villamayor, Bierstone–Milman).

### Theorem (Principalization ... N-T-W)

Let  $\mathcal{I}$  be an ideal on a log smooth  $Y$ . There is a functorial logarithmic morphism  $Y' \rightarrow Y$ , with  $Y'$  logarithmically smooth, and  $\mathcal{I}\mathcal{O}_{Y'}$  an invertible monomial ideal.



**Figure:** The ideal  $(u^2, x^2)$   
Here  $u$  is a monomial but  $x$  is not.

## $\dim B = 0$ : log resolution via principalization

- To resolve **log** singularities, one embeds  $X$  in a **log** smooth  $Y \dots$
- $\dots$  which can be done locally.
- One reduces to **principalization** of  $\mathcal{I}_X$  (Hironaka, Villamayor, Bierstone–Milman).

### Theorem (Principalization ... N-T-W)

Let  $\mathcal{I}$  be an ideal on a log smooth  $Y$ . There is a functorial logarithmic morphism  $Y' \rightarrow Y$ , with  $Y'$  logarithmically smooth, and  $\mathcal{I}\mathcal{O}_{Y'}$  an invertible monomial ideal.



**Figure:** The ideal  $(u^2, x^2)$  and the result of blowing up the origin,  $\mathcal{I}_E^2$ . Here  $u$  is a monomial but  $x$  is not.

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .



# Logarithmic order

Principlization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ .

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\text{logord}_p(u^2, x) =$$

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\text{logord}_p(u^2, x) = 1 \quad \left(\text{since } \frac{\partial}{\partial x} x = 1\right)$$

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\text{logord}_p(u^2, x) = 1 \quad (\text{since } \frac{\partial}{\partial x} x = 1)$$

$$\text{logord}_p(u^2, x^2) =$$

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\begin{aligned} \text{logord}_p(u^2, x) &= 1 && \text{(since } \frac{\partial}{\partial x} x = 1) \\ \text{logord}_p(u^2, x^2) &= 2 && \text{logord}_p(v, x^2) = \end{aligned}$$

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\begin{aligned} \text{logord}_p(u^2, x) &= 1 && \text{(since } \frac{\partial}{\partial x} x = 1) \\ \text{logord}_p(u^2, x^2) &= 2 && \text{logord}_p(v, x^2) = 2 \\ \text{logord}_p(v + u) &= \end{aligned}$$

# Logarithmic order

Principalization is done by **order reduction**, using **logarithmic derivatives**.

- for a monomial  $u$  we use  $u \frac{\partial}{\partial u}$ .
- for other variables  $x$  use  $\frac{\partial}{\partial x}$ .

## Definition

Write  $\mathcal{D}^{\leq a}$  for the sheaf of **logarithmic** differential operators of order  $\leq a$ . The **logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}^{\leq a} \mathcal{I} = (1)$ .

Take  $u, v$  monomials,  $x$  free variable,  $p$  the origin.

$$\begin{aligned} \text{logord}_p(u^2, x) &= 1 && \text{(since } \frac{\partial}{\partial x} x = 1) \\ \text{logord}_p(u^2, x^2) &= 2 && \text{logord}_p(v, x^2) = 2 \\ \text{logord}_p(v + u) &= \infty && \text{since } \mathcal{D}^{\leq 1} \mathcal{I} = \mathcal{D}^{\leq 2} \mathcal{I} = \dots = (u, v). \end{aligned}$$



## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

### Proposition (Kollár, X-T-W)

- (1) *In characteristic 0,  $\mathcal{M}(\mathcal{I}) = \mathcal{D}^\infty(\mathcal{I})$ . In particular  $\max_p \log \text{ord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M}(\mathcal{I}) \neq 1$ .*

## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

### Proposition (Kollár, Xi-T-W)

- (1) *In characteristic 0,  $\mathcal{M}(\mathcal{I}) = \mathcal{D}^\infty(\mathcal{I})$ . In particular  $\max_p \log \text{ord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M}(\mathcal{I}) \neq 1$ .*
- (2) *Let  $Y_0 \rightarrow Y$  be the normalized blowup of  $\mathcal{M}(\mathcal{I})$ .*

## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

### Proposition (Kollár, Xi-T-W)

- (1) *In characteristic 0,  $\mathcal{M}(\mathcal{I}) = \mathcal{D}^\infty(\mathcal{I})$ . In particular  $\max_p \log \text{ord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M}(\mathcal{I}) \neq 1$ .*
- (2) *Let  $Y_0 \rightarrow Y$  be the normalized blowup of  $\mathcal{M}(\mathcal{I})$ . Then  $\mathcal{M} := \mathcal{M}(\mathcal{I})\mathcal{O}_{Y_0} = \mathcal{M}(\mathcal{I}\mathcal{O}_{Y_0})$ , and it is an invertible monomial ideal,*

## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

### Proposition (Kollár, Xi-T-W)

- (1) *In characteristic 0,  $\mathcal{M}(\mathcal{I}) = \mathcal{D}^\infty(\mathcal{I})$ . In particular  $\max_p \log \text{ord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M}(\mathcal{I}) \neq 1$ .*
- (2) *Let  $Y_0 \rightarrow Y$  be the normalized blowup of  $\mathcal{M}(\mathcal{I})$ . Then  $\mathcal{M} := \mathcal{M}(\mathcal{I})\mathcal{O}_{Y_0} = \mathcal{M}(\mathcal{I}\mathcal{O}_{Y_0})$ , and it is an invertible monomial ideal, and so  $\mathcal{I}\mathcal{O}_{Y_0} = \mathcal{I}_0 \cdot \mathcal{M}$  with  $\max_p \log \text{ord}_p(\mathcal{I}_0) < \infty$ .*

## Key new ingredient: The monomial part of an ideal

### Definition

$\mathcal{M}(\mathcal{I})$  is the minimal monomial ideal containing  $\mathcal{I}$ .

### Proposition (Kollár, X-T-W)

- (1) *In characteristic 0,  $\mathcal{M}(\mathcal{I}) = \mathcal{D}^\infty(\mathcal{I})$ . In particular  $\max_p \log \text{ord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M}(\mathcal{I}) \neq 1$ .*
- (2) *Let  $Y_0 \rightarrow Y$  be the normalized blowup of  $\mathcal{M}(\mathcal{I})$ . Then  $\mathcal{M} := \mathcal{M}(\mathcal{I})\mathcal{O}_{Y_0} = \mathcal{M}(\mathcal{I}\mathcal{O}_{Y_0})$ , and it is an invertible monomial ideal, and so  $\mathcal{I}\mathcal{O}_{Y_0} = \mathcal{I}_0 \cdot \mathcal{M}$  with  $\max_p \log \text{ord}_p(\mathcal{I}_0) < \infty$ .*

### (1) $\Rightarrow$ (2)

$\mathcal{D}_{Y_0}$  is the pullback of  $\mathcal{D}_Y$ , so (2) follows from (1) since the ideals have the same generators.

# The monomial part of an ideal - proof

Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .

# The monomial part of an ideal - proof

## Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .
- The operators

$$1, u_1 \frac{\partial}{\partial u_1}, \dots, u_l \frac{\partial}{\partial u_l}$$

commute and have distinct systems of eigenvalues on the eigenspaces  $u \mathbb{C}[x_1, \dots, x_n]$ , for distinct monomials  $u$ .



# The monomial part of an ideal - proof

## Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .
- The operators

$$1, u_1 \frac{\partial}{\partial u_1}, \dots, u_l \frac{\partial}{\partial u_l}$$

commute and have distinct systems of eigenvalues on the eigenspaces  $u \mathbb{C}[x_1, \dots, x_n]$ , for distinct monomials  $u$ .

- Therefore  $\mathcal{M} = \bigoplus u \mathcal{M}_u$  with ideals  $\mathcal{M}_u \subset \mathbb{C}[x_1, \dots, x_n]$  stable under derivatives,

# The monomial part of an ideal - proof

## Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .
- The operators

$$1, u_1 \frac{\partial}{\partial u_1}, \dots, u_l \frac{\partial}{\partial u_l}$$

commute and have distinct systems of eigenvalues on the eigenspaces  $u \mathbb{C}[x_1, \dots, x_n]$ , for distinct monomials  $u$ .

- Therefore  $\mathcal{M} = \bigoplus u \mathcal{M}_u$  with ideals  $\mathcal{M}_u \subset \mathbb{C}[x_1, \dots, x_n]$  stable under derivatives,
- so each  $\mathcal{M}_u$  is either (0) or (1).

# The monomial part of an ideal - proof

## Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .
- The operators

$$1, u_1 \frac{\partial}{\partial u_1}, \dots, u_l \frac{\partial}{\partial u_l}$$

commute and have distinct systems of eigenvalues on the eigenspaces  $u \mathbb{C}[x_1, \dots, x_n]$ , for distinct monomials  $u$ .

- Therefore  $\mathcal{M} = \bigoplus u \mathcal{M}_u$  with ideals  $\mathcal{M}_u \subset \mathbb{C}[x_1, \dots, x_n]$  stable under derivatives,
- so each  $\mathcal{M}_u$  is either (0) or (1).
- In other words,  $\mathcal{M}$  is monomial.

# The monomial part of an ideal - proof

## Proof of (1), basic affine case.

- Let  $\mathcal{O}_Y = \mathbb{C}[x_1, \dots, x_n, u_1, \dots, u_m]$  and assume  $\mathcal{M} = \mathcal{D}(\mathcal{M})$ .
- The operators

$$1, u_1 \frac{\partial}{\partial u_1}, \dots, u_l \frac{\partial}{\partial u_l}$$

commute and have distinct systems of eigenvalues on the eigenspaces  $u \mathbb{C}[x_1, \dots, x_n]$ , for distinct monomials  $u$ .

- Therefore  $\mathcal{M} = \bigoplus u \mathcal{M}_u$  with ideals  $\mathcal{M}_u \subset \mathbb{C}[x_1, \dots, x_n]$  stable under derivatives,
- so each  $\mathcal{M}_u$  is either (0) or (1).
- In other words,  $\mathcal{M}$  is monomial.



The general case requires more commutative algebra.

## $\dim B = 0$ : sketch of argument

- In characteristic 0, if  $\log \text{ord}_p(\mathcal{I}) = a < \infty$ , then  $\mathcal{D}^{\leq a-1}\mathcal{I}$  contains an element  $x$  with derivative 1, a maximal contact element.

## $\dim B = 0$ : sketch of argument

- In characteristic 0, if  $\log \text{ord}_p(\mathcal{I}) = a < \infty$ , then  $\mathcal{D}^{\leq a-1}\mathcal{I}$  contains an element  $x$  with derivative 1, a maximal contact element.
- Carefully applying induction on dimension to an ideal on  $\{x = 0\}$  gives order reduction (Encinas–Villamayor, Bierstone–Milman, Włodarczyk):

## $\dim B = 0$ : sketch of argument

- In characteristic 0, if  $\log \text{ord}_p(\mathcal{I}) = a < \infty$ , then  $\mathcal{D}^{\leq a-1}\mathcal{I}$  contains an element  $x$  with derivative 1, a maximal contact element.
- Carefully applying induction on dimension to an ideal on  $\{x = 0\}$  gives order reduction (Encinas–Villamayor, Bierstone–Milman, Włodarczyk):

### Proposition (...N-T-W)

Let  $\mathcal{I}$  be an ideal on a logarithmically smooth  $Y$  with

$$\max_p \log \text{ord}_p(\mathcal{I}) = a.$$

## $\dim B = 0$ : sketch of argument

- In characteristic 0, if  $\log \text{ord}_p(\mathcal{I}) = a < \infty$ , then  $\mathcal{D}^{\leq a-1}\mathcal{I}$  contains an element  $x$  with derivative 1, a maximal contact element.
- Carefully applying induction on dimension to an ideal on  $\{x = 0\}$  gives order reduction (Encinas–Villamayor, Bierstone–Milman, Włodarczyk):

### Proposition (...N-T-W)

Let  $\mathcal{I}$  be an ideal on a logarithmically smooth  $Y$  with

$$\max_p \log \text{ord}_p(\mathcal{I}) = a.$$

There is a functorial logarithmic morphism  $Y_1 \rightarrow Y$ , with  $Y_1$  logarithmically smooth, such that  $\mathcal{I}\mathcal{O}_{Y'} = \mathcal{M} \cdot \mathcal{I}_1$  with  $\mathcal{M}$  an invertible monomial ideal and

$$\max_p \log \text{ord}_p(\mathcal{I}_1) < a.$$



Thank you for your attention!

## Adendum 1. Arbitrary $B$

(Work in progress)

### Main result (N-T-W)

Let  $Y \rightarrow B$  a logarithmically smooth morphism of logarithmically smooth schemes,  $\mathcal{I} \subset \mathcal{O}_Y$  an ideal. There is a log morphism  $B' \rightarrow B$  and functorial log morphism  $Y' \rightarrow Y$ , with  $Y' \rightarrow B'$  logarithmically smooth, and  $\mathcal{I}\mathcal{O}_{Y'}$  an invertible monomial ideal.

- This is done by **relative order reduction**, using **relative logarithmic derivatives**.

## Adendum 1. Arbitrary $B$

(Work in progress)

### Main result (N-T-W)

Let  $Y \rightarrow B$  a logarithmically smooth morphism of logarithmically smooth schemes,  $\mathcal{I} \subset \mathcal{O}_Y$  an ideal. There is a log morphism  $B' \rightarrow B$  and functorial log morphism  $Y' \rightarrow Y$ , with  $Y' \rightarrow B'$  logarithmically smooth, and  $\mathcal{I}\mathcal{O}_{Y'}$  an invertible monomial ideal.

- This is done by **relative order reduction**, using **relative logarithmic derivatives**.

### Definition

Write  $\mathcal{D}_{Y/B}^{\leq a}$  for the sheaf of **relative logarithmic differential operators** of order  $\leq a$ . The **relative logarithmic order** of an ideal  $\mathcal{I}$  is the minimum  $a$  such that  $\mathcal{D}_{Y/B}^{\leq a}\mathcal{I} = (1)$ .

## Adendum 1. The new step

- $\text{relord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M} := \mathcal{D}_{Y/B}^\infty \mathcal{I}$  is a nonunit ideal which is **monomial along the fibers**.
- Equivalently  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  is not the unit ideal.

## Adendum 1. The new step

- $\text{relord}_p(\mathcal{I}) = \infty$  if and only if  $\mathcal{M} := \mathcal{D}_{Y/B}^\infty \mathcal{I}$  is a nonunit ideal which is **monomial along the fibers**.
- Equivalently  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  is not the unit ideal.

### Monomialization Theorem [N-T-W]

Let  $Y \rightarrow B$  a logarithmically smooth morphism of logarithmically smooth schemes,  $\mathcal{M} \subset \mathcal{O}_Y$  an ideal with  $\mathcal{D}_{Y/B} \mathcal{M} = \mathcal{M}$ . There is a log morphism  $B' \rightarrow B$  with saturated pullback  $Y' \rightarrow B'$ , and  $\mathcal{M} \mathcal{O}_{Y'}$  a monomial ideal.

After this one can proceed as in the case “ $\dim B = 0$ ”.

## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .

## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .
- Once again the operators  $1, u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v}$  commute and have different eigenvalues on  $u^\alpha, v^\alpha$ .



## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .
- Once again the operators  $1, u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v}$  commute and have different eigenvalues on  $u^\alpha, v^\alpha$ .
- Expanding  $f = \sum u^\alpha f_\alpha + \sum v^\beta f_\beta$ , the condition  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  gives that only one term survives,

## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .
- Once again the operators  $1, u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v}$  commute and have different eigenvalues on  $u^\alpha, v^\alpha$ .
- Expanding  $f = \sum u^\alpha f_\alpha + \sum v^\beta f_\beta$ , the condition  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  gives that only one term survives,
- say  $f = u^\alpha f_\alpha$ , with  $f_\alpha \in \mathbb{C}[w]$ .

## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .
- Once again the operators  $1, u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v}$  commute and have different eigenvalues on  $u^\alpha, v^\alpha$ .
- Expanding  $f = \sum u^\alpha f_\alpha + \sum v^\beta f_\beta$ , the condition  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  gives that only one term survives,
- say  $f = u^\alpha f_\alpha$ , with  $f_\alpha \in \mathbb{C}[w]$ .
- Blowing up  $(f_\alpha)$  on  $B$  has the effect of making it monomial, so  $f$  becomes monomial.



## Adendum 1. Proof of Monomialization, special case

Let  $Y = \text{Spec } \mathbb{C}[u, v] \rightarrow B = \text{Spec } \mathbb{C}[w]$  with  $w = uv$ , and  $\mathcal{M} = (f)$ .

Proof in this special case.

- Every monomial is either  $u^\alpha w^k$  or  $v^\alpha w^k$ .
- Once again the operators  $1, u \frac{\partial}{\partial u} - v \frac{\partial}{\partial v}$  commute and have different eigenvalues on  $u^\alpha, v^\alpha$ .
- Expanding  $f = \sum u^\alpha f_\alpha + \sum v^\beta f_\beta$ , the condition  $\mathcal{M} = \mathcal{D}_{Y/B} \mathcal{M}$  gives that only one term survives,
- say  $f = u^\alpha f_\alpha$ , with  $f_\alpha \in \mathbb{C}[w]$ .
- Blowing up  $(f_\alpha)$  on  $B$  has the effect of making it monomial, so  $f$  becomes monomial.



The general case is surprisingly subtle.

## Adendum 1. In virtue of functoriality

### Theorem (Temkin)

*Resolution of singularities holds for excellent schemes, complex spaces, nonarchimedean spaces,  $p$ -adic spaces, formal spaces and for stacks.*

## Adendum 1. In virtue of functoriality

### Theorem (Temkin)

*Resolution of singularities holds for excellent schemes, complex spaces, nonarchimedean spaces,  $p$ -adic spaces, formal spaces and for stacks.*

- This is a consequence of resolution for varieties and schemes, **functorial** for smooth morphisms (submersions). Moreover

## Adendum 1. In virtue of functoriality


### Theorem (Temkin)

*Resolution of singularities holds for excellent schemes, complex spaces, nonarchimedean spaces,  $p$ -adic spaces, formal spaces and for stacks.*

- This is a consequence of resolution for varieties and schemes, **functorial** for smooth morphisms (submersions). Moreover
- Włodarczyk showed that if one seriously looks for a resolution **functor**, one is led to a resolution theorem.

## Adendum 2. Order reduction: Example 1

- Consider  $Y_1 = \text{Spec } \mathbb{C}[u, x]$  and  $D = \{u = 0\}$ .
- Let  $\mathcal{I} = (u^2, x^2)$ .
- If one blows up  $(u, x)$  the ideal is principalized:





## Adendum 2. Order reduction: Example 1

- Consider  $Y_1 = \text{Spec } \mathbb{C}[u, x]$  and  $D = \{u = 0\}$ .
- Let  $\mathcal{I} = (u^2, x^2)$ .
- If one blows up  $(u, x)$  the ideal is principalized:



## Adendum 2. Order reduction: Example 1

- Consider  $Y_1 = \text{Spec } \mathbb{C}[u, x]$  and  $D = \{u = 0\}$ .
- Let  $\mathcal{I} = (u^2, x^2)$ .
- If one blows up  $(u, x)$  the ideal is principalized:



- ▶ on the  $u$ -chart  $\text{Spec } \mathbb{C}[u, x']$  with  $x = x'u$  we have  $\mathcal{I}\mathcal{O}_{Y'_1} = (u^2)$ ,
- ▶ on the  $x$ -chart  $\text{Spec } \mathbb{C}[u', x]$  with  $u' = xu'$  we have  $\mathcal{I}\mathcal{O}_{Y'_1} = (x^2)$ ,
- ▶ which is exceptional hence monomial.

## Adendum 2. Order reduction: Example 1

- Consider  $Y_1 = \text{Spec } \mathbb{C}[u, x]$  and  $D = \{u = 0\}$ .
- Let  $\mathcal{I} = (u^2, x^2)$ .
- If one blows up  $(u, x)$  the ideal is principalized:



- ▶ on the  $u$ -chart  $\text{Spec } \mathbb{C}[u, x']$  with  $x = x'u$  we have  $\mathcal{IO}_{Y'_1} = (u^2)$ ,
  - ▶ on the  $x$ -chart  $\text{Spec } \mathbb{C}[u', x]$  with  $u' = xu'$  we have  $\mathcal{IO}_{Y'} = (x^2)$ ,
  - ▶ which is exceptional hence monomial.
- This is in fact the only functorial **admissible** blowing up.

## Adendum 2. Order reduction: Example 2

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$ .

## Adendum 2. Order reduction: Example 2

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$ .
- Example 1 is the pullback of this via  $v = u^2$ .
- Functoriality says: we need to blow up an ideal whose pullback is  $(u, x)$ .

## Adendum 2. Order reduction: Example 2

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$ .
- Example 1 is the pullback of this via  $v = u^2$ .
- Functoriality says: we need to blow up an ideal whose pullback is  $(u, x)$ .
- This means we need to blow up  $(v^{1/2}, x)$ .

## Adendum 2. Order reduction: Example 2

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$ .
- Example 1 is the pullback of this via  $v = u^2$ .
- Functoriality says: we need to blow up an ideal whose pullback is  $(u, x)$ .
- This means we need to blow up  $(v^{1/2}, x)$ .
- What is this? What is its blowup?

## Adendum 2. Kummer ideals

### Definition

- A **Kummer monomial** is a monomial in the Kummer-étale topology of  $Y$  (like  $v^{1/2}$ ).



## Adendum 2. Kummer ideals

### Definition

- A **Kummer monomial** is a monomial in the Kummer-étale topology of  $Y$  (like  $v^{1/2}$ ).
- A **Kummer monomial ideal** is a monomial ideal in the Kummer-étale topology of  $Y$ .

## Adendum 2. Kummer ideals

### Definition

- A **Kummer monomial** is a monomial in the Kummer-étale topology of  $Y$  (like  $v^{1/2}$ ).
- A **Kummer monomial ideal** is a monomial ideal in the Kummer-étale topology of  $Y$ .
- A **Kummer center** is the sum of a Kummer monomial ideal and the ideal of a log smooth subscheme.
- Locally  $(x_1, \dots, x_k, u_1^{1/d}, \dots, u_\ell^{1/d})$ .

## Adendum 2. Blowing up Kummer centers

### Proposition

*Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.*

## Adendum 2. Blowing up Kummer centers

### Proposition

*Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.*

### Example 0

$Y = \text{Spec } \mathbb{C}[v]$ , with toroidal structure associated to  $D = \{v = 0\}$ , and  $\mathcal{J} = (v^{1/2})$ .

## Adendum 2. Blowing up Kummer centers

### Proposition

*Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.*

### Example 0

$Y = \text{Spec } \mathbb{C}[v]$ , with toroidal structure associated to  $D = \{v = 0\}$ , and  $\mathcal{J} = (v^{1/2})$ .

- There is no log scheme  $Y'$  satisfying the proposition.

## Adendum 2. Blowing up Kummer centers

### Proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

### Example 0

$Y = \text{Spec } \mathbb{C}[v]$ , with toroidal structure associated to  $D = \{v = 0\}$ , and  $\mathcal{J} = (v^{1/2})$ .

- There is no log scheme  $Y'$  satisfying the proposition.
- There is a **stack**  $Y' = Y(\sqrt{D})$ , the **Cadman–Vistoli root stack**, satisfying the proposition!

## Adendum 2. Example 2 concluded

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$  and  $\mathcal{J} = (v^{1/2}, x)$ .

## Adendum 2. Example 2 concluded

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$  and  $\mathcal{J} = (v^{1/2}, x)$ .
- associated blowing up  $Y' \rightarrow Y_2$  with charts:
  - ▶  $Y'_x := \text{Spec } \mathbb{C}[v, x, v'] / (v'x^2 = v)$ , where  $v' = v/x^2$  (nonsingular scheme).
    - ★ Exceptional  $x = 0$ , now monomial.
    - ★  $\mathcal{I} = (v, x^2)$  transformed into  $(x^2)$ , invertible monomial ideal.
    - ★ Kummer ideal  $(v^{1/2}, x)$  transformed into **monomial ideal**  $(x)$ .



## Adendum 2. Example 2 concluded

- Consider  $Y_2 = \text{Spec } \mathbb{C}[v, x]$  and  $D = \{v = 0\}$ .
- Let  $\mathcal{I} = (v, x^2)$  and  $\mathcal{J} = (v^{1/2}, x)$ .
- associated blowing up  $Y' \rightarrow Y_2$  with charts:
  - ▶  $Y'_x := \text{Spec } \mathbb{C}[v, x, v'] / (v'x^2 = v)$ , where  $v' = v/x^2$  (nonsingular scheme).
    - ★ Exceptional  $x = 0$ , now monomial.
    - ★  $\mathcal{I} = (v, x^2)$  transformed into  $(x^2)$ , invertible monomial ideal.
    - ★ Kummer ideal  $(v^{1/2}, x)$  transformed into **monomial ideal**  $(x)$ .
  - ▶ The  $v^{1/2}$ -chart:
    - ★ stack quotient  $X'_{v^{1/2}} := [\text{Spec } \mathbb{C}[w, y] / \mu_2]$ ,
    - ★ where  $y = x/w$  and  $\mu_2 = \{\pm 1\}$  acts via  $(w, y) \mapsto (-w, -y)$ .
    - ★ Exceptional  $w = 0$  (monomial).
    - ★  $(v, x^2)$  transformed into invertible monomial ideal  $(v) = (w^2)$ .
    - ★  $(v^{1/2}, x)$  transformed into invertible monomial **ideal**  $(w)$ .

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

- There is a stack  $\tilde{Y}$  with coarse moduli space  $Y$  such that  $\tilde{\mathcal{J}} := \mathcal{J}\mathcal{O}_{\tilde{Y}}$  is an ideal.

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

- There is a stack  $\tilde{Y}$  with coarse moduli space  $Y$  such that  $\tilde{\mathcal{J}} := \mathcal{J}\mathcal{O}_{\tilde{Y}}$  is an ideal.
- Let  $\tilde{Y}' \rightarrow \tilde{Y}$  be the blowup of  $\tilde{\mathcal{J}}$  with exceptional  $E$ .

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

- There is a stack  $\tilde{Y}$  with coarse moduli space  $Y$  such that  $\tilde{\mathcal{J}} := \mathcal{J}\mathcal{O}_{\tilde{Y}}$  is an ideal.
- Let  $\tilde{Y}' \rightarrow \tilde{Y}$  be the blowup of  $\tilde{\mathcal{J}}$  with exceptional  $E$ .
- Let  $\tilde{Y}' \rightarrow B\mathbb{G}_m$  be the classifying morphism.

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

- There is a stack  $\tilde{Y}$  with coarse moduli space  $Y$  such that  $\tilde{\mathcal{J}} := \mathcal{J}\mathcal{O}_{\tilde{Y}}$  is an ideal.
- Let  $\tilde{Y}' \rightarrow \tilde{Y}$  be the blowup of  $\tilde{\mathcal{J}}$  with exceptional  $E$ .
- Let  $\tilde{Y}' \rightarrow B\mathbb{G}_m$  be the classifying morphism.
- $Y'$  is the relative coarse moduli space of  $\tilde{Y}' \rightarrow Y \times B\mathbb{G}_m$ .

## Adendum 2. Proof of proposition

Let  $\mathcal{J}$  be a Kummer center on a logarithmically smooth  $Y$ . There is a universal proper birational  $Y' \rightarrow Y$  such that  $Y'$  is logarithmically smooth and  $\mathcal{J}\mathcal{O}_{Y'}$  is an invertible ideal.

- There is a stack  $\tilde{Y}$  with coarse moduli space  $Y$  such that  $\tilde{\mathcal{J}} := \mathcal{J}\mathcal{O}_{\tilde{Y}}$  is an ideal.
- Let  $\tilde{Y}' \rightarrow \tilde{Y}$  be the blowup of  $\tilde{\mathcal{J}}$  with exceptional  $E$ .
- Let  $\tilde{Y}' \rightarrow B\mathbb{G}_m$  be the classifying morphism.
- $Y'$  is the relative coarse moduli space of  $\tilde{Y}' \rightarrow Y \times B\mathbb{G}_m$ .
- One shows this is independent of choices. ♠