

Stable varieties with a twist

Dan Abramovich and Brendan Hassett *

Abstract. We describe a new approach to the definition of the moduli functor of stable varieties. While there is wide agreement as to what classes of varieties should appear, the notion of a family of stable surfaces is quite subtle, as key numerical invariants may fail to be constant in flat families. Our approach is to add natural stack structure to stable varieties. For example, given a canonical model we take the global-quotient stack structure arising from its realization as Proj of the canonical ring. Deformations of the stack structure preserve key numerical invariants of the stable variety, including the top self-intersection of the canonical divisor. This approach yields a transparent construction of the moduli stack of stable varieties as a global quotient of a suitable Hilbert scheme of weighted projective stacks.

2010 Mathematics Subject Classification. Primary 14J10; Secondary 14D22, 14E30.

Keywords. moduli spaces, Gorenstein singularities, varieties of general type, quotient stacks

Contents

1	Introduction	1
2	Cyclotomic stacks and weighted projective stacks	6
3	Moduli of stacks with polarizing line bundles	15
4	Moduli of polarized orbispaces	20
5	Kollár families and stacks	24
6	Semilog canonical singularities and compactifications	29
A	The semilog canonical locus is open	33

1. Introduction

1.1. Moduli of stable varieties: the case of surfaces. In the paper [KSB88], Kollár and Shepherd-Barron introduced *stable surfaces* as a generalization of sta-

*Research of D.A. partially supported by NSF grants DMS-0335501 and DMS-0603284. Research of B.H. partially supported by NSF grants DMS-0196187, DMS-0134259, and DMS-0554491 and by the Sloan Foundation.

ble curves. This class is natural from the point of view of the minimal model program, which shows that any one-parameter family of surfaces of general type admits a unique stable limit. Indeed, the stable reduction process of Deligne and Mumford can be interpreted using the language of minimal models of surfaces. Stable surfaces admit *semilog canonical singularities*. Codimension-one semilog canonical singularities are nodes, but in codimension two more complicated singularities arise. However, these singularities are all reduced, satisfy Serre’s condition S_2 (hence are Cohen–Macaulay in codimension 2), and admit a well-defined \mathbb{Q} -Cartier canonical divisor K_S . Stable varieties in any dimension are similarly defined as proper varieties with semilog canonical singularities and ample canonical divisor.

Kollár and Shepherd-Barron proposed the moduli space of stable surfaces as the natural compactification of the moduli space of surfaces of general type. Much of this was established in [KSB88], which also offered a detailed classification of the singularities that arise. Boundedness of the class of stable surfaces with fixed invariant $(K_S)^2$ was shown by Alexeev [Ale94].

1.2. The problem of families. To a great extent this established the moduli space of stable surfaces as the ‘right’ geometric compactification of moduli of surfaces of general type. But one nagging question remains unresolved: *what flat families $\pi : X \rightarrow B$ of stable surfaces should be admitted in the compactification?*

The problem stems precisely from the fact that the dualizing sheaf $\omega_S = \mathcal{O}_S(K_S)$ is not necessarily invertible for a stable surface. Rather, for a suitable positive integer n depending on S , the sheaf

$$\mathcal{O}_S(nK_S) = \omega_S^{[n]} := (\omega_S^n)^{**}$$

is invertible. As it turns out, there exist flat families $\pi : X \rightarrow B$ over a smooth curve where the fibers are stable surfaces but $\omega_\pi^{[n]}$ is not locally free for any $n \neq 0$. For a local example, see Pinkham’s analysis of the deformation space of the cone over a rational normal quartic curve described in [KSB88, Example 2.8].

Two approaches were suggested to resolve this issue (see [HK04] for detailed discussion). In [Kol90, 5.2], Kollár proposed what has come to be called a *Kollár family* of stable surfaces, where the formation of *every* saturated power of ω_π is required to commute with base change. Viehweg [Vie95] suggested using families where *some* saturated power of ω_π is invertible (and in particular, commutes with base extension), i.e., there exists an integer $n > 0$ and an invertible sheaf on X restricting to $\omega_S^{[n]}$ on each fiber S of π .

It has been shown in [HK04] that Viehweg families give rise to a good moduli space. The purpose of this article is to address the case of Kollár families. While this question has been considered before (cf. [Hac04]), the solution we propose is perhaps more natural than previous approaches, and has implications beyond the question at hand.

We must note that Kollár’s [Kol08] resolves the issue by tackling it head on using the moduli spaces of Husks. Our approach goes by way of showing that, in the situation at hand, if we consider objects with an appropriate stack structure,

the problem does not arise. This does require us to show that moduli spaces of such stack structures are well behaved.

1.3. Canonical models and quotient stacks. Here is the basic idea: Let X be a smooth projective variety of general type. A fundamental invariant of X is the *canonical ring*

$$R(X) := \bigoplus_{n \geq 0} \Gamma(X, \omega_X^n),$$

a graded ring built up from the differential forms on X . This ring has long been known to be a birational invariant of X and has recently been shown to be finitely generated [BCHM10]. This was known classically for curves and surfaces, and established for threefolds by S. Mori in the 1980's. The associated projective variety

$$X^{\text{can}} := \text{Proj } R(X) = (\text{Spec } R(X) \setminus 0) / \mathbb{G}_m$$

is called the *canonical model* of X . Some saturated power $\omega_{X^{\text{can}}}^{[n]}$ is ample and invertible, essentially by construction. Therefore, deformations of canonical models (and stable surfaces) are most naturally expressed via deformations of the canonical ring.

In these terms, our main problem is that certain deformations of a stable surface do not yield deformations of its canonical ring. Our solution is to replace X^{can} with a more sophisticated geometric object that remembers its algebraic origins. Precisely, we consider quotient *stacks*

$$\mathcal{X}^{\text{can}} = \mathbf{Proj} R(X) := [(\text{Spec } R(X) \setminus 0) / \mathbb{G}_m],$$

which admit nontrivial stack-structure at precisely the points where $\omega_{X^{\text{can}}}$ fails to be invertible. Our main goal is to show that this formulation is equivalent to Kollár's, and yields a workable moduli space.

1.4. Moduli of stacks: the past. The idea that families with fibers having stack structure should admit workable moduli spaces is not new. On the level of deformations, Hacking's approach in [Hac04] uses precisely this idea. The case of *stable fibered surfaces*, namely stable surfaces fibered over curves with semistable fibers, was discussed in [AV00]. It was generalized to many moduli problems involving curves in [AV02] and to varieties of higher dimension with suitable "plurifibration" structures [Abr02]. The fibered and plurifibered examples are a little misleading, since the existence of canonical models and complete moduli in those cases are an outcome of [AV02] and do not require the minimal model program. Still they do reinforce the point that stacks do not pose a true obstacle in defining moduli.

We show in this paper that this is indeed the case for stacks of the form $\mathcal{X} = \mathbf{Proj} R$ for R a finitely generated graded ring, and moreover this works in exactly the same way it works for varieties, if one takes the appropriate viewpoint. Classical theory of moduli of projective varieties starts by considering varieties embedded in \mathbb{P}^n , forming Hilbert schemes, taking quotient stacks of the appropriate loci of normally embedded varieties by the action of $\mathbf{PGL}(n+1)$, and taking the union

of all the resulting stacks over all types of embeddings. Then one tries to carve out pieces of this “mother of all moduli spaces” (of varieties) which are bounded, and if lucky, proper, by fixing appropriate numerical invariants.

Miles Reid has been advocating over the years that, while most people insist on embedding projective varieties in projective space, the varieties really beg to be embedded in *weighted* projective spaces, denoted $\mathbb{P}(\rho)$ below. This is simply because graded rings are in general not generated in one degree; see [Rei78], [Rei97], and especially [Rei02]. For our stacks this point is absolutely essential, and in fact our stacks (as well as varieties) are embedded in the appropriate weighted projective *stacks*, denoted $\mathcal{P}(\rho)$ below.

1.5. Moduli of stacks: this paper. In section 2 we consider weighted projective stacks and their substacks. Following [OS03] substacks are parametrized by a Hilbert scheme. All substacks of $\mathcal{P}(\rho)$ are cyclotomic, in particular tame, and in analogy with the scheme case, we characterize in Proposition 2.4.2 pairs $(\mathcal{X}, \mathcal{L})$ consisting of a cyclotomic stack with a *polarizing line bundle* - one affording an embedding - in cohomological terms.

Similar results are obtained in [RT09], though the emphasis is different. Ross and Thomas embed smooth orbifolds in weighted projective stacks to study orbifold constant scalar curvature Kähler metrics. They develop a notion of Geometric Invariant Theory stability for polarized orbifolds and use the relation between stability and moment maps to formulate obstructions to existence of these metrics.

In section 3 we use these results to construct an algebraic stack $Sta^{\mathcal{L}}$ parametrizing pairs $(\mathcal{X}, \mathcal{L})$ of a proper stack with polarizing line bundle \mathcal{L} (see Theorem 3.1.4). This involves studying the effect of changing the range of weights on the moduli of normally embedded substacks (see Proposition 3.2.4).

In section 4 we introduce *orbispaces*, discuss the difference between *polarizing line bundle* and *polarization*, and construct algebraic stacks $Orb^{\mathcal{L}}$ and Orb^{λ} parametrizing *orbispaces with polarizing line bundles* and *polarized orbispaces*, respectively (see Proposition 4.2.1). We further discuss the canonical polarization, and the algebraic stack Orb^{ω} of canonically polarized orbispaces, in Theorem 4.4.4.

This ends the general discussion of moduli of stacks, and brings us to the questions related to Kollár families. We begin Section 5 by discussing Kollár families of \mathbb{Q} -line bundles (see Definition 5.2.1). We then define *uniformized twisted varieties* in Definition 5.3.4. An important point here is that *the conditions on flat families are entirely on geometric fibers*.

Finally we relate the two notions and prove an equivalence of categories between Kollár families $(X \rightarrow B, F)$ of \mathbb{Q} -line bundles and families of uniformized twisted varieties (see Theorem 5.3.6). In particular this means that we have an algebraic stack $\mathcal{K}^{\mathcal{L}}$ of Kollár families with a polarizing \mathbb{Q} -line bundle, which is at the same time the stack of twisted varieties with polarizing line bundles. The rigidification \mathcal{K}^{λ} of $\mathcal{K}^{\mathcal{L}}$ is naturally the stack of polarized twisted varieties. Passing to the locally closed substack where the initial homology group of the relative dualizing complex is invertible and polarizing, we obtain the substack \mathcal{K}^{ω} of *canonically polarized twisted varieties*.

We end the paper by applying the previous discussion to moduli of stable varieties in Section 6. We define the moduli functor in terms of families of twisted stable varieties and equivalently Kollár families of stable varieties. We show that this is open in \mathcal{K}^ω (see Proposition A.1.1) and discuss what is known about properness in section 6.1.

It may be of interest to develop a treatment of the *steps* of the minimal model program, and not only the end product $\mathbf{Proj}R(X)$, using stacks. It would require at least an understanding of positivity properties for the birational “contraction” $\mathcal{X} \rightarrow X$ from a twisted variety to its coarse moduli space, where we expect the notion of *age*, introduced in [IR96], to be salient. This however goes beyond the scope of the present paper.

1.6. Acknowledgments. We are grateful to Valery Alexeev, Kai Behrend, Tim Cochran, Alessio Corti, Barbara Fantechi, János Kollár, Sándor Kovács, Martin Olsson, Miles Reid, Julius Ross, Richard Thomas, Eckart Viehweg, and Angelo Vistoli for helpful conversations about these questions. We appreciate the hospitality of the Mathematisches Forschungsinstitut Oberwolfach and the Mathematical Sciences Research Institute in Berkeley, California.

1.7. Conventions. Through most of this paper we work over \mathbb{Z} . When we refer to schemes of finite type, we mean schemes of finite type over \mathbb{Z} . The main exceptions are Section 6 and the Appendix, where we assume characteristic zero. We leave it to the interested reader to supply the incantations needed to extend our results to an arbitrary base scheme.

In characteristic zero, we freely use the established literature for Deligne-Mumford stacks. In positive and mixed characteristics, we rely on the recent paper [AOV08a], which develops a new notion of ‘tame stacks’ with nice properties, e.g., they admit coarse moduli spaces that behave well under base extension.

1.8. Notation.

$\mathbb{P}(\rho)$	Weighted projective space of \mathbb{G}_m -representation ρ
$\mathcal{P}(\rho)$	Weighted projective <i>stack</i> of \mathbb{G}_m -representation ρ
$\mathcal{Sta}^{\mathcal{L}}$	Stack of <i>stacks</i> with polarizing line bundle
$\mathcal{Orb}^{\mathcal{L}}$	Stack of <i>orbispaces</i> with polarizing line bundle
\mathcal{Orb}^λ	– with polarization
\mathcal{Orb}^ω	– with canonical polarization
$\mathcal{K}^{\mathcal{L}}$	Stack of <i>twisted varieties</i> with polarizing line bundle
\mathcal{K}^λ	– with polarization
\mathcal{K}^ω	– with canonical polarization
\square_{cm}	Index indicating Cohen-Macaulay fibers
\square_{gor}	– Gorenstein fibers
\square_{can}	– canonical singularities (in the coarse fibers)
\square_{sic}	– semilog canonical singularities fibers
$\square_{\text{gor-slc}}$	– Gorenstein s.l.c. singularities (in the twisted fibers)
\square^{can}	indicating canonical model of something

2. Cyclotomic stacks and weighted projective stacks

2.1. Weighted projective stacks.

Definition 2.1.1. Fix a nondecreasing sequence of positive integer weights

$$0 < \rho_1 \leq \dots \leq \rho_r$$

and consider the associated linear action of \mathbb{G}_m on affine space

$$\begin{aligned} \rho : \mathbb{G}_m \times \mathbb{A}^r &\rightarrow \mathbb{A}^r \\ \rho^*(x_1, \dots, x_r) &= (t^{\rho_1}x_1, \dots, t^{\rho_r}x_r). \end{aligned}$$

Recall that *weighted projective space* associated to the representation ρ is defined as the quotient scheme

$$\mathbb{P}(\rho) = \mathbb{P}(\rho_1, \dots, \rho_r) = (\mathbb{A}^r \setminus 0) / \mathbb{G}_m.$$

We define the *weighted projective stack* as the quotient stack

$$\mathcal{P}(\rho) = \mathcal{P}(\rho_1, \dots, \rho_r) = [(\mathbb{A}^r \setminus 0) / \mathbb{G}_m].$$

Each weighted projective stack has the corresponding weighted projective space as its coarse moduli space. For each positive integer d we have

$$\mathbb{P}(d\rho_1, \dots, d\rho_r) \simeq \mathbb{P}(\rho_1, \dots, \rho_r),$$

however, the weighted projective stacks are not isomorphic unless $d = 1$. Indeed $\mathbb{P}(d\rho_1, \dots, d\rho_r) \rightarrow \mathbb{P}(\rho_1, \dots, \rho_r)$ is a gerbe banded by μ_d . Note also that a weighted projective stack is representable if and only if each weight $\rho_i = 1$; it contains a nonempty representable open subset if and only if $\gcd(\rho_1, \dots, \rho_r) = 1$.

Definition 2.1.2. Let B be a scheme of finite type and V a locally-free \mathcal{O}_B -module of rank r . Suppose that \mathbb{G}_m acts \mathcal{O}_B -linearly on V with positive weights (ρ_1, \dots, ρ_r) . Let $w_1 < \dots < w_m$ denote the distinct weights that occur, so we have a decomposition of \mathcal{O}_B -modules

$$V \simeq V_{w_1} \oplus \dots \oplus V_{w_m}$$

with

$$\rho(v) = t^{w_i}v, \quad v \in V_{w_i}.$$

Write $\mathbb{A} := \mathrm{Spec}_B(\mathrm{Sym}_B^\bullet V)$ for the associated vector bundle, with its zero section $0 \subset \mathbb{A}$, and

$$\rho : \mathbb{G}_m \times_B \mathbb{A} \rightarrow \mathbb{A}$$

for the associated group action. The *weighted projective stack* associated with V and ρ is the quotient stack

$$\mathcal{P}(V, \rho) := [(\mathbb{A} \setminus 0) / \mathbb{G}_m] \rightarrow B.$$

We will sometimes drop V from the notation when the meaning is unambiguous.

Graded coherent $\mathcal{O}_{\mathbb{A}}$ -modules descend to coherent sheaves on $\mathcal{P}(V, \rho)$; thus for each integer w , the twist $\mathcal{O}_{\mathbb{A}}(w)$ yields an *invertible* sheaf $\mathcal{O}_{\mathcal{P}(V, \rho)}(w)$ on $\mathcal{P}(V, \rho)$. The canonical graded homomorphisms

$$V_{w_i} \otimes_B \mathcal{O}_{\mathbb{A}} \rightarrow \mathcal{O}_{\mathbb{A}}(w_i)$$

induce

$$\phi_{w_i} : V_{w_i} \otimes_B \mathcal{O}_{\mathcal{P}(V, \rho)} \rightarrow \mathcal{O}_{\mathcal{P}(V, \rho)}(w_i), \quad i = 1, \dots, m.$$

Note that ϕ_{w_i} vanishes along the locus where the elements of V_{w_i} are simultaneously zero, i.e., where all the weight- w_i coordinates simultaneously vanish. The homomorphisms $\phi_{w_i}, i = 1, \dots, m$ do not vanish simultaneously anywhere.

The following lemma is well-known in the case of projective space. The proof for weighted-projective stacks is identical:

Lemma 2.1.3. *The stack $\mathcal{P}(V, \rho)$ is equivalent, as a category fibered over the category of B -schemes, with the following category:*

- (1) *Objects over a scheme T consist of*
 - (a) *a line bundle L over T , and*
 - (b) *\mathcal{O}_T -linear homomorphisms*

$$\phi_{w_i} : V_{w_i} \otimes_B \mathcal{O}_T \rightarrow L^{w_i}, \quad i = 1, \dots, m,$$

not vanishing simultaneously at any point of T .

- (2) *Arrows consist of fiber diagrams*

$$\begin{array}{ccc} L' & \rightarrow & L \\ \downarrow & & \downarrow \\ T' & \rightarrow & T \end{array}$$

compatible with the homomorphisms ϕ_{w_i} and ϕ'_{w_i} .

As we shall see, certain stacks admitting “uniformizing line bundles” have representable morphisms into weighted projective stacks.

2.2. Closed substacks of weighted projective stacks. By descent theory, closed substacks

$$\mathcal{X} \subset \mathcal{P}(\rho) = \mathcal{P}(\rho_1, \dots, \rho_r)$$

correspond to closed subschemes

$$\mathcal{X} \times_{\mathcal{P}(\rho)} (\mathbb{A}^r \setminus 0) \subset (\mathbb{A}^r \setminus 0),$$

equivariant under the action of \mathbb{G}_m . Let $C\mathcal{X}$ denote the closure of this scheme in \mathbb{A}^r , i.e., the cone over \mathcal{X} . Each such subscheme over a field k can be defined by a graded ideal

$$I \subset k[x_1, \dots, x_r].$$

The *saturation* of such an ideal is defined as

$$\{f \in k[x_1, \dots, x_r] : \langle x_1, \dots, x_r \rangle^d f \subset I \text{ for some } d \gg 0\}$$

and two graded ideals are *equivalent* if they have the same saturation. Equivalent graded ideals give identical ideal sheaves on $\mathbb{A}^r \setminus 0$, hence they define the same equivariant subschemes of $\mathbb{A}^r \setminus 0$ corresponding to the same substack of $\mathcal{P}(\rho)$.

There is one crucial distinction from the standard theory of projective varieties: when the weights $\rho_i = 1$ for all i , every graded ideal is equivalent to an ideal generated by elements in a single degree. This can fail when the weights are not all equal. However, if we set $N = \text{lcm}(\rho_1, \dots, \rho_r)$ then every graded ideal is equivalent to one with generators in degrees $n, n+1, \dots, n+N-1$, for n sufficiently large. (This can be shown by regarding graded ideals as modules over the subring of $k[x_1, \dots, x_r]$ generated by monomials of weights divisible by N .) An equivariant subscheme $C\mathcal{X} \subset \mathbb{A}^r$ is specified by the induced quotient homomorphisms on the graded pieces of the coordinate rings

$$k[x_1, \dots, x_r]_d \twoheadrightarrow k[C\mathcal{X}]_d, \quad d = n, n+1, \dots, n+N-1.$$

In sheaf-theoretic terms, $\mathcal{O}_{\mathcal{P}(\rho)}$ -module quotients

$$\mathcal{O}_{\mathcal{P}(\rho)} \twoheadrightarrow \mathcal{O}_{\mathcal{X}}$$

are determined by the induced quotient of $\mathcal{O}_{\mathbb{P}(\rho)}$ -modules

$$\pi_*[\oplus_{i=0}^{N-1} \mathcal{O}_{\mathcal{P}(\rho)}(i)] \twoheadrightarrow \pi_*[\oplus_{i=0}^{N-1} \mathcal{O}_{\mathcal{X}}(i)],$$

or any other interval of twists of length N , where $\pi : \mathcal{P}(\rho) \rightarrow \mathbb{P}(\rho)$ is the coarse moduli morphism.

Definition 2.2.1. Fix a representation (V, ρ) of \mathbb{G}_m with isotypical components V_{w_1}, \dots, V_{w_m} of positive weights $w_1 < \dots < w_m$.

A closed substack $\mathcal{X} \subset \mathcal{P}(\rho)$ is said to be *normally embedded* if

- (1) for each $1 \leq j \leq m$, the homomorphism $V_{w_j} \rightarrow H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(w_j))$ is an isomorphism,
- (2) for each $k \geq w_1$ and $i \geq 1$ we have $H^i(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(k)) = 0$, and
- (3) the natural homomorphism

$$\text{Sym } V \longrightarrow \oplus_{k \geq 0} H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(k))$$

surjects onto the component $H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(k))$ for each $k \geq w_1$.

We proceed, following Olsson–Starr [OS03] to classify substacks $\mathcal{X} \subset \mathcal{P}(\rho)$ by a Hilbert scheme. The following definition is [OS03, Definition 5.1].

Definition 2.2.2. A locally free sheaf \mathcal{E} on a stack \mathcal{Y} is *generating* if, for each quasi-coherent $\mathcal{O}_{\mathcal{Y}}$ -module \mathcal{F} , the natural homomorphism

$$\theta(\mathcal{F}) : \pi^* \pi_* \mathcal{H}om_{\mathcal{Y}}(\mathcal{E}, \mathcal{F}) \otimes_{\mathcal{Y}} \mathcal{E} \rightarrow \mathcal{F}$$

is surjective.

We note that

$$\mathcal{E} = \bigoplus_{i=0}^{N-1} \mathcal{O}_{\mathcal{P}(\rho)}(-i)$$

is a generating sheaf for $\mathcal{P}(\rho)$ where $N = \text{lcm}(w_1, \dots, w_m)$. The main insight of [OS03] §6 is to reduce the study of $\mathcal{O}_{\mathcal{P}(\rho)}$ -quotients of \mathcal{F} to an analysis of $\mathcal{O}_{\mathbb{P}(\rho)}$ -quotients of $\pi_* \mathcal{H}om_{\mathcal{P}(\rho)}(\mathcal{E}, \mathcal{F})$. The following is a direct application of [OS03, Theorem 1.5] to weighted projective stacks.

Theorem 2.2.3. *Let ρ be an action of \mathbb{G}_m on \mathbb{A}^r with positive weights and $\mathcal{P}(\rho)$ the resulting weighted projective stack. For each scheme T and stack \mathcal{Y} , write $\mathcal{Y}_T = \mathcal{Y} \times T$. Consider the functor*

$$\mathcal{H}ilb : \mathbb{Z}\text{-schemes} \rightarrow \text{Sets}$$

with

$$\begin{aligned} \mathcal{H}ilb_{\mathcal{P}(\rho)}(T) &= \left\{ \begin{array}{l} \text{Isomorphism classes of } \mathcal{O}_{\mathcal{P}(\rho)_T}\text{-linear quo-} \\ \text{tients } \mathcal{O}_{\mathcal{P}(\rho)_T} \twoheadrightarrow \mathcal{Q}, \text{ with } \mathcal{Q} \text{ flat over } T \end{array} \right\} \\ &= \left\{ \begin{array}{l} \text{Closed substacks } \mathcal{X} \subset \mathcal{P}(\rho)_T \text{ over } T, \text{ with } \mathcal{X} \\ \text{flat over } T \end{array} \right\}. \end{aligned}$$

Then $\mathcal{H}ilb_{\mathcal{P}(\rho)}$ is represented by a scheme (denoted also $\mathcal{H}ilb_{\mathcal{P}(\rho)}$).

Furthermore, for each function $\mathfrak{F} : \mathbb{Z} \rightarrow \mathbb{Z}$, the subfunctor consisting of closed substacks $\mathcal{X} \subset \mathcal{P}(\rho)_T$ with Hilbert-Euler characteristic

$$\chi(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}(m)) = \mathfrak{F}(m)$$

for all geometric points $t \in T$ is represented by a projective scheme $\mathcal{H}ilb_{\mathcal{P}(\rho), \mathfrak{F}}$ which is a finite union of connected components of $\mathcal{H}ilb_{\mathcal{P}(\rho)}$.

2.3. Cyclotomic stacks and line bundles. Given an algebraic stack \mathcal{X} , the *inertia stack* is an algebraic stack whose objects are pairs (ξ, σ) , with ξ an object of \mathcal{X} and $\sigma \in \text{Aut}(\xi)$. The morphism $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$ given by $(\xi, \sigma) \mapsto \xi$ is representable, and $\mathcal{I}_{\mathcal{X}}$ is a group-scheme over \mathcal{X} . If \mathcal{X} is separated, the morphism $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$ is proper.

Definition 2.3.1. A flat separated algebraic stack

$$f : \mathcal{X} \rightarrow B$$

locally of finite presentation over a scheme B is said to be a *cyclotomic stack* if it has cyclotomic stabilizers, i.e. if each geometric fiber of $\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$ is isomorphic to the finite group scheme μ_n for some n .

Keeping \mathcal{X} separated and with finite diagonal, let $\pi : \mathcal{X} \rightarrow X$ denote the coarse-moduli-space morphism; it is proper and quasi-finite [KM97]. The resulting family of coarse moduli spaces is denoted $\bar{f} : X \rightarrow B$.

Example 2.3.2. Let $Y \rightarrow B$ be an algebraic space, flat and separated over B . If \mathbb{G}_m acts properly on Y over B with finite stabilizers then the quotient stack $\mathcal{X} := [Y/\mathbb{G}_m] \rightarrow B$ is cyclotomic.

Indeed, the stabilizer scheme

$$Z = \{(y, h) : h \cdot y = y\} \subset Y \times_B \mathbb{G}_m$$

is equivariant with \mathbb{G}_m -action on $Y \times_B \mathbb{G}_m$

$$g \cdot (y, h) = (gy, ghg^{-1}) = (gy, h).$$

On taking quotients, we obtain

$$\mathcal{I}_{\mathcal{X}} = [Z/\mathbb{G}_m],$$

which is a closed substack of $[(Y \times_B \mathbb{G}_m)/\mathbb{G}_m] = [\mathcal{X} \times_B \mathbb{G}_m]$.

Definition 2.3.3. We say that a stack \mathcal{X} has index N if for each object $\xi \in \mathcal{X}(T)$ over a scheme T , and each automorphism $a \in \text{Aut } \xi$ we have $a^N = id$, and if N is the minimal positive integer satisfying this condition.

Lemma 2.3.4. *Let*

$$f : \mathcal{X} \rightarrow B$$

be a stack of finite presentation over a scheme B and having finite diagonal. There is a positive integer N such that \mathcal{X} has index N .

Proof. This is well known: we may assume \mathcal{X} is of finite type over a field k . Then the inertia stack $\mathcal{I}_{\mathcal{X}}$ is of finite type as well. The projection morphism

$$\mathcal{I}_{\mathcal{X}} \rightarrow \mathcal{X}$$

is finite, with fibers the automorphism group-schemes of the corresponding objects of \mathcal{X} . The degree is therefore bounded. \square

We note that cyclotomic stacks are tame [AOV08a, Theorem 3.2 (b)]. The following lemma is a special case of [AV02, Lemma 2.2.3] in the Deligne–Mumford case, and [AOV08a, Theorem 3.2 (d)] in general:

Lemma 2.3.5. *Consider a cyclotomic stack*

$$f : \mathcal{X} \rightarrow B$$

and a geometric closed point $\xi : \text{Spec } K \rightarrow \mathcal{X}$ with stabilizer μ_n . Let $\bar{\xi} : \text{Spec } K \rightarrow X$ be the corresponding point. Then, in a suitable étale neighborhood of $\bar{\xi}$, the stack \mathcal{X} is a quotient of an affine scheme by an action of μ_n .

The following is a special case of [AV02, Lemma 2.2.3] in the tame Deligne–Mumford case, and [AOV08a, Corollary 3.3] in general.

Lemma 2.3.6. *Let $f : \mathcal{X} \rightarrow B$ be a cyclotomic stack (assumed by definition flat).*

- (1) *The coarse moduli space $\bar{f} : X \rightarrow B$ is flat.*
- (2) *Formation of coarse moduli spaces commutes with base extension, i.e., for any morphism of schemes $X' \rightarrow X$, the coarse moduli space of $\mathcal{X} \times_X X'$ is X' . In particular, the geometric fibers of \bar{f} are coarse moduli spaces of the corresponding fibers of f .*

Lemma 2.3.7. *Let $\mathcal{X} \rightarrow B$ be a cyclotomic stack of index N and \mathcal{L} an invertible sheaf on \mathcal{X} . Then there is an invertible sheaf M on X and an isomorphism*

$$\mathcal{L}^N \simeq \pi^* M.$$

Proof. Each automorphism group acts trivially on the fibers of \mathcal{L}^N and hence trivially on the sheaf \mathcal{L}^N . Write \mathcal{X} locally near a geometric point $\bar{\xi}$ of X as $[U/\mu_r]$ with $r|N$; assume \mathcal{L} is trivial on U . Then the total space of \mathcal{L}^N is the quotient $[(U \times \mathbb{A}^1)/\mu_r]$, but the action on the factor \mathbb{A}^1 is trivial, and the total space is $[U/\mu_r] \times \mathbb{A}^1 = \mathcal{X} \times \mathbb{A}^1$. Its coarse moduli space in this neighborhood is $X \times \mathbb{A}^1$. It follows that $M = \pi_* \mathcal{L}^N$ is an invertible sheaf on the coarse moduli space and $\pi^* M \rightarrow \mathcal{L}$ is an isomorphism. See also [AOV08b, Lemma 2.9] \square

Definition 2.3.8. Let \mathcal{L} be an invertible sheaf over a stack \mathcal{X} . The stack

$$\mathcal{P}_{\mathcal{L}} := \mathrm{Spec}_{\mathcal{X}} \left(\bigoplus_{i \in \mathbb{Z}} \mathcal{L}^i \right) \rightarrow \mathcal{X}$$

is called the *principal bundle* of \mathcal{L} . The grading induces a \mathbb{G}_m -action on $\mathcal{P}_{\mathcal{L}}$ over \mathcal{X} , which gives $\mathcal{P}_{\mathcal{L}}$ the structure of a \mathbb{G}_m -principal bundle over \mathcal{X} .

This definition requires the spectrum of a finitely-generated quasi-coherent sheaf of algebras, whose existence is guaranteed by [LMB00] §14.2.

We shall require the following, see [AV02, Lemma 4.4.3] or [LMB00].

Lemma 2.3.9. *Let \mathcal{X} and \mathcal{Y} be algebraic stacks. Let $g : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism. The following are equivalent:*

- (1) *g is representable;*
- (2) *for any algebraically closed field K and geometric point $\xi : \mathrm{Spec}(K) \rightarrow \mathcal{X}$, the natural homomorphism of group schemes*

$$\mathrm{Aut} \xi \rightarrow \mathrm{Aut} g(\xi)$$

is a monomorphism.

Proof. Recall that an algebraic space is a stack with trivial stabilizers. Also note that if V is an algebraic space and $\phi : V \rightarrow \mathcal{Y}$ a morphism, then the automorphisms of a point ξ of $\mathcal{X} \times_{\mathcal{Y}} V$ are precisely the kernel of the map $\text{Aut } \phi(\xi) \rightarrow \text{Aut } g(\phi(\xi))$. So if the condition on automorphisms holds then $\mathcal{X} \times_{\mathcal{Y}} V$ has trivial stabilizers and therefore g representable. Conversely, if g is representable and $V \rightarrow \mathcal{Y}$ is surjective then $\mathcal{X} \times_{\mathcal{Y}} V$ is representable and $\mathcal{X} \times_{\mathcal{Y}} V \rightarrow \mathcal{X}$ surjective. So since $\mathcal{X} \times_{\mathcal{Y}} V$ has trivial stabilizers we have $\text{Aut } \phi(\xi) \rightarrow \text{Aut } g(\phi(\xi))$ injective for all ξ . \square

We can now state a result describing generating sheaves coming from powers of a line bundle.

Proposition 2.3.10. *Let $\mathcal{X} \rightarrow B$ be a cyclotomic stack of index N and \mathcal{L} an invertible sheaf on \mathcal{X} . The following conditions are equivalent.*

- (1) $\mathcal{P}_{\mathcal{L}}$ is representable;
- (2) the classifying morphism $\mathcal{X} \rightarrow \mathcal{B}\mathbb{G}_m$ associated to $\mathcal{P}_{\mathcal{L}}$ is representable;
- (3) $\bigoplus_{i=0}^{N-1} \mathcal{L}^{\otimes -i}$ is a generating sheaf for \mathcal{X} .
- (4) for each geometric point $\xi : \text{Spec } K \rightarrow \mathcal{X}$ the action of $\text{Aut } \xi$ on the fiber of \mathcal{L} is effective;

Definition 2.3.11. When any of the conditions in the proposition is satisfied, $\mathcal{X} \rightarrow B$ is said to be *uniformized by \mathcal{L}* , and \mathcal{L} is a *uniformizing line bundle* for $\mathcal{X} \rightarrow B$.

Proof of Proposition. We first show that the first two conditions are equivalent. The representable morphism

$$\text{Spec } \mathbb{Z} \rightarrow \mathcal{B}\mathbb{G}_m$$

is the universal principal bundle, thus

$$\mathcal{P}_{\mathcal{L}} = \mathcal{X} \times_{\mathcal{B}\mathbb{G}_m} \text{Spec } \mathbb{Z}.$$

Clearly if $\mathcal{X} \rightarrow \mathcal{B}\mathbb{G}_m$ is representable, so is $\mathcal{P}_{\mathcal{L}}$. Conversely, if $\mathcal{P}_{\mathcal{L}}$ is representable, the automorphism group of any geometric object $\text{Spec } K \rightarrow \mathcal{X}$ of \mathcal{X} acts on the \mathbb{G}_m -torsor $\text{Spec } K \times_{\mathcal{X}} \mathcal{P}_{\mathcal{L}}$ with trivial stabilizers; in particular, it acts effectively. This means that the morphism $\mathcal{X} \rightarrow \mathcal{B}\mathbb{G}_m$ induces a monomorphism on automorphism groups, and by Lemma 2.3.9 it is representable.

Recall from [OS03], Proposition 5.2 that a sheaf \mathcal{F} is a generating sheaf for \mathcal{X} if and only if for every geometric point $\xi : \text{Spec}(K) \rightarrow \mathcal{X}$ and every irreducible representation V of $\text{Aut}(\xi)$, the representation V occurs in \mathcal{F}_{ξ} . The third and fourth conditions are therefore equivalent.

We now show that the third condition is equivalent to the first.

Assume the third condition, and fix a geometric point $\xi : \text{Spec}(K) \rightarrow \mathcal{X}$. Then every irreducible representation of μ_r occurs in the fiber $\bigoplus_{i=0}^{N-1} \mathcal{L}_{\xi}^i$. This implies that the character \mathcal{L}_{ξ} is a generator of $\mathbb{Z}/r\mathbb{Z}$, and it clearly acts freely on the principal bundle of \mathcal{L} . Conversely, if $\mathcal{X} \xrightarrow{\mathcal{L}} \mathcal{B}\mathbb{G}_m$ is representable, then $\mu_r = \text{Aut}(\xi)$

injects into the automorphism group \mathbb{G}_m of \mathcal{L}_ξ , therefore the character of μ_r in \mathcal{L}_ξ is a generator of the dual group $\mathbb{Z}/r\mathbb{Z}$, and therefore all characters occur in $\bigoplus_{i=0}^{r-1} \mathcal{L}_\xi^{-i} \subset \bigoplus_{i=0}^{N-1} \mathcal{L}_\xi^{-i}$. \square

2.4. Embedding of a cyclotomic stack in a weighted projective bundle.

Definition 2.4.1. Let $f : \mathcal{X} \rightarrow B$ be a proper cyclotomic stack, with moduli space $\bar{f} : X \rightarrow B$. A *polarizing line bundle* \mathcal{L} on \mathcal{X} is a uniformizing line bundle, such that there is an \bar{f} -ample invertible sheaf M on X , an integer N and an isomorphism

$$\mathcal{L}^N \simeq \pi^* M.$$

We have the following analogue of the standard properties of ample bundles:

Proposition 2.4.2. *let $\mathcal{X} \rightarrow B$ be a proper family of stacks over a base of finite type, and let \mathcal{L} be a line bundle. The following conditions are equivalent:*

- (1) *The stack \mathcal{X} is cyclotomic, and \mathcal{L} is a polarizing line bundle.*
- (2) *For every coherent sheaf \mathcal{F} on \mathcal{X} there exist integers n_0 and N such that for any $n \geq n_0$ the sheaf homomorphism*

$$\bigoplus_{j=n}^{n+N-1} H^0(\mathcal{X}, \mathcal{L}^j \otimes \mathcal{F}) \otimes \mathcal{L}^{-j} \rightarrow \mathcal{F}$$

is surjective.

- (3) *The stack \mathcal{X} is cyclotomic, uniformized by \mathcal{L} , and for every coherent sheaf \mathcal{F} on \mathcal{X} there exists an integer n_0 such that for any $n \geq n_0$ and any $i \geq 1$ we have*

$$H^i(\mathcal{X}, \mathcal{L}^n \otimes \mathcal{F}) = 0.$$

Proof. Assume the first condition and let \mathcal{F} be a coherent sheaf. Let N be the index of \mathcal{X} . Fix k such that $\mathcal{L}^k = \pi^* M$ with M ample and consider the sheaves $\mathcal{G}_i = \pi_*(\mathcal{L}^i \otimes \mathcal{F})$, $i = 0, \dots, N-1$ on M . There exists m_0 such that for all $m \geq m_0$ the sheaf homomorphisms $H^0(\mathcal{X}, M^m \otimes \mathcal{G}_i) \otimes \mathcal{O}_X \rightarrow \mathcal{G}_i$ are surjective. Note that $M^m \otimes \mathcal{G}_i \simeq \pi_*(\mathcal{L}^{mk+i} \otimes \mathcal{F})$. Also by Proposition 2.3.10 we have that $\pi^* \pi_*(\bigoplus_{i=0}^{N-1} \mathcal{L}^i \otimes \mathcal{F}) \otimes (\bigoplus_{i=0}^{N-1} \mathcal{L}^{-i}) \rightarrow \mathcal{F}$ is surjective. The second condition follows. The reverse implication also follows: \mathcal{L} is uniformizing since the sheaf $\bigoplus_{i=-(N-1)}^0 \mathcal{L}^i$ is necessarily generating by Proposition 2.3.10, and M ample by Serre's criterion.

Since \mathcal{X} is tame we have that π_* is exact on coherent sheaves. Writing $n = mk + j$ we have $H^i(\mathcal{X}, \mathcal{L}^n \otimes \mathcal{F}) = H^i(X, \pi_*(\mathcal{L}^n \otimes \mathcal{F})) = H^i(X, M^m \otimes \pi_*(\mathcal{L}^i \otimes \mathcal{F}))$. The equivalence of the first two conditions with the third follows from Serre's criterion for ampleness applied to M . \square

Proposition 2.4.3. *Let $f : \mathcal{X} \rightarrow B$ be as above, and \mathcal{L} a polarizing line bundle. There are positive integers $n < m$ such that*

(1) for every geometric point $s \in B$ with residue field K , we have

$$H^i(\mathcal{X}_s, \mathcal{L}^j) = 0 \quad \forall i > 0, j \geq n;$$

(2) the sheaf $f_*\mathcal{L}^j$ is locally free $\forall j \geq n$;

(3) for every geometric point $s \in B$, the space

$$\bigoplus_{j=n}^m H^0(\mathcal{X}_s, \mathcal{L}^j)$$

generates the algebra

$$K \oplus \bigoplus_{j \geq n} H^0(\mathcal{X}_s, \mathcal{L}^j).$$

Proof. Part (1) is a special case of part 3 of Proposition 2.4.2.

Part (2) follows from (1) by Grothendieck's "cohomology and base change" applied directly to the f -flat sheaves \mathcal{L}^j since $H^1(\mathcal{X}_s, \mathcal{L}^j) = 0$. Alternatively, consider the sheaves $\pi_*\mathcal{L}^j$. These sheaves are flat over B since \mathcal{X} is tame: since \mathcal{L}^j is flat we have that for any \mathcal{O}_B -ideal I the induced homomorphism $I \otimes_{\mathcal{O}_B} \mathcal{L}^j \rightarrow \mathcal{L}^j$ is injective. Since \mathcal{X} is tame we have π_* exact, therefore $I \otimes_{\mathcal{O}_B} \pi_*\mathcal{L}^j \rightarrow \pi_*\mathcal{L}^j$ is injective, as needed. Since $H^1(X_s, \pi_*\mathcal{L}^j) = 0$ we can apply "cohomology and base change" to $X \rightarrow B$.

For (3), we first note that a range $n \leq j \leq m$ as required exists for any point s of B : for each $i = 0, \dots, N-1$ there exists an l such that

$$H^0(\mathcal{X}, \pi^*M^l \otimes \mathcal{L}^i) \otimes H^0(\mathcal{X}, \pi^*M^r) \rightarrow H^0(\mathcal{X}, \pi^*M^{l+r} \otimes \mathcal{L}^i)$$

is surjective for all r . We may choose a single l that works for all $i = 0, \dots, N-1$ and take $n = lN$.

If at the same time we choose n so that part (1) holds, then elements of $H^0(\mathcal{X}_s, \mathcal{L}^j)$ in the given finite range lift to a neighborhood of s . Also the algebra $H^0(X, \bigoplus_{l \geq 0} M^l)$ is finitely generated and for each j , $H^0(X, (\pi_*\mathcal{L}^j) \oplus M^l)$ is a finite module over the algebra. So a given range for X_s works for a neighborhood of s . Since B is Noetherian, finitely many such neighborhoods cover B and the maximal choice of m works over all B . \square

Corollary 2.4.4. *Let n, m be as in Proposition 2.4.3. Consider the locally free sheaves $W_j = f_*\mathcal{L}^j$. Then we have a closed embedding*

$$\mathcal{X} \hookrightarrow \mathcal{P}\left(\bigoplus_{j=n}^m W_j\right).$$

Proof. Write

$$\mathcal{R}_{n,m} := \mathrm{Sym}_B\left(\bigoplus_{j=n}^m W_j\right),$$

and denote $\mathbb{A} := \mathrm{Spec}_B \mathcal{R}_{n,m}$. We have a \mathbb{G}_m -equivariant map of

$$\mathcal{P}_{\mathcal{L}} \rightarrow \mathbb{A},$$

and since $\mathcal{L}^N = M$ is ample, the image is disjoint from the zero section $0 \subset \mathbb{A}$. In order to show that the morphism of quotient stacks

$$[\mathcal{P}_{\mathcal{L}}/\mathbb{G}_m] = \mathcal{X} \rightarrow [(\mathbb{A} \setminus 0)/\mathbb{G}_m]$$

is an embedding, it suffices to show that the morphism $\mathcal{P}_{\mathcal{L}} \rightarrow \mathbb{A}$ is an embedding.

Let

$$\mathcal{R}_n = \mathcal{O}_B \oplus \bigoplus_{j \geq n} W_j.$$

We have a surjective algebra homomorphism $\mathcal{R}_{n,m} \rightarrow \mathcal{R}_n$, implying that

$$\mathrm{Spec}_B \mathcal{R}_n \rightarrow \mathbb{A}$$

is a closed embedding. It suffices to show that $\mathcal{P}_{\mathcal{L}} \rightarrow \mathrm{Spec}_B \mathcal{R}_n$ is an embedding.

Next, let $\mathcal{R}_{\mathcal{L}} = \mathcal{R}_0 = \bigoplus_{j \geq 0} W_j$. The morphism

$$\mathrm{Spec}_B \mathcal{R}_{\mathcal{L}} \rightarrow \mathrm{Spec}_B \mathcal{R}_n$$

induced by the natural inclusion $\mathcal{R}_n \subset \mathcal{R}_{\mathcal{L}}$ is finite and birational, having its conductor supported along the zero section. It therefore suffices to show that the morphism $\mathcal{P}_{\mathcal{L}} \rightarrow \mathrm{Spec}_B \mathcal{R}_{\mathcal{L}}$ is an embedding.

We now use the finite morphism $\mathcal{P}_{\mathcal{L}} \rightarrow \mathcal{P}_M$, given by the finite ring extension $\bigoplus_{l \in \mathbb{Z}} M^l \hookrightarrow \bigoplus_{j \in \mathbb{Z}} \pi_* \mathcal{L}^j$.

Since M is ample, we have an open embedding $\mathcal{P}_M \hookrightarrow \mathrm{Spec}_B \mathcal{R}_M$. Taking the normalization $\hat{\mathcal{P}}_{\mathcal{L}}$ of $\mathrm{Spec}_B \mathcal{R}_M$ in the structure sheaf of $\mathcal{P}_{\mathcal{L}}$ we get an embedding $\mathcal{P}_{\mathcal{L}} \rightarrow \hat{\mathcal{P}}_{\mathcal{L}}$. Note that $\hat{\mathcal{P}}_{\mathcal{L}}$ is affine over B . Now sections over affines in B of the structure sheaf of $\hat{\mathcal{P}}_{\mathcal{L}}$ restrict to sections of $\pi_* \mathcal{L}^j$, in other words they come from global sections of $\pi_* \mathcal{L}^j$. It follows that $\hat{\mathcal{P}}_{\mathcal{L}} = \mathrm{Spec}_B \bigoplus_{j \in \mathbb{Z}} f_* \mathcal{L}^j$, and since these vanish for $j < 0$ we get $\hat{\mathcal{P}}_{\mathcal{L}} = \mathrm{Spec}_B \mathcal{R}_{\mathcal{L}}$. \square

3. Moduli of stacks with polarizing line bundles

3.1. The category $\mathbf{Sta}^{\mathcal{L}}$. We are now poised to define a moduli stack. In order to avoid cumbersome terminology and convoluted statements, we will define the objects of the stack over schemes of finite type. The extension to arbitrary schemes is standard but less than illuminating.

Definition 3.1.1. We define a category $\mathbf{Sta}^{\mathcal{L}}$, fibered in groupoids over the category of schemes, as follows:

- (1) An object $\mathbf{Sta}^{\mathcal{L}}(B)$ over scheme of finite type consists of a proper family $\mathcal{X} \rightarrow B$ of cyclotomic stacks, with polarizing line bundle \mathcal{L} .

- (2) An arrow from $(\mathcal{X} \rightarrow B, \mathcal{L})$ to $(\mathcal{X}_1 \rightarrow B_1, \mathcal{L}_1)$ consists of a fiber diagram

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\phi} & \mathcal{X}_1 \\ \downarrow & & \downarrow \\ B & \longrightarrow & B_1 \end{array}$$

along with an isomorphism $\alpha : \mathcal{L} \rightarrow \phi^* \mathcal{L}_1$.

Remark 3.1.2. As defined, this is really a 2-category, because arrows of stacks in general have automorphisms, but because \mathcal{L} is polarizing it is easy to see (see, e.g. [AGV08], §3.3.2) that it is equivalent to the associated category, whose morphisms are isomorphism classes of 1-morphisms in the 2-category. We can realize this category directly using the principal bundle $\mathcal{P}_{\mathcal{L}}$ as follows: an object of $\mathit{Sta}^{\mathcal{L}}(B)$ consists of a \mathbb{G}_m -scheme $\mathcal{P} \rightarrow B$, such that $\mathcal{X} := [\mathcal{P}/\mathbb{G}_m]$ is a proper cyclotomic stack polarized by the line bundle associated to \mathcal{P} ; an arrow is a \mathbb{G}_m -equivariant fiber diagram

$$\begin{array}{ccc} \mathcal{P} & \longrightarrow & \mathcal{P}_1 \\ \downarrow & & \downarrow \\ B & \longrightarrow & B_1. \end{array}$$

The fact that $\mathit{Sta}^{\mathcal{L}}$ parametrizes schemes with extra structure can be used to show that it is a stack. We will however show that it is an algebraic stack via a different route - as suggested by the theorem below.

It is also worth noting that $\mathit{Sta}^{\mathcal{L}}$ is highly non-separated and far from finite type.

Remark 3.1.3. Note that the group \mathbb{G}_m acts by automorphisms on every object.

Theorem 3.1.4. *The category $\mathit{Sta}^{\mathcal{L}}$ is an algebraic stack, locally of finite type over \mathbb{Z} . In fact, $\mathit{Sta}^{\mathcal{L}}$ has an open covering by $\sqcup \mathcal{Q}_{\alpha} \rightarrow \mathit{Sta}^{\mathcal{L}}$, where*

- (1) each $\mathcal{Q}_{\alpha} = [H_{\alpha}/G_{\alpha}]$ is a global quotient,
- (2) $H_{\alpha} \subset \mathcal{H}ilb_{\mathcal{P}(\rho_{\alpha})}$ is a quasi-projective open subscheme (described explicitly below), and
- (3) $G_{\alpha} = \text{Aut}_{\rho_{\alpha}}(V)$, where V is the representation space of ρ_{α} .

This theorem is a direct generalization of its well-known analogue, due to Grothendieck, in the context of moduli of projective schemes endowed with an ample sheaf. The proof encompasses the rest of this section.

3.2. Normally embeddable stacks. Before we start the proof in earnest, we give some preliminary results.

Definition 3.2.1. A stack \mathcal{X} uniformized by \mathcal{L} is *normally embeddable* in $\mathcal{P}(\rho)$ if there exists a normal embedding $\mathcal{X} \subset \mathcal{P}(\rho)$ such that $\mathcal{O}_{\mathcal{X}}(1) \simeq \mathcal{L}$.

Proposition 3.2.2. *Fix a function $\mathfrak{F} : \mathbb{Z} \rightarrow \mathbb{Z}$ and two positive integers $n < m$. For $n \leq i \leq m$ fix free modules V_i of rank $f(i)$ and consider the natural representation ρ of \mathbb{G}_m on $V = \bigoplus_{i=n}^m V_i$.*

- (1) *There is an open subscheme $H \subset \text{Hilb}_{\mathcal{P}(\rho), \mathfrak{F}}$ parametrizing normally embedded substacks.*
- (2) *The subcategory $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho) \subset \text{Sta}^{\mathcal{L}}$ of objects with fibers normally embeddable in $\mathcal{P}(\rho)$ and Hilbert–Euler characteristic $\chi(\mathcal{X}, \mathcal{L}^m) = \mathfrak{F}(m)$ satisfies*

$$\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho) \simeq [H/G],$$

with $G = \text{Aut}_{\rho}(V)$.

Remark 3.2.3. Note that G is the group of automorphisms of the free module representation rather than automorphisms of its projectivization. This accounts for the fact that \mathbb{G}_m acts on all objects of the quotient $[H/G]$.

Proof of Proposition. For (1), note that each of the conditions in definition 2.2.1 is an open condition, by the theorem on cohomology and base change.

We prove (2). The data of a map to the quotient stack $B \rightarrow [H/G]$ is equivalent to a family $\mathcal{Y} \rightarrow Q \rightarrow B$ where Q is a principal G -bundle over B and $\mathcal{Y} \subset \mathcal{P}(\rho) \times Q$ is a family of normally embedded substacks. Recall (see Definition 2.3.8) the principal \mathbb{G}_m -bundle $\mathcal{P}_{\mathcal{O}_{\mathcal{Y}}(1)} \rightarrow \mathcal{Y}$ associated to $\mathcal{O}_{\mathcal{Y}}(1)$. The free action of G lifts canonically to $\mathcal{O}_{\mathcal{Y}}(1)$ and to its principal bundle, and commutes with the \mathbb{G}_m action on the line bundle by scaling. To show that $\mathcal{Y} \rightarrow Q$ descends to B , consider the quotient $\mathcal{P}_{\mathcal{O}_{\mathcal{Y}}(1)}/G$. This retains a \mathbb{G}_m action, with finite stabilizers, whose quotient is denoted \mathcal{X} . Let \mathcal{L} denote the line bundle associated to the principal \mathbb{G}_m -bundle $\mathcal{P}_{\mathcal{O}_{\mathcal{Y}}(1)}/G \rightarrow \mathcal{X}$. While we have scrupulously avoided the subtle procedure of taking the quotient of a stack under a free group action, we can identify $\mathcal{X} = \mathcal{Y}/G$ and $\mathcal{L} = \mathcal{O}_{\mathcal{Y}}(1)/G$, namely \mathcal{X} parametrizes principal G bundles carrying an object of \mathcal{Y} and similarly for \mathcal{L} . A diagram summarizing the objects in this construction, where the parallelograms are cartesian squares, is as follows:

$$\begin{array}{ccccccc}
 \mathcal{P}_{\mathcal{O}_{\mathcal{Y}}(1)} & \hookrightarrow & \mathcal{O}_{\mathcal{Y}}(1) & \hookrightarrow & \mathcal{O}(1) \times Q & \longrightarrow & \mathcal{O}(1) \times \mathcal{H} \\
 \downarrow & \searrow & \downarrow & & \downarrow & & \downarrow \\
 \mathcal{P}_{\mathcal{L}} & & \mathcal{Y} & \hookrightarrow & \mathcal{P}(\rho) \times Q & \longrightarrow & \mathcal{P}(\rho) \times \mathcal{H} \\
 & \searrow & \downarrow & \searrow & \downarrow & & \downarrow \\
 & & \mathcal{X} & & Q & \longrightarrow & \mathcal{H} \\
 & & & \searrow & \downarrow & & \downarrow \\
 & & & & B & \longrightarrow & [H/G].
 \end{array}$$

Now $\mathcal{X} \rightarrow B$ is a cyclotomic stack polarized by \mathcal{L} , with geometric fibers embeddable in $\mathcal{P}(\rho)$: indeed the geometric fibers of $\mathcal{X} \rightarrow B$ are isomorphic to those of $\mathcal{Y} \rightarrow Q$,

which are cyclotomic, polarized by $\mathcal{O}_{\mathcal{Y}}(1)$, and embedded in $\mathcal{P}(\rho)$. The whole construction works over arrows $B' \rightarrow B \rightarrow [H/G]$, and therefore gives a functor $[H/G] \rightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho)$.

It is crucial in this construction that we work with the quotient of H by the linear group G and not its projective version - otherwise we would not have a functorial construction of the line bundle \mathcal{L} .

We now consider the opposite direction. Let $\mathcal{X} \rightarrow B$ be a cyclotomic stack polarized by \mathcal{L} . Consider the locally-free sheaf $\oplus_n^m W_j$, with $W_j = f_* \mathcal{L}^j$. Each fiber of W is isomorphic as \mathbb{G}_m -space to a fixed free module V with representation ρ . By the embeddability assumption and Corollary 2.4.4, we have an embedding $\mathcal{X} \hookrightarrow \mathcal{P}(W)$.

Consider the principal G -bundle $\mathcal{Q} = \text{Isom}_{\mathbb{G}_m}(W, V_B)$. We have a canonical isomorphism $W \times_B \mathcal{Q} \simeq V_B$, giving an isomorphism $\mathcal{P}(W) \times_B \mathcal{Q} \simeq \mathcal{P}(\rho) \times \mathcal{Q}$. Write $\mathcal{Y} = \mathcal{X} \times_B \mathcal{Q}$. We have a canonical induced embedding $\mathcal{Y} \hookrightarrow \mathcal{P}(\rho) \times \mathcal{Q}$. This is a normal embedding by definition, and it is clearly G -equivariant. We thus obtain an object of $[H/G]$ as required.

Again the fact that the construction is canonical gives a construction for arrows in $B' \rightarrow B \rightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho)$. The proof that the two functors are inverse to each other is standard and left to the reader. \square

We now consider what happens when we change the range of integers $n \leq i \leq m$.

Proposition 3.2.4. *Fix again a function $\mathfrak{F} : \mathbb{Z} \rightarrow \mathbb{Z}$ and free modules V_i of rank $\mathfrak{F}(i)$. For positive integers $n < m$ we again consider the natural representation $\rho_{n,m}$ of \mathbb{G}_m on $V_{n,m} = \oplus_{i=n}^m V_i$. Then*

- (1) $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \rightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+1})$ is an open embedding.
- (2) For any $n < m$ there exists a canonical open embedding

$$\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \hookrightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n+1,m+n}).$$

- (3) For any $n_1 < m_1$ and $n_2 < m_2$, there exist an integer $m \geq \max(m_1, m_2)$ and canonical open embeddings

$$\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n_1,m_1}) \hookrightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m})$$

$$\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n_2,m_2}) \hookrightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m})$$

with $n = \max(n_1, n_2)$.

Proof of proposition. For (1), it is clear that we have an inclusion $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \hookrightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+1})$. Given a family of graded rings generated in degrees $n, \dots, m+1$, it is an open condition on the base that they are generated in degrees n, \dots, m , as required.

We prove (2). We have an open embedding $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \hookrightarrow \text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+n})$ by (1). In $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+n})$ we have an open substack $\text{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{(n),n+1,m+n})$ of objects

whose projection to the factors in degrees $n + 1, \dots, n + m$ is still an embedding. This also has an open embedding $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{(n),n+1,m+n}) \hookrightarrow \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n+1,m+n})$ as it corresponds to objects such that $H^i(\mathcal{X}, \mathcal{L}^n) = 0$ for $i > 0$.

$$\begin{array}{ccc} \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) & \hookrightarrow & \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+n}) \\ & \searrow & \uparrow \\ & & \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{(n),n+1,m+n}) \hookrightarrow \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n+1,m+n}) \end{array}$$

Now consider a graded ring $R = \bigoplus R_i$. Suppose R is generated in degrees n, \dots, m over R_0 . Then the truncation

$$R' = R_0 \oplus \bigoplus_{i=n+1}^{\infty} R_i$$

is generated in degrees $n + 1, \dots, n + m$. Indeed if $g_1 \cdots g_k$ is a product of homogeneous terms in degrees between n and m , of total degree $> n + m$, and is of minimal degree requiring a term g_1 of degree n as a product of terms in the range $n, \dots, m + n$, then the other terms g_2, \dots, g_k are of degree between $n + 1$ and m , so we may replace g_1 by $g'_1 = g_1 g_k$ and write $g = g'_1 g_2 \cdots g_{k-1}$ as a product of homogeneous terms in the range $n + 1, \dots, n + m$.

This implies that the truncation of the ring of an object in $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+n})$ coming from the open substack $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m})$ is still normally generated. This means that the embedding $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \subset \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m+n})$ factors through the stack $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{(n),n+1,m+n})$. This induces the required open embedding $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \hookrightarrow \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n+1,m+n})$, as required.

Part (3) follows by iterating the construction of part (2). \square

3.3. Proof of Theorem 3.1.4. Given an object $f : \mathcal{X} \rightarrow B$ in $\mathcal{Sta}^{\mathcal{L}}(B)$, there is a maximal open and closed subset $B_0 \subset B$ so that $f_0 : \mathcal{X} \times_B B_0 \rightarrow B_0$ is an object in $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(B_0)$; the formation of B_0 commutes with arbitrary base change. Thus we can decompose

$$\mathcal{Sta}^{\mathcal{L}} = \bigsqcup_{\mathfrak{F}} \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}},$$

where $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}$ is the open and closed fibered subcategory parametrizing cyclotomic stacks polarized by a line bundle with Hilbert-Euler characteristic $\mathfrak{F} : \mathbb{Z} \rightarrow \mathbb{Z}$. We can thus fix \mathfrak{F} and focus on $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}$.

Proposition 3.2.4 gives for each pair of integers $n < m$ a fibered subcategory $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \subset \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}$. This fibered subcategory is open: first note that the condition that the higher cohomologies of \mathcal{L}^i vanish for $n \leq i \leq m$ is open by cohomology and base change. Second, the condition that the ring be normally generated in these degrees (including vanishing of cohomology in higher degrees) is open as well. These open embeddings are compatible with the embeddings in Proposition 3.2.4. It follows that $\cup \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m}) \subset \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}$ is an open subcategory. But by Corollary 2.4.4 every geometric point of $\mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}$ is in $\cup \mathcal{Sta}_{\mathfrak{F}}^{\mathcal{L}}(\rho_{n,m})$, which implies that

$\cup Sta_{\mathfrak{S}}^{\mathcal{L}}(\rho_{n,m}) = Sta_{\mathfrak{S}}^{\mathcal{L}}$ as fibered categories. Since $\cup Sta_{\mathfrak{S}}^{\mathcal{L}}(\rho_{n,m})$ is a direct limit of open subcategories which are algebraic stacks, it is also an algebraic stack. \square

4. Moduli of polarized orbispaces

4.1. Orbispaces.

Definition 4.1.1. An *orbispace* \mathcal{X} is a separated stack with finite diagonal, of finite type over a field, equidimensional, geometrically connected and reduced, and admitting a dense open $U \subset \mathcal{X}$ where U is an algebraic space.

The assumption that \mathcal{X} be connected is made mainly for convenience.

Given two flat families of orbispaces $\mathcal{X}_i \rightarrow B_i$ there exists a natural notion of a 1-morphism of families between them, namely a *cartesian square*

$$\begin{array}{ccc} \mathcal{X}_1 & \rightarrow & \mathcal{X}_2 \\ \downarrow & & \downarrow \\ B_1 & \rightarrow & B_2. \end{array}$$

There is a natural notion of 2-morphism, making families of orbispaces into a 2-category. However by [AV00], Lemma 4.2.3, such 2-morphisms are unique when they exist. It follows that this 2-category is equivalent to the associated category, where morphisms consist of isomorphism classes of 1-morphisms. We call this the *category of families of orbispaces*.

As an example, we have that the weighted projective stack $\mathcal{P}(\rho_1, \dots, \rho_r)$ is an orbispace if and only if $\gcd(\rho_1, \dots, \rho_r) = 1$.

We can consider cyclotomic orbispaces with uniformizing line bundle. Adding the conditions in $Sta^{\mathcal{L}}$ we can define another category - the subcategory $Orb^{\mathcal{L}} \subset Sta^{\mathcal{L}}$ of proper cyclotomic orbispaces with polarizing line bundles. We have

Proposition 4.1.2. *The subcategory $Orb^{\mathcal{L}} \subset Sta^{\mathcal{L}}$ forms an open substack*

Proof. Let $\mathcal{X} \rightarrow B$ be a family of cyclotomic stacks polarized by \mathcal{L} . The locus where the fibers are reduced and connected is open in B : see [Gro66, 12.2.1], for “no embedded points” and then cohomology and base change for “connected”. Since the inertia stack is finite over \mathcal{X} , the same holds for the locus where the fibers have a dense open with trivial inertia. \square

4.2. Polarizations. There is an important distinction to be made between a polarizing line bundle and a polarization - the difference between the data of a line bundle and an element of the Picard group.

Let $(\mathcal{X}_i \rightarrow B_i, \mathcal{L}_i)$ be two families with polarizing line bundles \mathcal{L}_i ; then a morphism comes from a 1-morphism $f_{\mathcal{X}} : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ sitting in a cartesian diagram as above, together with an isomorphism $\alpha : \mathcal{L}_1 \rightarrow f_{\mathcal{X}}^* \mathcal{L}_2$. But a morphism of polarizations should ignore the \mathbb{G}_m -action on the line bundles. The issue is treated extensively in the literature. A procedure for removing the redundant action, called

Proposition 4.3.1. *Consider a proper family of stacks $f : \mathcal{X} \rightarrow B$ and $\mathcal{L}, \mathcal{L}'$ line bundles on \mathcal{X} . Assume the fibers of $\mathcal{X} \rightarrow B$ are reduced and connected.*

There exists a locally closed subscheme $B^0 \subset B$ and a line bundle M on B^0 , over which there exists an isomorphism $\mathcal{L}|_{B^0} \rightarrow \mathcal{L}'|_{B^0} \otimes f_{B^0}^ M$, and is universal with respect to this property.*

By ‘universal’ we mean that if $B^1 \rightarrow B$ is a morphism and M_1 a line bundle on B^1 such that there exists an isomorphism $\mathcal{L}|_{B^1} \rightarrow \mathcal{L}'|_{B^1} \otimes f_{B^1}^* M_1$, then $B^1 \rightarrow B$ factors uniquely through B^0 .

The following is the outcome for polarizations.

Corollary 4.3.2. *Consider a proper family of orbispaces $\mathcal{X} \rightarrow B$ and λ, λ' polarizations on \mathcal{X} . There exists a locally closed subscheme $B^0 \subset B$, over which there exists an isomorphism $\lambda_{B^0} \rightarrow \lambda'_{B^0}$, and is universal with respect to this property.*

Proof. Fix an étale covering $C \rightarrow B$ and line bundles $\mathcal{L}, \mathcal{L}'$ on \mathcal{X}_C representing λ_C, λ'_C . By the proposition there is $C^0 \subset C$ and a line bundle M on C^0 , over which there exists an isomorphism $\mathcal{L}|_{C^0} \rightarrow \mathcal{L}'|_{C^0} \otimes f_C^* M$, and is universal with respect to this property. Since it is universal, the two inverse images of C^0 in $C \times_B C$ coincide, therefore C^0 descends to $B^0 \subset B$, and the data $\mathcal{L}|_{C^0} \rightarrow \mathcal{L}'|_{C^0} \otimes f_C^* M$ gives an isomorphism $\lambda_{B^0} \rightarrow \lambda'_{B^0}$ by definition. \square

Corollary 4.3.3. *Consider a proper family of orbispaces $\mathcal{X} \rightarrow B$, \mathcal{L} a line bundle, λ' a polarization on \mathcal{X} .*

There exists a locally closed subscheme $B^0 \subset B$, over which there exists an isomorphism $\lambda_{B^0} \rightarrow \lambda'_{B^0}$, with λ the polarization induced by \mathcal{L} , and is universal with respect to this property.

Proof. This is immediate from the previous corollary. \square

4.4. Moduli of canonically polarized orbispaces.

Lemma 4.4.1. *Let $\mathcal{X} \rightarrow B$ be a proper family of orbispaces. There exists open subschemes $B_{\text{gor}} \subset B_{\text{cm}} \subset B$ where the fibers of $\mathcal{X} \rightarrow B$ are Gorenstein and Cohen-Macaulay respectively.*

Proof. Recall the following fact [Gro66, 12.2.1]: Suppose $X \rightarrow B$ is a flat morphism of finite type with pure dimensional fibers and F is a coherent \mathcal{O}_X -module flat over B . Then for each $r \geq 0$ the locus

$$\{b \in B : F|_{X_b} \text{ is } S_r\}$$

is open in B . Take $F = \mathcal{O}_X$ and $r = \dim(X/B)$ to see that the locus where the fibers are Cohen-Macaulay is open.

When the fibers of $X \rightarrow B$ are Cohen-Macaulay, the shifted relative dualizing complex $\omega_{X/B}^\bullet[-n]$ is a sheaf $\omega_{X/B}$ that is invertible precisely on the open subset where the fibers are Gorenstein (see [Con00], Theorem 3.5.1, Corollary 3.5.2, and the subsequent discussion).

It remains to check this analysis applies to the stack \mathcal{X} . In characteristic 0, note that the dualizing sheaf is insensitive to étale localization [Con00, Th. 4.4.4 and p. 214], and thus descends canonically to \mathcal{X} . In arbitrary characteristic one has to use smooth covers instead. However, it is a fundamental property of the dualizing complex that it behaves well under smooth morphisms, i.e., if $g : Y \rightarrow Z$ is smooth and $Z \rightarrow B$ is flat then (up to shifts)

$$\omega_{Y/B}^\bullet = g^* \omega_{Z/B}^\bullet \otimes \omega_{Y/Z}.$$

Indeed, these formulas are fundamental tools in showing that the dualizing complex is well-defined (see [Con00, pp. 29-30, Th. 2.8.1 and 3.5.1]). In particular, we can define $\omega_{\mathcal{X}/B}^\bullet$ via smooth covers.

In the particular case at hand with $\mathcal{X} = [\mathcal{P}/\mathbb{G}_m]$, the smooth cover is given by \mathcal{P} and the descent datum is given by the \mathbb{G}_m -equivariant structure on $\omega_{\mathcal{P}}$. (Duality for Artin stacks is developed in [Nir08]. However we will only need the existence of a unique $\omega_{\mathcal{X}}$.) \square

The lemma implies:

Proposition 4.4.2. *There are open substacks $\text{Orb}_{\text{gor}}^\lambda \subset \text{Orb}_{\text{cm}}^\lambda \subset \text{Orb}^\lambda$ parametrizing Gorenstein and Cohen-Macaulay polarized orbispaces.*

For most applications to compactifications of moduli spaces these substacks should be sufficient (see Remark 6.1.6). However, in some situations more general singularities might be needed:

Definition 4.4.3. A family $\mathcal{X} \rightarrow B$ of orbispaces is *canonically polarized* if

- the fibers are pure-dimensional, satisfy Serre's condition S_2 , and are Gorenstein in codimension one;
- $\omega_{\mathcal{X}/B}^\circ$, namely the component of the dualizing complex $\omega_{\mathcal{X}/B}^\bullet$ in degree $\dim B - \dim X$, is locally free and polarizing.

The *canonical polarization* on $\mathcal{X} \rightarrow B$ is the polarization induced by $\omega_{\mathcal{X}/B}^\circ$.

Theorem 4.4.4. *There is a locally closed substack $\text{Orb}^\omega \subset \text{Orb}^\lambda$ parametrizing canonically polarized orbispaces.*

Proof. Lemma 4.4.1 guarantees the conditions on the fibers are open. The change-of-rings spectral sequence gives a homomorphism of sheaves

$$\omega_{\mathcal{X}/B}^\circ|_{\mathcal{X}_b} \rightarrow \omega_{\mathcal{X}_b}^\circ.$$

This is injective because the relative dualizing sheaf is invertible and \mathcal{X}_b is reduced; its cokernel is supported on a subset of \mathcal{X}_b of codimension two. The dualizing sheaf $\omega_{\mathcal{X}_b}^\circ$ is automatically saturated (cf. [Rei87, 1.6]), so any injection from an invertible sheaf that is an isomorphism in codimension one is an isomorphism. Thus on the open locus where $\omega_{\mathcal{X}/B}^\circ$ is invertible, its formation commutes with arbitrary base change. Applying Corollary 4.3.3, we obtain the theorem. \square

We denote the intersection $\mathcal{O}rb_{\text{gor}}^\omega := \mathcal{O}rb^\omega \cap \mathcal{O}rb_{\text{gor}}^\lambda$, the substack of canonically polarized Gorenstein orbispaces.

Note that over $\mathcal{O}rb^\omega$, the universal polarization is represented by an invertible sheaf, namely, the relative dualizing sheaf of the universal family. This gives us a lifting

$$\begin{array}{ccc} & & \mathcal{O}rb^\mathcal{L} \\ & \nearrow \text{dotted} & \downarrow \\ \mathcal{O}rb^\omega & \longrightarrow & \mathcal{O}rb^\lambda. \end{array}$$

Indeed we can describe $\mathcal{O}rb^\omega$ or $\mathcal{O}rb_{\text{gor}}^\omega$ directly in terms of line bundles as follows: an object of $\mathcal{O}rb^\omega$ is a triple $(\mathcal{X}, \mathcal{L}, \phi)$ where \mathcal{X} is an orbispace on which $\omega_{\mathcal{X}/B}^\circ$ is invertible, \mathcal{L} a line bundle, and $\phi : \mathcal{L} \rightarrow \omega_{\mathcal{X}/B}^\circ$ an isomorphism. Arrows are fiber diagram as for $\mathcal{O}rb^\omega$. But note that an arrow does not involve the choice of an arrow on $\omega_{\mathcal{X}/B}^\circ$: such an arrow is canonically given as the unique arrow on $\omega_{\mathcal{X}/B}^\circ$ respecting the trace map $\mathbf{R}^n \pi_* \omega_{\mathcal{X}/B}^\circ \rightarrow \mathcal{O}_B$. The relative dualizing sheaf (and complex) does have \mathbb{G}_m acting as automorphisms, but this automorphism group acts effectively on the trace map.

5. Kollár families and stacks

5.1. Reflexive sheaves, saturation and base change. Let X be a reduced scheme of finite type over a field, having pure dimension d , satisfying Serre's condition S_2 .

Lemma 5.1.1. (1) *Let F be a coherent sheaf on X . Then the sheaf $F^* := \text{Hom}(F, \mathcal{O}_X)$ is S_2 .*

(2) *Suppose $\psi : F \rightarrow G$ is a morphism of S_2 -sheaves which is an isomorphism on the complement of a closed subscheme of codimension ≥ 2 . Then ψ is an isomorphism.*

(3) *Let F be an S_2 coherent sheaf on X . Then the morphism $F \rightarrow F^{**}$ is an isomorphism.*

Proof. The problems being local, we assume X is affine.

(1) Let ϕ be a section of $\text{Hom}(F, \mathcal{O}_X)$ over an open set U with $\text{codim}(X \setminus U) \geq 2$. Let f be a section of F on X and consider $\phi(f_U) \in H^0(U, \mathcal{O}_U)$. Since X is S_2 this extends uniquely to a regular function g on X . We define $\tilde{\phi}(f) = g$.

(2) Let g be a section of \mathcal{G} . Let U be an open subset with codimension ≥ 2 complement on which ψ is an isomorphism. Set $f_U = \psi^{-1}(g_U)$, then f_U is uniquely the restriction of a section f of F , and $f = \psi^{-1}g$.

(3) Since F and F^{**} are S_2 , the homomorphism $F \rightarrow F^{**}$ is an isomorphism. □

Definition 5.1.2. A coherent sheaf F on X is said to be *reflexive* if the morphism $F \rightarrow F^{**}$ is an isomorphism.

Definition 5.1.3. Let F be a coherent sheaf on X , and n a *positive* integer. We define

$$F^{[-n]} = \text{Hom}(F^{\otimes n}, \mathcal{O}_X).$$

and

$$F^{[n]} = \text{Hom}(F^{[-n]}, \mathcal{O}_X).$$

Theorem 10 of [Kol95] implies

Proposition 5.1.4. *Let $X \rightarrow B$ be a flat morphism of finite type with S_2 fibers of pure dimension d . Let $U \subset X$ be an open subscheme, dense in each fiber. Let F be a coherent sheaf on X , locally free on U . If the formation of $F^{[n]}$ commutes with arbitrary base extension then $F^{[n]}$ is flat over B .*

5.2. Kollár families of \mathbb{Q} -line bundles.

Definition 5.2.1. (1) By a *Kollár family of \mathbb{Q} -line bundles* we mean

- (a) $\bar{f} : X \rightarrow B$ a flat family of equidimensional connected reduced schemes satisfying Serre's condition S_2 ,
- (b) F a coherent sheaf on X , such that
 - (i) for each fiber X_b , the restriction $F|_{X_b}$ is reflexive of rank 1;
 - (ii) for every n , the formation of $F^{[n]}$ commutes with arbitrary base change;
 - (iii) for each geometric point s of B there is an integer $N_s \neq 0$ such that $F^{[N_s]}|_{X_s}$ is invertible.

- (2) A *morphism* from a Kollár family $(X \rightarrow B, F)$ to another $(X_1 \rightarrow B_1, F_1)$ consists of a cartesian diagram

$$\begin{array}{ccc} X & \xrightarrow{\bar{\phi}} & X_1 \\ \downarrow & & \downarrow \\ B & \longrightarrow & B_1 \end{array}$$

along with an isomorphism $\bar{\alpha} : F \rightarrow \bar{\phi}^* F_1$.

- (3) Parts (1) and (2) above define the objects and arrows of a category of *Kollár families of \mathbb{Q} -line bundles*, fibered over the category Sch_k of k -schemes. It has an important open subcategory $\mathcal{K}^{\mathcal{L}}$ of *Kollár families of polarizing \mathbb{Q} -line bundles*, where $X \rightarrow B$ is *proper* and $F^{[N_s]}|_{X_s}$ is *ample*.

Note that (1.b.i) and (1.b.ii) imply that $F = F^{[1]}$. Furthermore, (1.b.i) and (1.b.iii) imply that F is invertible in codimension 1. A reflexive rank-one sheaf on a reduced one-dimensional scheme X is a fractional ideal J . Indeed, express \mathcal{F}^* locally as a quotient $\mathcal{O}_X^{\oplus r} \rightarrow F \rightarrow 0$, which realizes \mathcal{F} as a subsheaf of $\mathcal{O}_X^{\oplus r}$. Choose a projection $\mathcal{O}_X^{\oplus r} \rightarrow \mathcal{O}_X$ such that the composed homomorphism $\mathcal{F} \rightarrow \mathcal{O}_X$ has rank one at each generic point. This is injective since our scheme is reduced. The saturated powers $F^{[n]}$ equal J^n , i.e., the powers of J as an ideal. In particular, J^N is locally principal. The blow-up of X along J is

$$\mathrm{Proj}_X \oplus_{n \geq 0} J^n \simeq \mathrm{Proj}_X \oplus_{n \geq 0} J^{Nn} = X,$$

so J is locally principal and F is invertible.

One can show that the category of *proper* Kollár families is an algebraic stack. We will show below that the subcategory $\mathcal{K}^{\mathcal{L}}$ is an algebraic stack by identifying it as an algebraic substack of $\mathcal{O}rb^{\mathcal{L}}$. Kollár families were introduced in the canonical case $F = \omega_{X/B}$ in [Kol90, 5.2] and [HK04, 2.11]. In [Hac04] it was shown that they admit, at least in the canonical case, a good deformation-obstruction theory. As we see below, this holds in complete generality. Indeed Hacking's approach is via the associated stacks, as is ours. In [Kol08] Kollár provides an approach without stacks.

We note that the index N_s is bounded in suitable open sets:

Lemma 5.2.2. *Let $(X \rightarrow B, F)$ be a Kollár family of \mathbb{Q} -line bundles, and $s \in B$ a geometric point. Let N_s be an integer such that $F^{[N_s]}|_{X_s}$ is invertible. Then there is an open neighborhood U of s such that $F^{[N_s]}|_{X_U}$ is invertible.*

Proof. Since $F^{[N]}$ are flat and their formation commutes with base change, the assumption that $F^{[N_s]}|_{X_s}$ is invertible implies $F^{[N_s]}$ is invertible in a neighborhood of X_s . Since $X \rightarrow B$ is of finite type the assertion follows. \square

5.3. Kollár families and uniformized twisted varieties.

Definition 5.3.1. Let $(\bar{f}: X \rightarrow B, F)$ be a Kollár family of \mathbb{Q} -line bundles.

- (1) The \mathbb{G}_m -space of F is the X -scheme

$$\mathcal{P}_F = \mathrm{Spec}_X \left(\bigoplus_{j \in \mathbb{Z}} F^{[j]} \right).$$

Note that $\mathcal{P}_F \rightarrow B$ is flat by Proposition 5.1.4.

- (2) The *associated stack* is

$$\mathcal{X}_F = [\mathcal{P}_F / \mathbb{G}_m].$$

This comes with the *natural line bundle* $\mathcal{L} = \mathcal{L}_F$ associated to the principal bundle

$$\mathcal{P}_F \rightarrow \mathcal{X}_F.$$

We drop the index F and write \mathcal{X} and \mathcal{P} when no confusion is likely. We denote by $\pi : \mathcal{X} \rightarrow X$ the resulting morphism.

Proposition 5.3.2. (1) *The family $\mathcal{X} \rightarrow B$ is a family of cyclotomic orbispaces uniformized by \mathcal{L} , with fibers satisfying Serre's condition S_2 .*

(2) *The morphism $\pi : \mathcal{X} \rightarrow X$ makes X into the coarse moduli space of \mathcal{X} . This morphism is an isomorphism on the open subset where F is invertible, the complement of which has codimension > 1 in each fiber.*

(3) *For any integer a , we have $\pi_*(\mathcal{L}^a) = F^{[a]}$ (in particular $\pi_*(\mathcal{L}^a)$ is reflexive).*

(4) *This construction is functorial, that is, given a morphism of Kollár families $(\bar{\phi}, \bar{\alpha})$ from $(X \rightarrow B, F)$ to $(X_1 \rightarrow B_1, F_1)$ we have canonically $\mathcal{P}_F \simeq \bar{\phi}^* \mathcal{P}_{F_1}$ and $\mathcal{X}_F \simeq \bar{\phi}^* \mathcal{X}_{F_1}$.*

Proof. To verify that $\mathcal{X} \rightarrow B$ is cyclotomic and uniformized by \mathcal{L} , we just need to check that \mathbb{G}_m acts on \mathcal{P}_F with finite stabilizers. Lemma 5.2.2 allows us to assume that $F^{[N]}$ is locally free for some $N > 0$. We have natural homomorphisms

$$m_a : (F^{[N]})^a \longrightarrow F^{[Na]}$$

for each $a \in \mathbb{Z}$. For each $b \in B$, the sheaf F is locally free on an open set $U_b \hookrightarrow X_b$ with codimension-two complement, so m_a is an isomorphism over U_b , and hence over all X_b . It follows that $(F^{[N]})^a = F^{[Na]}$.

Consider the \mathbb{G}_m -equivariant morphism

$$\mathcal{P}_F \longrightarrow \mathrm{Spec}_X \left(\bigoplus_{a \in \mathbb{Z}} F^{[aN]} \right).$$

Note that \mathbb{G}_m acts on the target with stabilizer μ_N , which implies the stabilizers on \mathcal{P}_F are subgroups of μ_N .

To complete part (1), we note that a fiber in \mathcal{P}_F over $b \in B$ is S_2 as the spectrum of an algebra with reflexive components over an S_2 base. Also $\mathcal{P}_F \rightarrow \mathcal{X}$ is smooth and surjective. Smoothness implies that the quotient stack \mathcal{X} is S_2 as well.

Since the coarse moduli space is obtained by taking invariants (see the proof of Lemma 2.3.6), and since the invariant part of $\bigoplus F^{[a]}$ is $F^{[0]} = \mathcal{O}_X$, we have that X is the coarse moduli space of \mathcal{X} . Over the locus U where F is invertible, the scheme \mathcal{P}_F is a principal \mathbb{G}_m -bundle, so

$$\mathcal{X}_U = [(\mathcal{P}_F)|_U / \mathbb{G}_m] = U.$$

Part (3) follows since \mathcal{L}^a is the degree- a component of the graded $\mathcal{O}_{\mathcal{X}}$ -algebra $\bigoplus \mathcal{L}^i$, whose spectrum is \mathcal{P}_F . The degree- a component of the graded \mathcal{O}_X algebra $\bigoplus_{i \in \mathbb{Z}} F^{[i]}$ is $F^{[a]}$.

Part (4) follows from the functoriality of Spec and of the formation of quotient stack. □

Remark 5.3.3. If we assume for each $a \in \mathbb{Z}$ and $b \in B$ that $F^{[a]}$ is S_r (in particular the fibers of X are S_r), it follows that the fibers of \mathcal{X} are S_r as well.

Definition 5.3.4. A family of *uniformized twisted varieties* $(\mathcal{X} \rightarrow B, \mathcal{L})$ is a flat family of S_2 cyclotomic orbispaces uniformized by \mathcal{L} , such that the morphism $\pi : \mathcal{X} \rightarrow X$ to the coarse moduli space is an isomorphism away from a subset of codimension > 1 in each fiber.

We define 1-morphisms of families of uniformized twisted varieties as fibered diagrams. Of course, there is a notion of 2-morphism making these into a 2-category, but since these are orbispaces, a 2-isomorphism is unique when it exists [AV02, Lemma 4.2.3], so this 2-category is equivalent to the associated category, whose morphisms are isomorphism classes of 1-morphisms.

Remark 5.3.5. We insist on $\pi : \mathcal{X} \rightarrow X$ an isomorphism in codimension-1 to obtain an equivalence with Kollár families as in Theorem 5.3.6 below. To describe uniformized orbispaces in terms of sheaves on X in general, it is necessary to specify an algebra of sheaves, which can be quite delicate especially when $\mathcal{X} \rightarrow X$ is branched along singular codimension-1 loci. For a subtle example see [Jar00, Definition 2.5] as well as [AJ03, §4.2-4.4], where an equivalence analogous to our Theorem 5.3.6 below is given.

Theorem 5.3.6. *The category of Kollár families of \mathbb{Q} -line bundles is equivalent to the category of uniformized twisted varieties via the base preserving functors*

$$(X \rightarrow B, F) \quad \mapsto \quad (\mathcal{X}_F \rightarrow B, \mathcal{L}_F),$$

with $\mathcal{X}_F = [\mathcal{P}_F / \mathbb{G}_m]$, and its inverse

$$(\mathcal{X} \rightarrow B, \mathcal{L}) \quad \mapsto \quad (X \rightarrow B, \pi_* \mathcal{L}),$$

where X is the coarse moduli space of \mathcal{X} .

Proof. Proposition 5.3.2 gives the functor from the category of Kollár families of \mathbb{Q} -line bundles to the category of uniformized twisted varieties. We now give an inverse. Fix a uniformized twisted variety $(\mathcal{X} \rightarrow B, \mathcal{L})$, with coarse moduli space $\pi : \mathcal{X} \rightarrow X$. Since the formation of π commutes with arbitrary base change (Lemma 2.3.6), the universal property of coarse moduli spaces guarantees that each fiber of $X \rightarrow B$ is reduced. If N is the index of \mathcal{X} then, by Lemma 2.3.7, \mathcal{L}^N descends to an invertible sheaf on X , which coincides with $\pi_* \mathcal{L}^N$.

Each geometric point of \mathcal{X} admits an étale neighborhood which is isomorphic to a quotient of an affine S_2 -scheme V by the action of μ_r . Locally, the coarse moduli space is the scheme-theoretic quotient V/μ_r . However, since μ_r is linearly reductive, any quotient of an S_2 -scheme by μ_r is also S_2 , because the invariants are a direct summand in the coordinate ring of V . Similarly the sheaves $\pi_* \mathcal{L}^j$ are direct summands in the algebra of \mathcal{P} , which is affine over X , flat over B with S_2 fibers. It follows that these sheaves are flat over B , saturated, and their formation commutes with base change on B . \square

Corollary 5.3.7. *The category $\mathcal{K}^{\mathcal{L}}$ is an algebraic stack, isomorphic to the open substack of $\mathcal{O}rb^{\mathcal{L}}$ where $(\mathcal{X} \rightarrow B, \mathcal{L})$ are uniformized twisted varieties.*

Proof. Using Theorem 5.3.6 note that $\mathcal{K}^{\mathcal{L}}$ indeed parametrizes orbispaces with polarizing line bundles which are at the same time uniformized twisted varieties. This is open in $\mathcal{O}rb^{\mathcal{L}}$ since the conditions of being S_2 is open ([Gro66, 12.2.1]) and the condition on the fibers of inertia having support in codimension > 1 is open by semicontinuity of fiber dimensions in proper morphisms. \square

Definition 5.3.8. We define $\mathcal{K}^{\lambda} = \mathcal{K}^{\mathcal{L}} // \mathbb{G}_m$, where \mathbb{G}_m acts by scalars on \mathcal{L} . We define $\mathcal{K}^{\omega} \subset \mathcal{K}^{\lambda}$ as the locally closed subcategory corresponding to canonically polarized twisted varieties $(f : \mathcal{X} \rightarrow B, \lambda)$ where λ is given by $\omega_{\mathcal{X}/B}^{\circ}$.

We define $\mathcal{K}_{\text{gor}}^{\omega} \subset \mathcal{K}^{\omega}$ to be the open substack where the fibers of f are Cohen-Macaulay. Since the dualizing sheaf is invertible—it is the polarizing line bundle—the fibers are automatically Gorenstein.

Note that Corollary 4.3.3 guarantees that the second condition is locally closed.

Remark 5.3.9. We can interpret \mathcal{K}^{ω} in terms of Kollár families $(\bar{f} : X \rightarrow B, F)$ with isomorphism $F \rightarrow \omega_{X/B}^{\circ}$. The substack $\mathcal{K}_{\text{gor}}^{\omega}$ is characterized by

- (1) the fibers of f are Cohen-Macaulay;
- (2) each power $\omega_{X/B}^{[n]}$ is Cohen-Macaulay.

Indeed, if a sheaf \mathcal{G} on \mathcal{X} is Cohen-Macaulay then $\pi_*\mathcal{G}$ is also Cohen-Macaulay: locally write \mathcal{X} as a quotient of an affine scheme V by μ_r , and \mathcal{G} corresponding to an equivariant module G . The complex computing local cohomology of G can be taken μ_r -equivariant, and the invariant subcomplex splits. Therefore, if the local cohomology vanishes, then the invariant part, which computes local cohomology of $G^{\mu_r} = \pi_*\mathcal{G}$ on X , vanishes as well.

In general, the second condition is not a logical consequence of the first. There are examples [Sin03, §6] of Cohen-Macaulay log canonical threefolds whose index-one covers are not Cohen-Macaulay.

Remark 5.3.10. In characteristic zero, varieties with canonical (or even log terminal) singularities admit index-one covers with canonical singularities [Rei80], which are therefore rational [Elk81] and Cohen-Macaulay [KM98]. However, in positive characteristic there exist log terminal surface singularities with index-one covers that are not canonical [Kaw99b]. Generally, index-one covers present technical difficulties when the index of the singularity is divisible by the characteristic, especially in small characteristics. Our approach sidesteps these issues.

6. Semilog canonical singularities and compactifications

From here on we assume our schemes are over a field of characteristic 0.

6.1. Properness results and questions. Recall that canonical singularities deform to canonical singularities [Kaw99a]. This fact and Proposition A.1.1 of the appendix allow us to formulate the following:

Definition 6.1.1. We define $\mathcal{K}_{\text{can}}^\omega \subset \mathcal{K}^\omega$ as the open substack corresponding to canonically polarized twisted varieties where *both* \mathcal{X} and X have canonical singularities.

We define $\mathcal{K}_{\text{slc}}^\omega, \mathcal{K}_{\text{gor-slc}}^\omega \subset \mathcal{K}^\omega$ as the open substack corresponding to canonically polarized twisted varieties where \mathcal{X} has semilog canonical (resp. Gorenstein semilog canonical) singularities.

Objects of $\mathcal{K}_{\text{slc}}^\omega$ are called *families of twisted stable varieties*.

Theorem 5.3.6 induces equivalences of the categories above with the respective categories of Kollár families. Accordingly, we sometimes call $\mathcal{K}_{\text{slc}}^\omega$ the stack of *Kollár families of stable varieties*, with the understanding that the \mathbb{Q} -line bundle is the canonical polarization.

We review what is known about properness of moduli spaces of stable varieties:

Proposition 6.1.2. *Each connected component of the closure of $\mathcal{K}_{\text{can}}^\omega$ in $\mathcal{K}_{\text{slc}}^\omega$ is proper with projective coarse moduli space.*

Proof. Most of this is contained in [Kar00], though our definition of families differs from Karu's.

We first check that the closure of $\mathcal{K}_{\text{can}}^\omega$ satisfies the valuative criterion for properness.

Let Δ be the spectrum of a discrete valuation ring, with special point s and generic point η . Let \mathcal{X}_η be a canonically-polarized orbispace with Gorenstein canonical singularities over η . Let X_η denote its coarse moduli space.

Apply resolution of singularities and semistable reduction [KKMSD73] to obtain

- a finite branched cover $\Delta' \rightarrow \Delta$ with generic point η' and special point s' ;
- a nonsingular variety Y and a flat projective morphism $\phi : Y \rightarrow \Delta'$, with $Y_{s'}$ reduced simple normal crossings;
- a birational morphism $Y_{\eta'} \rightarrow X_{\eta'}$.

Consider the canonical model of Y relative to ϕ

$$\psi : W \rightarrow \Delta',$$

which exists due to [BCHM10, Theorem 1.1]. It has the following properties:

- $K_{W/\Delta'}$ is \mathbb{Q} -Cartier and ample relative to ψ ;
- W has canonical singularities, so every exceptional divisor has discrepancy ≥ 0 .

As a consequence, we deduce $W_{\eta'} \simeq X_{\eta'}$ and W is Cohen-Macaulay.

Furthermore, every exceptional divisor in the special fiber has log discrepancy ≥ 0 with respect to $(W, W_{s'})$, thus W_s has normal crossings in codimension one. In particular, the pair $(W, W_{s'})$ is log canonical. Applying ‘Adjunction’ [Kaw07] and criterion A.1.3, we conclude that $W_{s'}$ has semilog canonical singularities. The variety $W_{s'}$ is the desired stable limit of X_{η} ; it is unique by the uniqueness of the canonical models.

Now we construct our desired stack-theoretic stable limit. The point is that the stack-theoretic canonical model has canonical singularities for the same reason W has. In classical terms, since W has canonical singularities, its index-one cover does as well [KM98, 5.20,5.21]. Furthermore, the direct-image sheaves

$$\psi_*\omega_{\psi}^{[n]}, \quad n \geq 0,$$

are locally free and commute with restriction to the fibers, i.e.,

$$\psi_*\omega_{\psi}^{[n]}|_{s'} = \Gamma(W_t, \omega_{W_{s'}}^{[n]}).$$

This formulation of ‘deformation invariance of plurigenera’ follows from the existence of minimal models [Nak86, Cor. 3, Thm. 8] and the good behavior of direct images of dualizing sheaves [Kol86, Thm 2.1].

Since W is a canonical model, we can express

$$W = \text{Proj}_{\Delta'} \oplus_{n \geq 0} \psi_*\omega_{\psi}^{[n]}.$$

The corresponding affine cone

$$CW := \text{Spec}_{\Delta'} \oplus_{n \geq 0} \psi_*\omega_{\psi}^{[n]}$$

comes with a \mathbb{G}_m -action associated with the grading, and we define

$$\mathcal{W} = [(CW \setminus 0) / \mathbb{G}_m].$$

Again, we have $\mathcal{W}_{\eta'} \simeq \mathcal{X}_{\eta'}$ (as polarized orbispaces) and thus $\mathcal{W}_{s'}$ has Gorenstein semilog canonical singularities. Indeed, \mathcal{W} has canonical singularities (it is locally modeled by the index-one cover of W) and thus is Cohen-Macaulay; the dualizing sheaf $\omega_{\mathcal{W}/\Delta'}$ is invertible by the \mathbb{G}_m -quotient construction. Repeating the previous adjunction analysis yields that $\mathcal{W}_{s'}$ is semilog canonical.

The uniqueness of the stack-theoretic limit follows from the uniqueness of the canonical model, via Theorem 5.3.6.

We turn now to boundedness: [Kar00, §3] still shows that each connected component of the closure of $\mathcal{K}_{\text{can}}^{\omega}$ is of finite type: Matsusaka’s big theorem shows the open locus $\mathcal{K}_{\text{can}}^{\omega}$ is of finite type. We apply weak semistable reduction to an arbitrary compactification of a tautological family, and obtain a family $X \rightarrow B$ which is weakly semistable. In particular, B is smooth. Applying finite generation ([BCHM10]) we can take the relative canonical model $X^{\text{can}} = \text{Proj}_B R(X/B) \rightarrow B$, or its orbispace version $\mathcal{X}^{\text{can}} = \mathbf{Proj}_B R(X/B) \rightarrow B$. The total space has

canonical singularities as it is a canonical model - this is true for either X^{can} or \mathcal{X}^{can} . Deformation invariance of plurigeners shows that this is flat. An argument similar to the one above shows that the fibers are semilog canonical. This gives a compactified family over a base of finite type as required.

The argument there, citing Kollár's projectivity criterion [Kol90], shows that properness of the closure of $\mathcal{K}_{\text{can}}^\omega$ implies projectivity of its coarse moduli space. \square

We would like the following formulation of 'semi-canonical models':

Assumption 6.1.3. Suppose that Δ is the spectrum of a discrete valuation ring with special point s and generic point η . Let X_η be a stable variety over η . Then there exists a finite ramified base-change $\Delta' \rightarrow \Delta$ with special point s' and generic point η' , a unique variety W , and a flat morphism $\psi : W \rightarrow \Delta'$ satisfying the following:

- $W_{\eta'}$ is isomorphic to the base-change $X_{\eta'} = X \times_\eta \eta'$.
- The relative dualizing sheaf ω_ψ is \mathbb{Q} -Cartier and ample.
- Each reflexive power $\omega_\psi^{[n]}$ is S_2 relative to ψ , i.e., S_2 on restriction to the fibers. If X_η is Cohen-Macaulay, we expect the sheaves $\omega_\psi^{[n]}$ to be Cohen-Macaulay as well. [KK09]
- $W_{s'}$ has semilog canonical singularities.
- The sheaves $\psi_* \omega_\psi^{[n]}$, $n \geq 0$ are locally free and satisfy

$$\psi_* \omega_\psi^{[n]}|_{s'} = \Gamma(W_{s'}, \omega_{W_{s'}}^{[n]}).$$

Proposition 6.1.4. *Under Assumption 6.1.3, $\mathcal{K}_{\text{slc}}^\omega$ satisfies the valuative criterion for properness. In particular, each connected component of finite type is proper.*

Remark 6.1.5. To say that each connected component of $\mathcal{K}_{\text{slc}}^\omega$ is of finite type is a boundedness assertion about the stable varieties with given numerical invariants. These assertions have been proven in dimensions ≤ 2 [Ale94], but remain open in higher dimensions.

Remark 6.1.6. The discussion following Corollary 1.3 of [KK09] implies that the stable limit of a one-parameter family of Cohen-Macaulay stable varieties (resp. Gorenstein twisted stable varieties) is Cohen-Macaulay (resp. Gorenstein). Thus the $\mathcal{K}_{\text{gor-slc}}^\omega \subset \mathcal{K}_{\text{slc}}^\omega$ is a union of connected components.

Nevertheless, there are good reasons to consider non-Cohen-Macaulay singularities. We have already mentioned non-Cohen-Macaulay log canonical singularities in Remark 5.3.9. In characteristic zero, log terminal singularities and their index-one covers are automatically Cohen-Macaulay [KM98].

A. The semilog canonical locus is open

A.1. Statement. The following result is known for surfaces ([KSB88, Kol90]). The present proof was suggested to us by V. Alexeev. The second part can be found in [Kar00].

Proposition A.1.1. *Let B be a scheme of finite type over a field of characteristic zero and $f : X \rightarrow B$ be a flat morphism of finite type with connected reduced equidimensional fibers. Assume that the fibers are S_2 and Gorenstein in codimension one.*

Suppose there exists an invertible sheaf \mathcal{L} on X such that for each $b \in B$ we have

$$\mathcal{L}|_{X_b} \simeq j_* \omega_U^{\otimes n}, \quad n \in \mathbb{N},$$

where $j : U \hookrightarrow X_b$ is the open subset where X_b is Gorenstein. Then the locus where the fibers have semilog canonical singularities

$$B^{slc} = \{b \in B : X_b \text{ is semilog canonical}\}$$

is open in B .

Remark A.1.2. In the case where the fibers are Gorenstein, the existence of \mathcal{L} is automatic. Indeed, the relative dualizing sheaf $\omega_{X/B}$ is itself invertible.

Before starting the proof proper, we recall a characterization of semilog canonical singularities, which can be found in chapter 2 of [Kol92]:

Lemma A.1.3. *Let (Y, D) be a pair consisting of a connected equidimensional S_2 scheme Y and a reduced effective divisor D with no components contained in the singular locus of Y . Suppose that Y has normal crossings in codimension one and that the Weil divisor $K_Y + D$ is \mathbb{Q} -Cartier. Let $\nu : \tilde{Y} \rightarrow Y$ denote the normalization and $C \subset \tilde{Y}$ its conductor, i.e., the reduced effective divisor such that*

$$\nu^*(K_Y + D) = K_{\tilde{Y}} + \tilde{D} + C,$$

where \tilde{D} is the proper transform of D . Then the following conditions are equivalent:

- (Y, D) is semilog canonical.
- $(\tilde{Y}, \tilde{D} + C)$ is log canonical.

The condition of having normal crossings in codimension 1 is also open. Indeed, this means that the only singularities in codimension 1 are nodes, which can only deform to smooth points. We therefore assume that the fibers of f have normal crossings in codimension one.

A.2. Constructibility.

Lemma A.2.1. *Under the assumptions of Proposition A.1.1, the locus B_{slc} is constructible in B .*

Proof. The main ingredient is resolution of singularities. The image of a constructible set is constructible, so we may replace B by a resolution of singularities of its normalization $B' \rightarrow B$. We have the pull back family

$$X' = X \times_B B' \rightarrow B'.$$

We claim we can stratify B' by locally-closed subsets

$$B' = \coprod_{j \in J} B'_j$$

such that the restriction over each stratum

$$f_j : X'_j = X'|_{B'_j} \rightarrow B'_j$$

admits a simultaneous good resolution of singularities. By definition, a good resolution of singularities is one where the preimage of the singular locus is a simple normal crossings divisor; in a simultaneous good resolution, each intersection of components of the simple normal crossings divisor is smooth over the base.

The stratification is constructed inductively: Choose a good resolution of singularities $W \rightarrow X'$, which induces a good resolution of the geometric generic fiber of $X' \rightarrow B'$. Since we are working over a field of characteristic zero, such fiber is smooth. We can choose a dense open subset $B'' \subset B'$ such that $W \rightarrow X'$ is a simultaneous good resolution of the fibers X'_b for $b \in B''$. However, $B' \setminus B'' \subsetneq B'$ is closed, so we are done by Noetherian induction.

The fibers of f_j have invertible dualizing sheaves, so Lemma A.1.3 allows us to determine whether the fibers have SLC singularities by computing discrepancies on the good resolutions. However, discrepancies are constant over families with a simultaneous good resolution, so if one fiber of f_j is SLC then all fibers are SLC. In particular, the fibers are SLC over a constructible subset of the base. \square

A.3. Proof of Proposition. We complete the proof that B_{slc} is open by proving that it is stable under one-parameter generalizations. Precisely, let T be nonsingular connected curve, $t \in T$ a closed point, $\mu : T \rightarrow B$ a morphism, and

$$f_T : X_T = X \times_B T \rightarrow T$$

the base change of our family to T . Then the set

$$\{s \in T : X_s \text{ is semilog canonical}\}$$

is open. The existence of the sheaf \mathcal{L} guarantees that the canonical class of X_T is \mathbb{Q} -Cartier.

Suppose that $s \in T$ is such that X_s has semilog canonical singularities. Let $\nu : \tilde{X}_T \rightarrow X_T$ denote the normalization and $g : \tilde{X} \rightarrow T$ the induced morphism. Let C denote the conductor divisor, which is also flat over T . We may write

$$\nu^* K_X = K_{\tilde{X}} + C,$$

which is also \mathbb{Q} -Cartier. If S is the normalization of some irreducible component of \tilde{X}_s and $\iota : S \rightarrow \tilde{X}_s$ the induced morphism, then

$$\iota^*(K_{\tilde{X}_s} + C_s) = K_S + \Theta,$$

where Θ contains the conductor and the preimage of C_s . Lemma A.1.3 implies that (S, Θ) is log canonical, and therefore that (\tilde{X}_s, C_s) is semilog canonical. By ‘Inversion of Adjunction’ [Kaw07], the pair $(\tilde{X}, \tilde{X}_s + C)$ is log canonical as well, at least in a neighborhood of \tilde{X}_s . Fix a good log resolution

$$\rho : (Y, D) \rightarrow (\tilde{X}, \tilde{X}_s + C),$$

i.e., one where the union of the exceptional locus and the proper transform of the boundary is simple normal crossings. All discrepancies are computed with respect to this resolution. Since \tilde{X}_s is a fiber of $\tilde{X} \rightarrow T$, we have

- the log discrepancies of (\tilde{X}, C) for exceptional divisors with center in \tilde{X}_s are ≥ 0 ;
- the log discrepancies of (\tilde{X}, C) for exceptional divisors dominating T are ≥ -1 ;

For $t \neq s$ in a neighborhood of s , the fiber \tilde{X}_t is normal. Furthermore, ρ induces a good log resolution of (\tilde{X}_t, C_t) . Consider the pairs $(\tilde{X}, \tilde{X}_t + C)$, which are also log canonical by the discrepancy analysis above. Applying ‘Adjunction’, we conclude that (\tilde{X}_t, C_t) is also log canonical. Lemma A.1.3 implies that X_t is therefore semilog canonical. \square

References

- [Abr02] Dan Abramovich, Canonical models and stable reduction for plurifibered varieties. Unpublished, [arXiv:math/0207004](https://arxiv.org/abs/math/0207004), 2002.
- [ACV03] Dan Abramovich, Alessio Corti, and Angelo Vistoli, Twisted bundles and admissible covers. *Comm. Algebra* **31** (2003), 3547–3618. Special issue in honor of Steven L. Kleiman.
- [AGV08] Dan Abramovich, Tom Graber, and Angelo Vistoli, Gromov-Witten theory of Deligne-Mumford stacks. *Amer. J. Math.* **130** (2008), 1337–1398.
- [AJ03] Dan Abramovich and Tyler J. Jarvis, Moduli of twisted spin curves, *Proc. Amer. Math. Soc.* **131** (2003), 685–699.
- [Ale94] Valery Alexeev, Boundedness and K^2 for log surfaces, *Internat. J. Math.* **5** (1994), 779–810.

- [AOV08a] Dan Abramovich, Martin Olsson, and Angelo Vistoli, Tame stacks in positive characteristic. *Ann. Inst. Fourier (Grenoble)* **58** (2008), 1057–1091.
- [AOV08b] Dan Abramovich, Martin Olsson, and Angelo Vistoli, Twisted stable maps to tame Artin stacks. preprint [arXiv:0801.3040v2](https://arxiv.org/abs/0801.3040v2), 2008, to appear in *J. Algebraic Geom.*
- [Art74] M. Artin, Versal deformations and algebraic stacks. *Invent. Math.* **27** (1974), 165–189.
- [AV00] Dan Abramovich and Angelo Vistoli, Complete moduli for fibered surfaces. In *Recent progress in intersection theory (Bologna, 1997)*, Trends Math., pages 1–31. Birkhäuser Boston, Boston, MA, 2000.
- [AV02] Dan Abramovich and Angelo Vistoli, Compactifying the space of stable maps. *J. Amer. Math. Soc.* **15** (2002), 27–75.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan, Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.* **23** (2010), 405–468.
- [Con00] Brian Conrad, *Grothendieck duality and base change*, Lecture Notes in Mathematics, Vol. 1750. Springer-Verlag, Berlin, 2000.
- [Elk81] Renée Elkik, Rationalité des singularités canoniques. *Invent. Math.* **64** (1981), 1–6.
- [Gro66] A. Grothendieck, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III. *Inst. Hautes Études Sci. Publ. Math.* **28** (1966).
- [Hac04] Paul Hacking, Compact moduli of plane curves. *Duke Math. J.* **124** (2004), 213–257.
- [HK04] Brendan Hassett and Sándor J. Kovács, Reflexive pull-backs and base extension. *J. Algebraic Geom.* **13** (2004), 233–247.
- [IR96] Yukari Ito and Miles Reid, The McKay correspondence for finite subgroups of $SL(3, \mathbf{C})$. In *Higher-dimensional complex varieties (Trento, 1994)*, pages 221–240. de Gruyter, Berlin, 1996.
- [Jar00] Tyler J. Jarvis, Geometry of the moduli of higher spin curves. *Internat. J. Math.* **11** (2000), 637–663.
- [Kar00] Kalle Karu, Minimal models and boundedness of stable varieties. *J. Algebraic Geom.* **9** (2000), 93–109.
- [Kaw99a] Yujiro Kawamata, Deformations of canonical singularities. *J. Amer. Math. Soc.* **12** (1999), 85–92.
- [Kaw99b] Yujiro Kawamata, Index 1 covers of log terminal surface singularities, 1999.
- [Kaw07] Masayuki Kawakita, Inversion of adjunction on log canonicity. *Invent. Math.* **167** (2007), 129–133.
- [KK09] János Kollár and Sandor J. Kovács, Log canonical singularities are Du Bois, 2009. preprint [arXiv:0902.0648](https://arxiv.org/abs/0902.0648).
- [KKMSD73] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings. I*. Lecture Notes in Mathematics, Vol. 339. Springer-Verlag, Berlin, 1973.

- [KM97] Seán Keel and Shigefumi Mori, Quotients by groupoids. *Ann. of Math. (2)* **145** (1997), 193–213.
- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kol86] János Kollár, Higher direct images of dualizing sheaves. I. *Ann. of Math. (2)* **123** (1986), 11–42.
- [Kol90] János Kollár, Projectivity of complete moduli. *J. Differential Geom.* **32** (1990), 235–268.
- [Kol92] János Kollár, editor, *Flips and abundance for algebraic threefolds*. Astérisque No. 211, Société Mathématique de France, Paris, 1992. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991.
- [Kol95] János Kollár, Flatness criteria. *J. Algebra* **175** (1995), 715–727.
- [Kol08] János Kollár, Hulls and husks. preprint [arXiv:0801.3040v2](https://arxiv.org/abs/0801.3040v2), 2008.
- [KSB88] J. Kollár and N. I. Shepherd-Barron, Threefolds and deformations of surface singularities. *Invent. Math.* **91** (1988), 299–338.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly, *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [Nak86] Noboru Nakayama, Invariance of the plurigenera of algebraic varieties under minimal model conjectures. *Topology* **25** (1986), 237–251.
- [Nir08] Fabio Nironi, Grothendieck duality for Deligne-Mumford stacks. preprint [arXiv:0811955v2](https://arxiv.org/abs/0811955v2), 2008.
- [OS03] Martin Olsson and Jason Starr, Quot functors for Deligne-Mumford stacks. *Comm. Algebra* **31** (2003), 4069–4096. Special issue in honor of Steven L. Kleiman.
- [Rei78] Miles Reid, Surfaces with $p_g = 0$, $K^2 = 1$. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **25** (1978), 75–92.
- [Rei80] Miles Reid, Canonical 3-folds. In *Journées de Géométrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry, Angers, 1979*, pages 273–310. Sijthoff & Noordhoff, Alphen aan den Rijn, 1980.
- [Rei87] Miles Reid, Young person's guide to canonical singularities. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 345–414. Amer. Math. Soc., Providence, RI, 1987.
- [Rei97] Miles Reid, Chapters on algebraic surfaces. In *Complex algebraic geometry (Park City, UT, 1993)*, volume 3 of *IAS/Park City Math. Ser.*, pages 3–159. Amer. Math. Soc., Providence, RI, 1997.
- [Rei02] Miles Reid, Graded rings and varieties in weighted projective space, 2002. unpublished chapter from an upcoming book on surfaces, available at <http://www.warwick.ac.uk/~masda/surf/more/grad.pdf>.

- [Rom05] Matthieu Romagny, Group actions on stacks and applications. *Michigan Math. J.* **53** (2005), 209–236.
- [RT09] Julius Ross and Richard Thomas, Weighted projective embeddings, stability of orbifolds and constant scalar curvature Kähler metrics. preprint [arXiv:0907.5412](https://arxiv.org/abs/0907.5412), 2009.
- [Sin03] Anurag K. Singh, Cyclic covers of rings with rational singularities. *Trans. Amer. Math. Soc.* **355** (2003), 1009–1024.
- [Vie95] Eckart Viehweg, *Quasi-projective moduli for polarized manifolds*, volume 30 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1995.

Department of Mathematics, Box 1917, Brown University, Providence, RI, 02912, U.S.A

E-mail: abrmovic@math.brown.edu

Department of Mathematics, MS 136, Rice University, Houston, TX 77251, U.S.A

E-mail: hassett@math.rice.edu