

Resolution and logarithmic resolution by weighted blowing up

Dan Abramovich, Brown University

Work with Michael Tëmkin and Jarosław Włodarczyk
and work by Ming Hao Quek

Also parallel work by M. McQuillan

Algebraic geometry and Moduli Seminar

J. Lee Anne Fröhbers - Krüger

ETH Zürich, July 15, 2020

How to resolve

To resolve a singular variety X one wants to

- (1) find the worst singular locus $S \subset X$,
- (2) Hopefully S is smooth - blow it up.

How to resolve

To resolve a singular variety X one wants to

- (1) find the worst singular locus $S \subset X$,
- (2) Hopefully S is smooth - blow it up.

Fact

This works for curves but not in general.

Example: Whitney's umbrella

Consider $X = V(x^2 - y^2z)$

Example: Whitney's umbrella

Consider $X = V(x^2 - y^2z)$

(1) The worst singularity is the origin.

(2) In the z chart we get

$x = x'z$, $y = y'z$, giving

$$x'^2z^2 - y'^2z^3 = 0, \quad \text{or} \quad z^2(x'^2 - y'^2z) = 0.$$

Example: Whitney's umbrella

Consider $X = V(x^2 - y^2z)$

(1) The worst singularity is the origin.

(2) In the z chart we get

$x = x'z$, $y = y'z$, giving

$$x'^2z^2 - y'^2z^3 = 0, \quad \text{or} \quad z^2(x'^2 - y'^2z) = 0.$$

The **first term** is exceptional, the **second** is the same as X .

Two theorems

Nevertheless:

Theorem (N-T-W, McQuillan, 2019, characteristic 0)

There is a *functor* F associating to a singular subvariety $X \subset Y$ of a smooth variety Y , a *center* \bar{J} with *stack theoretic weighted blowing up* $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\text{maxinv}(X') < \text{maxinv}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n smooth.

Two theorems

Nevertheless:

Theorem (N-T-W, McQuillan, 2019, characteristic 0)

There is a **functor** F associating to a singular subvariety $X \subset Y$ of a smooth variety Y , a **center** \bar{J} with **stack theoretic weighted blowing up** $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\max_{\text{inv}}(X') < \max_{\text{inv}}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n smooth.

Theorem (Quek, 2020, characteristic 0)

There is a functor F associating to a **logarithmically** singular subvariety $X \subset Y$ of a **logarithmically** smooth variety Y , a **logarithmic** center \bar{J} with stack theoretic **logarithmic** blowing up $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\max_{\text{loginv}}(X') < \max_{\text{loginv}}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n **logarithmically smooth**.

- context
- definitions

~ ideas of proofs
- indicate differences

Context: families

Hironaka's theorem resolves varieties. What can you do with families of varieties $X \rightarrow B$?

Theorem (Kawamata-Matsuda-Mumford, 1977)

There is a modification $X' \rightarrow B'$ which is *logarithmically smooth*.

Context: families

Hironaka's theorem resolves varieties. What can you do with families of varieties $X \rightarrow B$?

Theorem (Kawamata-Matsuda-Matsuki, 1981)

There is a modification $X' \rightarrow B'$ which is *logarithmically smooth*.

Logarithmically smooth = toroidal:

Context: families

Hironaka's theorem resolves varieties. What can you do with families of varieties $X \rightarrow B$?

Theorem (Kawamata-Matsuda-Matsuki, 1981)

There is a modification $X' \rightarrow B'$ which is *logarithmically smooth*.

Logarithmically smooth = toroidal:

- A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.

Context: families

Hironaka's theorem resolves varieties. What can you do with families of varieties $X \rightarrow B$?

Theorem (Kawakita-Karu, 2000)

There is a modification $X' \rightarrow B'$ which is *logarithmically smooth*.

Logarithmically smooth = toroidal:

- A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.
- A **toroidal embedding** $U_X \subset X$ is an open embedding étale locally isomorphic to toric $T \subset V$.

$K \subset \mu \subset S \subset \Delta$

Context: families

Hironaka's theorem resolves varieties. What can you do with families of varieties $X \rightarrow B$?

Theorem (Karu, 2000)

There is a modification $X' \rightarrow B'$ which is *logarithmically smooth*.

Logarithmically smooth = toroidal:

- A **toric** morphism $X \rightarrow B$ of toric varieties is a *dominant* torus equivariant morphism.
- A **toroidal embedding** $U_X \subset X$ is an open embedding étale locally isomorphic to toric $T \subset V$.
- A **toroidal** morphism $X \rightarrow B$ of toroidal embeddings is étale locally isomorphic to a toric morphism.

Examples of toroidal morphisms

A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.

Examples of toroidal morphisms

A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.e.g.



$$\mathrm{Spec} \mathbb{C}[x, y, z]/(xy - z^2) \rightarrow \mathrm{Spec} \mathbb{C},$$

Examples of toroidal morphisms

A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.e.g.



$$\mathrm{Spec} \mathbb{C}[x, y, z]/(xy - z^2) \rightarrow \mathrm{Spec} \mathbb{C},$$



$$\mathrm{Spec} \mathbb{C}[x] \rightarrow \mathrm{Spec} \mathbb{C}[x^2],$$

Examples of toroidal morphisms

A **toric** morphism $X \rightarrow B$ of toric varieties is a torus equivariant morphism.e.g.

- $$\text{Spec } \mathbb{C}[x, y, z]/(xy - z^2) \rightarrow \text{Spec } \mathbb{C},$$

- $$\text{Spec } \mathbb{C}[x] \rightarrow \text{Spec } \mathbb{C}[x^2],$$

- toric blowups



Context: functoriality

- Hironaka's theorem is functorial.
- [Karu 2000] is not: relied on deJong's method.

Context: functoriality

- Hironaka's theorem is functorial.
- [Karu 2000] is not: relied on deJong's method.
- For K-S-B or K-moduli want functoriality.

Context: functoriality

- Hironaka's theorem is functorial.
- [Karu 2000] is not: relied on deJong's method.
- For K-S-B or K-moduli want functoriality.

Theorem (T-W 2020)

Given $X \rightarrow B$ there is a *relatively functorial* logarithmically smooth modification $X' \rightarrow B'$.

Context: functoriality

- Hironaka's theorem is functorial.
- [Karu 2000] is not: relied on deJong's method.
- For K-S-B or K-moduli want functoriality.

Theorem (T-W 2020)

Given $X \rightarrow B$ there is a *relatively functorial* logarithmically smooth modification $X' \rightarrow B'$.

- This respects $\text{Aut}_B X$.
- Does not modify log smooth fibers.

Context: principalization

- Following Hironaka, the above theorem is based on embedded methods:

Theorem (N-T-W 2020)

Given $Y \rightarrow B$ logarithmically smooth and $\mathcal{I} \subset \mathcal{O}_Y$, there is a relatively functorial logarithmically smooth modification $Y' \rightarrow B'$ such that $\mathcal{I}\mathcal{O}_{Y'}$ is *monomial*.

Context: principalization


- Following Hironaka, the above theorem is based on embedded methods:

Theorem (K-T-W 2020)

Given $Y \rightarrow B$ logarithmically smooth and $\mathcal{I} \subset \mathcal{O}_Y$, there is a relatively functorial logarithmically smooth modification $Y' \rightarrow B'$ such that $\mathcal{I}\mathcal{O}_{Y'}$ is *monomial*.

- This is done by a sequence of logarithmic modifications,
- where in each step E becomes part of the divisor $D_{Y'}$.

Example 1





- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad B = \text{Spec } k;$

Example 1

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad B = \text{Spec } k; \quad \mathcal{I} = (x^2, u^2).$

Example 1

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad B = \text{Spec } k; \quad \mathcal{I} = (x^2, u^2).$
- Blow up $J = (x, u)$
- $\mathcal{I}\mathcal{O}_{Y'} = \mathcal{O}(-2E)$



Example 1/2

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad \mathcal{I} = (x^2, u^2)$

Example 1/2

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad \mathcal{I} = (x^2, u^2)$
- $Y_0 = \text{Spec } k[x, v]; \quad D_{Y_0} = V(v); \quad \mathcal{I}_0 = (x^2, v)$
- $f : Y \rightarrow Y_0 \quad v = u^2 \quad \text{so} \quad \mathcal{I} = f^*\mathcal{I}_0$

Example 1/2

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad \mathcal{I} = (x^2, u^2)$
- $Y_0 = \text{Spec } k[x, v]; \quad D_{Y_0} = V(v); \quad \mathcal{I}_0 = (x^2, v),$
- $f : Y \rightarrow Y_0 \quad v = u^2 \quad \text{so} \quad \mathcal{I} = f^*\mathcal{I}_0$
- By functoriality blow up J_0 so that $f^*J_0 = J = (x, u).$

Example 1/2

- $Y = \text{Spec } k[x, u]; \quad D_Y = V(u); \quad \mathcal{I} = (x^2, u^2)$
- $Y_0 = \text{Spec } k[x, v]; \quad D_{Y_0} = V(v); \quad \mathcal{I}_0 = (x^2, v),$
- $f : Y \rightarrow Y_0 \quad v = u^2 \quad \text{so} \quad \mathcal{I} = f^*\mathcal{I}_0$
- By functoriality blow up J_0 so that $f^*J_0 = J = (x, u)$.
- Blow up $J_0 = (x, \sqrt{v})$
- Whatever J_0 is, the blowup is a stack.

Example 1/2: charts

$$(x, \sqrt{v})$$

- **x chart:** $v = v'x^2$:

$$(x^2, v) = (x^2, v'x^2) = (x^2)$$

exceptional, so monomial.


- **\sqrt{v} chart:** $v = w^2$, $x = x'w$, with ± 1 action $(x', w) \mapsto (-x', -w)$:

$$(x^2, v) = (x'^2 w^2, w^2) = (w^2)$$

$$\left[\text{Spec } k[x', w] / \pm 1 \right]$$

exceptional, so monomial.

- The schematic quotient of the above is **not toroidal**.



Resolution again

Theorem (K-T-W, McQuillan, characteristic 0)

There is a *functor* F associating to a singular subvariety $X \subset Y$ of a smooth variety Y , a *center* J with *stack theoretic weighted blowing up* $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\text{maxinv}(X') < \text{maxinv}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n smooth.

Resolution again

Theorem (K-T-W, McQuillan, characteristic 0)

There is a *functor* F associating to a singular subvariety $X \subset Y$ of a smooth variety Y , a *center* \bar{J} with *stack theoretic weighted blowing up* $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\text{maxinv}(X') < \text{maxinv}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n smooth.

Example

For $X = V(x^2 - y^2z)$ we have $\text{inv}_p(X) = (2, 3, 3)$

Resolution again

Theorem (N-T-W, McQuillan, characteristic 0)

There is a functor F associating to a singular subvariety $X \subset Y$ of a smooth variety Y , a center \bar{J} with stack theoretic weighted blowing up $Y' \rightarrow Y$ and proper transform $(X' \subset Y') = F(X \subset Y)$ such that $\max \text{inv}(X') < \max \text{inv}(X)$. In particular, for some n the iterate $(X_n \subset Y_n) := F^{\circ n}(X \subset Y)$ of F has X_n smooth.

Example

For $X = V(x^2 - y^2z)$ we have $\text{inv}_p(X) = (2, 3, 3)$

We read it from the degrees of terms.

The center is:

$$J = (x^2, y^3, z^3); \bar{J} = (x^{1/3}, y^{1/2}, z^{1/2}).$$

Example: blowing up Whitney's umbrella $x^2 = y^2z$

The blowing up $Y' \rightarrow Y$ makes $\bar{J} = (x^{1/3}, y^{1/2}, z^{1/2})$ principal. Explicitly:

- The z chart has $x = w^3x_3, y = w^2y_3, z = w^2$ with chart

$$Y' = [\text{Spec } \mathbb{C}[x_3, y_3, w] / (\pm 1)],$$

with action of (± 1) given by $(x_3, y_3, w) \mapsto (-x_3, y_3, -w)$.

Example: blowing up Whitney's umbrella $x^2 = y^2z$

The blowing up $Y' \rightarrow Y$ makes $\bar{J} = (x^{1/3}, y^{1/2}, z^{1/2})$ principal. Explicitly:

- The z chart has $x = w^3x_3, y = w^2y_3, z = w^2$ with chart

$$Y' = [\text{Spec } \mathbb{C}[x_3, y_3, w] / (\pm 1)],$$

with action of (± 1) given by $(x_3, y_3, w) \mapsto (-x_3, y_3, -w)$.

The transformed equation is

$$w^6(x_3^2 - y_3^2),$$

Example: blowing up Whitney's umbrella $x^2 = y^2z$

The blowing up $Y' \rightarrow Y$ makes $\bar{J} = (x^{1/3}, y^{1/2}, z^{1/2})$ principal. Explicitly:

- The z chart has $x = w^3x_3, y = w^2y_3, z = w^2$ with chart

$$Y' = [\text{Spec } \mathbb{C}[x_3, y_3, w] / (\pm 1)],$$

with action of (± 1) given by $(x_3, y_3, w) \mapsto (-x_3, y_3, -w)$.

The transformed equation is

$$w^6(x_3^2 - y_3^2),$$

and the invariant of the proper transform $(x_3^2 - y_3^2)$ is $(2, 2) < (2, 3, 3)$.

Order (following Kollár's book)

We fix Y smooth and $\mathcal{I} \subset \mathcal{O}_Y$.

Definition

For $p \in Y$ let $\text{ord}_p(\mathcal{I}) = \max\{a : \mathcal{I} \subseteq \mathfrak{m}_p^a\}$.

Order (following Kollár's book)

We fix Y smooth and $\mathcal{I} \subset \mathcal{O}_Y$.

Definition

For $p \in Y$ let $\text{ord}_p(\mathcal{I}) = \max\{a : \mathcal{I} \subseteq \mathfrak{m}_p^a\}$.

- We denote by \mathcal{D}^a the sheaf of a -th order differential operators.

Order (following Kollár's book)

We fix Y smooth and $\mathcal{I} \subset \mathcal{O}_Y$.

Definition

For $p \in Y$ let $\text{ord}_p(\mathcal{I}) = \max\{a : \mathcal{I} \subseteq \mathfrak{m}_p^a\}$.

- We denote by \mathcal{D}^a the sheaf of a -th order differential operators.
- We note that $\text{ord}_p(\mathcal{I}) = \min\{a : \mathcal{D}^a(\mathcal{I}_p)\} = (1)$.
- The invariant starts with $a_1 = \text{ord}_p(\mathcal{I})$.

Order (following Kollár's book)

We fix Y smooth and $\mathcal{I} \subset \mathcal{O}_Y$.

Definition

For $p \in Y$ let $\text{ord}_p(\mathcal{I}) = \max\{a : \mathcal{I} \subseteq \mathfrak{m}_p^a\}$.

- We denote by \mathcal{D}^a the sheaf of a -th order differential operators.
- We note that $\text{ord}_p(\mathcal{I}) = \min\{a : \mathcal{D}^a(\mathcal{I}_p)\} = (1)$.
- The invariant starts with $a_1 = \text{ord}_p(\mathcal{I})$.

Proposition

The order is upper semicontinuous.

Proof.

$$V(\mathcal{D}^{a-1}\mathcal{I}) = \{p : \text{ord}_p(\mathcal{I}) \geq a\}.$$



Maximal contact (following Kollár's book)

Definition

A regular parameter $x_1 \in \mathcal{D}^{a_1-1}\mathcal{I}_p$ is called a **maximal contact** element.

The center starts with $(x_1^{a_1}, \dots)$.

Maximal contact (following Kollár's book)

Definition

A regular parameter $x_1 \in \mathcal{D}^{a_1-1}\mathcal{I}_p$ is called a **maximal contact** element.

The center starts with $(x_1^{a_1}, \dots)$.

Lemma (Hironaka, Giraud)

In characteristic 0 a maximal contact exists on an open neighborhood of p .

Since $1 \in \mathcal{D}^{a_1}\mathcal{I}_p$ there is x_1 with derivative 1. This derivative is a unit in a neighborhood.

$$\mathcal{I} = (x^p)$$

Maximal contact (following Kollár's book)

Definition

A regular parameter $x_1 \in \mathcal{D}^{a_1-1}\mathcal{I}_p$ is called a **maximal contact** element.

The center starts with $(x_1^{a_1}, \dots)$.

Lemma (Hironaka, Giraud)

In characteristic 0 a maximal contact exists on an open neighborhood of p .

Since $1 \in \mathcal{D}^{a_1}\mathcal{I}_p$ there is x_1 with derivative 1. This derivative is a unit in a neighborhood.

Example

For $\mathcal{I} = (x^2 - y^2z)$ we have $\text{ord}_p \mathcal{I} = 2$ with $x_1 = x$
(or $\alpha x + \text{h.o.t.}$ in $\mathcal{D}(\mathcal{I})$).

Coefficient ideals (treated following Kollár)

We must restrict to $x_1 = 0$ the data of all

$$\mathcal{I}, \mathcal{D}\mathcal{I}, \dots, \mathcal{D}^{a_1-1}\mathcal{I}$$

with corresponding weights $a_1, a_1 - 1, \dots, 1$.

We combine these in

$$C(\mathcal{I}, a_1) := \sum f(\mathcal{I}, \mathcal{D}\mathcal{I}, \dots, \mathcal{D}^{a_1-1}\mathcal{I}),$$

where f runs over monomials $f = t_0^{b_0} \cdots t_{a_1-1}^{b_{a_1-1}}$ with weights

$$\sum b_i(a_1 - i) \geq a_1!.$$

Define $\mathcal{I}[2] = C(\mathcal{I}, a_1)|_{x_1=0}$.

Defining $J_{\mathcal{I}}$

Again $a_1 = \text{ord}_p \mathcal{I}$ and x_1 maximal contact.

We denoted $\mathcal{I}[2] = C(\mathcal{I}, a_1)|_{x_1=0}$ (with order $\geq a_1!$).

Defining $J_{\mathcal{I}}$

Again $a_1 = \text{ord}_p \mathcal{I}$ and x_1 maximal contact.

We denoted $\mathcal{I}[2] = C(\mathcal{I}, a_1)|_{x_1=0}$ (with order $\geq a_1!$).

Definition

Suppose $\mathcal{I}[2]$ has invariant $\text{inv}_p(\mathcal{I}[2])$ defined with parameters $\bar{x}_2, \dots, \bar{x}_k$, with lifts x_2, \dots, x_k .

Defining $J_{\mathcal{I}}$

Again $a_1 = \text{ord}_p \mathcal{I}$ and x_1 maximal contact.

We denoted $\mathcal{I}[2] = C(\mathcal{I}, a_1)|_{x_1=0}$ (with order $\geq a_1!$).

Definition

Suppose $\mathcal{I}[2]$ has invariant $\text{inv}_p(\mathcal{I}[2])$ defined with parameters $\bar{x}_2, \dots, \bar{x}_k$, with lifts x_2, \dots, x_k . Set

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1 - 1)!} \right)$$

and

$$J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Questi:
need to
consider
variables

Defining $J_{\mathcal{I}}$

Again $a_1 = \text{ord}_p \mathcal{I}$ and x_1 maximal contact.

We denoted $\mathcal{I}[2] = C(\mathcal{I}, a_1)|_{x_1=0}$ (with order $\geq a_1!$).

Definition

Suppose $\mathcal{I}[2]$ has invariant $\text{inv}_p(\mathcal{I}[2])$ defined with parameters $\bar{x}_2, \dots, \bar{x}_k$, with lifts x_2, \dots, x_k . Set

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1 - 1)!} \right)$$

and

$$J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Write $(a_1, \dots, a_k) = \ell(1/w_1, \dots, 1/w_k)$ with $w_i, \ell \in \mathbb{N}$ and $\text{gcd}(w_1, \dots, w_k) = 1$. We set

$$\bar{J}_{\mathcal{I}} = (x_1^{1/w_1}, \dots, x_k^{1/w_k}).$$

Examples of $J_{\mathcal{I}}$

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1-1)!} \right), \quad \text{with } J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Example

(0) for $X = V(x^2 + y^2z)$

$$(x^2, y^3, z^3) \quad (2, 3, 3)$$
$$C(\mathbb{I}, 2) \Big|_{x=0} = (y^2, z^2)$$

Examples of $J_{\mathcal{I}}$

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1-1)!} \right), \quad \text{with} \quad J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Example

(0) for $X = V(x^2 + y^2z)$ we have $\mathcal{I}[2] = (y^2z)$, leading to

$$J_{\mathcal{I}} = (x^2, y^3, z^3), \quad \bar{J}_{\mathcal{I}} = (x^{1/3}, y^{1/2}, z^{1/2})$$

(1) for $X = V(x^5 + x^3y^3 + y^8)$

$$(x^5, y^{5/2})$$
$$\mathcal{I}[2] = (y^{180}) \quad \frac{1}{(5 \cdot 1)!}$$

Examples of $J_{\mathcal{I}}$

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1-1)!} \right), \quad \text{with} \quad J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Example

- (0) for $X = V(x^2 + y^2z)$ we have $\mathcal{I}[2] = (y^2z)$, leading to
 $J_{\mathcal{I}} = (x^2, y^3, z^3)$, $\bar{J}_{\mathcal{I}} = (x^{1/3}, y^{1/2}, z^{1/2})$
- (1) for $X = V(x^5 + x^3y^3 + y^8)$ we have $\mathcal{I}[2] = (y)^{180}$, so
 $J_{\mathcal{I}} = (x^5, y^{180/24}) = (x^5, y^{15/2})$, $\bar{J}_{\mathcal{I}} = (x^{1/3}, y^{1/2})$.
- (2) for $X = V(x^5 + x^3y^3 + y^7)$

Examples of $J_{\mathcal{I}}$

$$\text{inv}_p(\mathcal{I}) = (a_1, \dots, a_k) := \left(a_1, \frac{\text{inv}_p(\mathcal{I}[2])}{(a_1-1)!} \right), \quad \text{with} \quad J_{\mathcal{I}} = (x_1^{a_1}, \dots, x_k^{a_k}).$$

Example

- (0) for $X = V(x^2 + y^2z)$ we have $\mathcal{I}[2] = (y^2z)$, leading to $J_{\mathcal{I}} = (x^2, y^3, z^3)$, $\bar{J}_{\mathcal{I}} = (x^{1/3}, y^{1/2}, z^{1/2})$
- (1) for $X = V(x^5 + x^3y^3 + y^8)$ we have $\mathcal{I}[2] = (y)^{180}$, so $J_{\mathcal{I}} = (x^5, y^{180/24}) = (x^5, y^{15/2})$, $\bar{J}_{\mathcal{I}} = (x^{1/3}, y^{1/2})$.
- (2) for $X = V(x^5 + x^3y^3 + y^7)$ we have $\mathcal{I}[2] = (y)^{7 \cdot 24}$, so $J_{\mathcal{I}} = (x^5, y^7)$, $\bar{J}_{\mathcal{I}} = (x^{1/7}, y^{1/5})$.

Implementation: Jonghyun Lee, Anne Frühbis-Krüger.

Properties of the invariant

Proposition

- inv_p is well-defined.
- inv_p is lexicographically upper-semi-continuous.
- inv_p is functorial.
- inv_p takes values in a well-ordered set.

We define $\text{maxinv}(X) = \max_p \text{inv}_p(X)$.

Properties of the invariant

Proposition

- inv_p is well-defined.
- inv_p is lexicographically upper-semi-continuous.
- inv_p is functorial.
- inv_p takes values in a well-ordered set.

We define $\text{maxinv}(X) = \max_p \text{inv}_p(X)$.

The invariant is well defined because of the **MC-invariance** property of $C(\mathcal{I}, a_1)$. The rest is induction!

Properties of the invariant

Proposition

- inv_p is well-defined.
- inv_p is lexicographically upper-semi-continuous.
- inv_p is functorial.
- inv_p takes values in a well-ordered set.

We define $\text{maxinv}(X) = \max_p \text{inv}_p(X)$.

The invariant is well defined because of the **MC-invariance** property of $C(\mathcal{I}, a_1)$. The rest is induction!

Theorem (MC-invariance [Włodarczyk, Kollár])

Given maximal contacts x_1, x'_1 there are étale $\pi, \pi' : \tilde{Y} \rightrightarrows Y$ such that $\pi^* x_1 = \pi'^* x'_1 \dots$ and $\pi^* C(\mathcal{I}, a_1) = \pi'^* C(\mathcal{I}, a_1)$.

Definition of $Y' \rightarrow Y$

Let $\bar{J} = (x_1^{1/w_1}, \dots, x_k^{1/w_k})$. Define the graded algebra

$$\mathcal{A}_{\bar{J}} \subset \mathcal{O}_Y[T]$$

as the integral closure of the image of

$$\begin{array}{ccc} \mathcal{O}_Y[Y_1, \dots, Y_n] & \longrightarrow & \mathcal{O}_Y[T] \\ Y_i & \longmapsto & x_i T^{w_i}. \end{array}$$

Definition of $Y' \rightarrow Y$

Let $\bar{J} = (x_1^{1/w_1}, \dots, x_k^{1/w_k})$. Define the graded algebra

$$\mathcal{A}_{\bar{J}} \subset \mathcal{O}_Y[T]$$

as the integral closure of the image of

$$\begin{aligned} \mathcal{O}_Y[Y_1, \dots, Y_n] &\longrightarrow \mathcal{O}_Y[T] \\ Y_i &\longmapsto x_i T^{w_i}. \end{aligned}$$

Let

$$S_0 \subset \operatorname{Spec}_Y \mathcal{A}_{\bar{J}}, \quad S_0 = V((\mathcal{A}_{\bar{J}})_{>0}).$$

Then

$$Bl_{\bar{J}}(Y) := \operatorname{Proj}_Y \mathcal{A}_{\bar{J}} := [(\operatorname{Spec} \mathcal{A}_{\bar{J}} \setminus S_0) / \mathbb{G}_m].$$

Description of $Y' \rightarrow Y$

- **Charts:** The x_1 -chart is

$$[\mathrm{Spec} k[u, x'_2, \dots, x'_n] / \mu_{w_1}],$$

with $x_1 = u^{w_1}$ and $x_i = u^{w_i} x'_i$ for $2 \leq i \leq k$, and induced action:

$$(u, x'_2, \dots, x'_n) \mapsto (\zeta u, \zeta^{-w_2} x'_2, \dots, \zeta^{-w_k} x'_k, x'_{k+1}, \dots, x'_n).$$

Description of $Y' \rightarrow Y$

- **Charts:** The x_1 -chart is

$$[\mathrm{Spec} k[u, x'_2, \dots, x'_n] / \mu_{w_1}],$$

with $x_1 = u^{w_1}$ and $x_i = u^{w_i} x'_i$ for $2 \leq i \leq k$, and induced action:

$$(u, x'_2, \dots, x'_n) \mapsto (\zeta u, \zeta^{-w_2} x'_2, \dots, \zeta^{-w_k} x'_k, x'_{k+1}, \dots, x'_n).$$

- **Toric stack:** Consider $\mathrm{Spec} k[x_1, \dots, x_n, T]$ with \mathbb{G}_m action with weights $(w_1, \dots, w_n, -1)$. Let U be the open set where one of the x_i is a unit. Then $Y' = [U/\mathbb{G}_m]$.
It is an example of a *fantastack* [Geraschenko-Satriano], the stack quotient of a Cox construction.

What is J ?

Definition

Consider the Zariski-Riemann space $\mathbf{ZR}(Y)$ with its sheaf of ordered groups Γ , and associated sheaf of rational ordered group $\Gamma \otimes \mathbb{Q}$.

- A **valuative \mathbb{Q} -ideal** is

$$\gamma \in H^0(\mathbf{ZR}(Y), (\Gamma \otimes \mathbb{Q})_{\geq 0}).$$

What is J ?

Definition

Consider the Zariski-Riemann space $\mathbf{ZR}(Y)$ with its sheaf of ordered groups Γ , and associated sheaf of rational ordered group $\Gamma \otimes \mathbb{Q}$.

- A **valuative \mathbb{Q} -ideal** is

$$\gamma \in H^0(\mathbf{ZR}(Y), (\Gamma \otimes \mathbb{Q})_{\geq 0}).$$

- $\mathcal{I}_\gamma := \{f \in \mathcal{O}_Y : v(f) \geq \gamma_v \forall v\}$.
- $v(\mathcal{I}) := (\min v(f) : f \in \mathcal{I})_v$.

What is J ?

Definition

Consider the Zariski-Riemann space $\mathbf{ZR}(Y)$ with its sheaf of ordered groups Γ , and associated sheaf of rational ordered group $\Gamma \otimes \mathbb{Q}$.

- A **valuative \mathbb{Q} -ideal** is

$$\gamma \in H^0(\mathbf{ZR}(Y), (\Gamma \otimes \mathbb{Q})_{\geq 0}).$$

- $\mathcal{I}_\gamma := \{f \in \mathcal{O}_Y : v(f) \geq \gamma \forall v\}$.
- $v(\mathcal{I}) := (\min v(f) : f \in \mathcal{I})_v$.

A center is in particular a valuative \mathbb{Q} -ideal. It is also an **idealistic exponent** or **graded sequence of ideals**.

$$v\left(\left(x, y\right)^2\right)_v = v\left(x^2, y^2\right)$$

$$\mathcal{I}_{u, \delta}$$

Admissibility and coefficient ideals

Definition

J is \mathcal{I} -admissible if $J \leq v(\mathcal{I})$.

Admissibility and coefficient ideals

Definition

J is \mathcal{I} -admissible if $J \leq v(\mathcal{I})$.

Lemma

This is equivalent to $\mathcal{I}\mathcal{O}_{Y'} = E^\ell \mathcal{I}'$, with $J = \bar{J}^\ell$ and \mathcal{I}' an ideal.

Indeed, on Y' the center J becomes E^ℓ , in particular principal.
This is more subtle in Quek's theorem!

Admissibility and coefficient ideals

Definition

J is \mathcal{I} -admissible if $J \leq v(\mathcal{I})$.

Lemma

This is equivalent to $\mathcal{I}\mathcal{O}_{Y'} = E^\ell \mathcal{I}'$, with $J = \bar{J}^\ell$ and \mathcal{I}' an ideal.

Indeed, on Y' the center J becomes E^ℓ , in particular principal. This is more subtle in Quek's theorem!

Proposition

A center J is \mathcal{I} -admissible if and only if $J^{(a_1-1)!}$ is $C(\mathcal{I}, a_1)$ -admissible.

The key theorems

Theorem

$\text{inv}_p(\mathcal{I})$ is the maximal invariant of an \mathcal{I} -admissible center.

Theorem

$J_{\mathcal{I}}$ is well-defined: it is the unique admissible center of maximal invariant.

The key theorems

Theorem

$\text{inv}_p(\mathcal{I})$ is the maximal invariant of an \mathcal{I} -admissible center.

Theorem

$J_{\mathcal{I}}$ is well-defined: it is the unique admissible center of maximal invariant.

Theorem

$C(\mathcal{I}, a_1)\mathcal{O}_{Y'} = E^{\ell'} C'$ with $\text{inv}_{p'} C' < \text{inv}_p(C(\mathcal{I}, a_1))$.

The key theorems

Theorem

$\text{inv}_p(\mathcal{I})$ is the maximal invariant of an \mathcal{I} -admissible center.

Theorem

$J_{\mathcal{I}}$ is well-defined: it is the unique admissible center of maximal invariant.

Theorem

$C(\mathcal{I}, a_1)\mathcal{O}_{Y'} = E^{\ell'} C'$ with $\text{inv}_{p'} C' < \text{inv}_p(C(\mathcal{I}, a_1))$.

Theorem

$\mathcal{I}\mathcal{O}_{Y'} = E^{\ell'} \mathcal{I}'$ with $\text{inv}_{p'} \mathcal{I}' < \text{inv}_p(\mathcal{I})$.

This is a consequence of Kollár's \mathcal{D} -balanced property of $C(\mathcal{I}, a_1)$.

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.
- Then $\text{maxinv}(\mathcal{I}) = (4, 4, 4)$ with center $J = (x^4, y^4, z^4)$, a usual blowup.

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.
- Then $\text{maxinv}(\mathcal{I}) = (4, 4, 4)$ with center $J = (x^4, y^4, z^4)$, a usual blowup.
- The z -chart has $\mathcal{I}' = (y(x^2 + z))$. The new invariant is $(2, 2)$ with reduced center $(y, x^2 + z)$, which is tangent to the exceptional $z = 0$.

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.
- Then $\text{maxinv}(\mathcal{I}) = (4, 4, 4)$ with center $J = (x^4, y^4, z^4)$, a usual blowup.
- The z -chart has $\mathcal{I}' = (y(x^2 + z))$. The new invariant is $(2, 2)$ with reduced center $(y, x^2 + z)$, which is tangent to the exceptional $z = 0$.
- **Instead work with logarithmic derivative in z .**

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.
- Then $\text{maxinv}(\mathcal{I}) = (4, 4, 4)$ with center $J = (x^4, y^4, z^4)$, a usual blowup.
- The z -chart has $\mathcal{I}' = (y(x^2 + z))$. The new invariant is $(2, 2)$ with reduced center $(y, x^2 + z)$, which is tangent to the exceptional $z = 0$.
- **Instead work with logarithmic derivative in z .** $y x^2 \neq y z$
- The logarithmic invariant is $(3, 3, \infty)$ with center $(y^3, x^3, z^{3/2})$ and reduced logarithmic center $(y, x, z^{1/2})$.

Quek's theorem is necessary

- $\mathcal{I} = (x^2yz + yz^4) \subset \mathbb{C}[x, y, z]$.
- Then $\text{maxinv}(\mathcal{I}) = (4, 4, 4)$ with center $J = (x^4, y^4, z^4)$, a usual blowup.
- The z -chart has $\mathcal{I}' = (y(x^2 + z))$. The new invariant is $(2, 2)$ with reduced center $(y, x^2 + z)$, which is tangent to the exceptional $z = 0$.
- **Instead work with logarithmic derivative in z .**
- The logarithmic invariant is $(3, 3, \infty)$ with center $(y^3, x^3, z^{3/2})$ and reduced logarithmic center $(y, x, z^{1/2})$.
- This reduces logarithmic invariants respecting logarithmic, hence exceptional, divisors.

The end

Thank you for your attention