

# FUNCTORIAL FACTORIZATION OF BIRATIONAL MAPS FOR QE SCHEMES IN CHARACTERISTIC 0

DAN ABRAMOVICH AND MICHAEL TEMKIN

ABSTRACT. We prove functorial weak factorization of projective birational morphisms of regular quasi-excellent schemes in characteristic 0 based on the existing line of proof for varieties. From this we deduce factorization of any blowing up of formal schemes, complex analytic germs, Berkovich analytic or rigid analytic spaces.

## CONTENTS

1. Introduction	1
2. Qe schemes and functoriality	5
3. Functorial toroidal factorization	11
4. Birational cobordisms	16
5. Factoring the map	21
6. Extending the factorization to other categories	26
Appendix A. Construction of a birational cobordism via deformation to the normal cone	32
Appendix B. Germs of complex analytic spaces	35
Appendix C. The complex relative GAGA Theorem	37
References	41

## 1. INTRODUCTION

1.1. The class of qe schemes (originally “quasi excellent schemes”) is the natural class of schemes on which problems around resolution of singularities are of interest. They can also be used as a bridge for studying the same type of problems in other geometric categories, see [Tem12, Section 5]. In this paper we address the problem of functorial factorization of birational morphisms between regular qe schemes of characteristic 0 into blowings up and down of regular schemes along regular centers. We rely on general foundations developed in [AT15a, AT15b] and the approach for varieties of [Wło00, AKMW02]. As a consequence of both this generality of qe schemes and of functoriality, we are able to deduce factorization of birational or bimeromorphic morphisms in other geometric categories of interest.

---

*Date:* February 17, 2016.

This research is supported by BSF grant 2010255.

Sec: def-factor

**1.2. Blowings up and weak factorizations.** We start with a morphism of noetherian qe regular schemes  $\phi: X_1 \rightarrow X_2$  given as the blowing up of a coherent sheaf of ideals  $I$  on the qe scheme  $X_2$ . In addition, we provide  $\phi$  with a *boundary*  $(D_1, D_2)$ , where each  $D_i$  is a normal crossings divisor in  $X_i$  and  $D_1 := \phi^{-1}D_2$ . Let  $U = X_2 \setminus (D_2 \cup V(I))$  be the maximal open subscheme of  $X_2$  such that  $I$  is the unit ideal on  $U$  and the boundary is disjoint from  $U$ . The restriction of  $\phi$  on  $U$  is the trivial blowing up (i.e. the blowing up of the empty center), in particular, we canonically have an isomorphism  $\phi^{-1}U \rightarrow U$ . We often keep the ideal  $I$  implicit in the notation, even though it determines  $\phi$  (but see Section 2.3.8 for a construction in the reverse direction). The reader may wish to focus on the following two cases of interest: (i)  $D_2 = \emptyset$ ; (ii)  $V(I) \subseteq D_2$ .

A *weak factorization* of a blowing up  $\phi: X_1 \rightarrow X_2$  is a diagram of regular qe schemes

$$X_1 = V_0 \xrightarrow{-\varphi_1} V_1 \xrightarrow{-\varphi_2} \dots \xrightarrow{-\varphi_{l-1}} V_{l-1} \xrightarrow{-\varphi_l} V_l = X_2$$

along with regular schemes  $Z_i$  for  $i = 1, \dots, l$  and ideal sheaves  $J_i$  for  $i = 1, \dots, (l-1)$  satisfying the following conditions:

- (1)  $\phi = \varphi_l \circ \varphi_{l-1} \circ \dots \circ \varphi_2 \circ \varphi_1$ .
- (2) The maps  $V_i \dashrightarrow X_2$  are morphisms; these maps as well as  $\varphi_i$  induce isomorphisms on  $U$ .
- (3) For every  $i = 1, \dots, l$  either  $\varphi_i: V_{i-1} \dashrightarrow V_i$  or  $\varphi_i^{-1}: V_i \dashrightarrow V_{i-1}$  is a morphism given as the blowing up of  $Z_i$ , which is respectively a subscheme of  $V_i$  or  $V_{i-1}$  disjoint from  $U$ .
- (4) The inverse image  $D_{V_i} \subset V_i$  of  $D_2 \subset X_2$  is a normal crossings divisor and  $Z_i$  has normal crossings with  $D_{V_i}$ .
- (5) For every  $i = 1, \dots, (l-1)$ , the morphism  $V_i \rightarrow X_2$  is given as the blowing up of the corresponding coherent ideal sheaf  $J_i$  on  $X_2$ , which is the unit ideal on  $U$ .

To include  $V_0 \rightarrow X_2$ , we define  $J_0 = I$ . The ideals  $J_i$  are a convenient way to encode functoriality, especially when we later pass to other geometric categories.

These conditions are the same as (1)–(5) in [AKMW02, Theorem 0.3.1], except that here the centers of blowing up and ideal sheaves are specified. Condition (2) is formulated for convenience; it is a consequence of (3) and (5). Note that here, as in [AKMW02, Theorem 0.3.1], the centers are not assumed irreducible, in contrast with [AKMW02, Theorem 0.1.1]. With these condition, the most basic form of our main theorem is as follows:

Th: factor-simple

**Theorem 1.2.1** (Weak factorization). *Every birational blowing up  $\phi: X_1 \rightarrow X_2$  of a noetherian qe regular  $\mathbb{Q}$ -scheme has a weak factorization  $X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2$ .*

The adjective “weak” serves to indicate that blowings up and down may alternate arbitrarily among the maps  $\varphi_i$ , as opposed to a *strong factorization*, where one has a sequence of blowings up followed by a sequence of blowings down. We note that at present strong factorization is not known even for toric threefolds.

Theorem 1.2.1 generalizes [Wlo03, 0.0.1] and [AKMW02, Theorem 0.1.1], where the case of varieties is considered. But we wish to prove a more precise theorem.

**1.3. Functorial weak factorization.** The class of data  $(X_2, I, D_2)$ , namely morphisms  $\phi: X_1 \rightarrow X_2$  of noetherian qe regular schemes given as blowings up of ideals  $I$ , with divisor  $D_2$  as in Section 1.2, can be made into *the regular surjective category of blowings up*, denoted  $\text{Bl}_{\text{rs}}$ , by defining arrows as follows:

**Def:Bl**

**Definition 1.3.1.** An arrow from the blowing up  $\phi': X'_1 = \text{Bl}_{I'}(X'_2) \rightarrow X'_2$  to  $\phi: X_1 = \text{Bl}_I(X_2) \rightarrow X_2$  is a regular and surjective morphism  $g: X'_2 \rightarrow X_2$  such that  $g^*I = I'$  and  $g^{-1}D_2 = D'_2$ . In particular,  $g$  induces a canonical isomorphism  $X'_1 \rightarrow X_1 \times_{X_2} X'_2$  and  $D'_1$  is the preimage of  $D_1$  under  $X'_1 \rightarrow X_1$ .

Similarly, weak factorizations can be made into the regular surjective category of weak factorizations, denoted  $\text{Fact}_{\text{rs}}$ , by defining arrows as follows:

**Def:factcat**

**Definition 1.3.2.** A morphism in  $\text{Fact}_{\text{rs}}$  from a weak factorization

$$X'_1 = V'_0 \dashrightarrow V'_1 \dashrightarrow \dots \dashrightarrow V'_{l-1} \dashrightarrow V'_l = X'_2$$

of  $\phi': X'_1 \rightarrow X'_2$ , with centers  $Z'_i$  and ideals  $J'_i$  to a weak factorization

$$X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2$$

of  $\phi: X_1 \rightarrow X_2$ , with centers  $Z_i$  and ideals  $J_i$  consists of a regular surjective morphism  $g: X'_2 \rightarrow X_2$  such that  $g^*I = I'$ ,  $g^*J_i = J'_i$  inducing  $g_i: V'_i \rightarrow V_i$ , such that  $Z'_i = g_i^{-1}Z_i$  or  $g_i^{-1}Z_i$  as appropriate. In particular  $\varphi_i \circ g_{i-1} = g_i \circ \varphi_i$  and  $g_i^{-1}D_{V_i} = D_{V'_i}$ .

Note that given a factorization of  $\phi$ , any morphism from a factorization of  $\phi'$  is uniquely determined by  $g: X'_2 \rightarrow X_2$ .

If we wish to restrict to schemes in a given characteristic  $p$  we denote the categories  $\text{Bl}_{\text{rs}}(\text{char} = p)$  and  $\text{Fact}_{\text{rs}}(\text{char} = p)$  respectively. If we wish to restrict the dimension we write  $\text{Bl}_{\text{rs}}(\text{char} = p, \dim \leq d)$  and  $\text{Fact}_{\text{rs}}(\text{char} = p, \dim \leq d)$ .

There is an evident forgetful functor  $\text{Fact}_{\text{rs}} \rightarrow \text{Bl}_{\text{rs}}$  taking a weak factorization  $X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2$  to its composition  $\phi: X_1 \rightarrow X_2$ . The weak factorization theorem provides a section, when strong resolution of singularities holds:

**Th:main**

**Theorem 1.3.3.** (1) **FUNCTORIAL WEAK FACTORIZATION:** *There is a functor*

$$\text{Bl}_{\text{rs}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}(\text{char} = 0)$$

*assigning to a blowing up  $\phi: X_1 \rightarrow X_2$  in characteristic 0 a weak factorization*

$$X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2,$$

*so that the composite  $\text{Bl}_{\text{rs}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}(\text{char} = 0) \rightarrow \text{Bl}_{\text{rs}}(\text{char} = 0)$  is the identity.*

(2) **CONDITIONAL FACTORIZATION IN POSITIVE AND MIXED CHARACTERISTICS:** *If functorial embedded resolution of singularities applies in characteristic  $p$  (respectively, over  $\mathbb{Z}$ ) for schemes of dimension  $\leq d + 1$ , then there is a functor*

$$\text{Bl}_{\text{rs}}(\text{char} = p, \dim \leq d) \rightarrow \text{Fact}_{\text{rs}}(\text{char} = p, \dim \leq d)$$

*(respectively, a functor*

$$\text{Bl}_{\text{rs}}(\dim \leq d) \rightarrow \text{Fact}_{\text{rs}}(\dim \leq d)$$

which is a section of  $\text{Fact}_{\text{rs}}(\text{char} = p, \dim \leq d) \rightarrow \text{Bl}_{\text{rs}}(\text{char} = p, \dim \leq d)$  (respectively,  $\text{Fact}_{\text{rs}}(\dim \leq d) \rightarrow \text{Bl}_{\text{rs}}(\dim \leq d)$ ).

This generalizes a theorem for *varieties* in characteristic 0, [AKMW02, Theorem 0.3.1 and Remark (3) thereafter], [Wlo06, Theorem 1.1], [Wlo09, Theorem 0.0.1], where the factorization is only shown to be functorial for isomorphisms. The precise statements we need for part (2) are spelled out below as Hypothetical Statements 2.1.4 and 2.2.4.

**Remark 1.3.4** (Preservation of  $G$ -normality). In [BL05, Definition 3.1] Borisov and Libgober introduce  $G$ -normal divisors and in [BL05, Theorem 3.8] they show that this condition can be preserved in the algorithm of [AKMW02]. The same holds true here, using the same argument of [BL05, Theorem 3.8], by performing the sequence of blowings up associated to the barycentric subdivision on the schemes  $W_{i\pm}^{\text{res}}$  obtained in Section 5.2. Details are left to the interested reader.

**1.4. Applications of functoriality.** We need to justify the somewhat heavy functorial treatment. Of course functoriality may be useful if one wants to make sure the factorization is equivariant under group actions and separable field extensions; this has been of use already in the case of varieties. But it also serves as a tool to transport our factorization result to other geometric spaces.

Blowings up of regular objects is a concept which exists in categories other than schemes, for instance: Artin stacks,  $qe$  formal schemes, complex semianalytic germs (see Appendix B), Berkovich  $k$ -analytic spaces, rigid  $k$ -analytic spaces. For brevity we denote the full subcategory of  $qe$  *noetherian* objects in any of these categories by  $\mathfrak{Sp}$ . Functoriality, as well as the generality of  $qe$  schemes, is crucial in proving the following:

Th:factor-Sp

**Theorem 1.4.1** (Factorization in other categories). *Any blowing up  $X_1 \rightarrow X_2$  of either noetherian  $qe$  regular algebraic stacks, or regular objects of  $\mathfrak{Sp}$ , in characteristic 0 has a weak factorization  $X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{i-1} \dashrightarrow V_i = X_2$ . The same holds in positive and mixed characteristics (when relevant) if functorial embedded resolution of singularities for  $qe$  schemes applies in positive and mixed characteristics.*

See Theorem 6.1.3 for the case of stacks and Theorem 6.4.5 for other categories, where functoriality is also shown, in other words Theorem 1.3.3 applies in each of the categories  $\mathfrak{Sp}$ . In addition, the argument deducing Theorem 6.1.3 from Theorem 1.3.3 is a formal one based on functoriality, so the same argument can be used to extend Theorem 6.4.5 to stacks in the categories of formal schemes, Berkovich spaces, etc., once an appropriated theory of stacks is constructed, see for instance [Sim96, Noo05, Uli15, Yu14, PY14].

Sec:strong-functoriality

**1.5. The question of stronger functoriality.** It is natural to replace the category  $\text{Bl}_{\text{rs}}$  by the category  $\text{Bl}_r$  with the same objects but where arrows  $g : X'_2 \rightarrow X_2$  as in Definition 1.3.1 are not required to be surjective, only regular. In a similar way one can replace the category  $\text{Fact}_{\text{rs}}$  by a category  $\text{Fact}_r$ . As explained in [Tem12, §2.3.3] for resolution of singularities, removing the surjectivity assumption requires imposing an equivalence relation on factorizations, in which two factorizations which differ by a step which is the blowing up of the unit ideal are considered equivalent. It is conceivable that the analogue of Theorem 1.3.3 may hold for  $\text{Fact}_r \rightarrow \text{Bl}_r$ .

**1.6. Factorization of birational and bimeromorphic maps.** Our results for projective morphism imply results for birational and bimeromorphic maps. We start with the case of schemes. By a proper birational map  $f: X_1 \dashrightarrow X_2$  of reduced schemes we mean an isomorphism  $f_0: U_1 \rightarrow U_2$  of dense open subschemes such that the closure  $Y \subset X_1 \times X_2$  of the graph of  $f_0$  is proper over each  $X_i$ . Assume that  $X_i$  are regular. The factorization problem for the birational map  $f$  reduces to factorization of the proper morphisms  $Y^{\text{res}} \rightarrow X_i$ , where  $Y^{\text{res}}$  is a resolution of  $Y$ . Assume, now, that  $f: X_1 \rightarrow X_2$  is a proper birational morphism. By a blow up version of Chow's lemma (e.g., it follows from the flattening of Raynaud-Gruson) there exists a blowing up  $Y = \text{Bl}_I(X_1) \rightarrow X_2$  that factors through  $X_1$ . Then  $Y = \text{Bl}_{f^{-1}I}(X_1)$  and hence the resolution  $Y^{\text{res}}$ , which is the blowing up of  $Y$ , is also a blowing up of both  $X_i$ . Thus, factorization of  $f$  reduces to the factorization for blowings up, which was dealt with in Theorem 1.3.3.

Now, assume that  $\mathfrak{Sp}$  is any geometric category. The definition of a proper bimeromorphic map  $f: X_1 \rightarrow X_2$  is similar to the definition of a proper birational map with two addenda: in the case of stacks we require that the morphisms  $Y \rightarrow X_i$  are representable, and in the case of analytic spaces or formal schemes we require that  $U$  is open  $Y$  (in particular,  $Y \rightarrow X_i$  are bimeromorphic). Then the general factorization problem immediately reduces to the case when  $f$  is a proper morphism. Furthermore, if objects of  $\mathfrak{Sp}$  are compact and if Chow's lemma holds in  $\mathfrak{Sp}$  then the problem reduces further to the case when  $f$  is a blowing up. For complex analytic spaces, Chow's lemma was proved by Hironaka in [Hir75, Corollary 2]. It extends immediately to the complex analytic germs we consider in this paper, and these are indeed compact. Most probably, it also holds in all other categories  $\mathfrak{Sp}$  we mentioned, but this does not seem to be worked out so far.

## 2. QE SCHEMES AND FUNCTORIALITY

Sec:qe-resolution

### 2.1. Qe schemes and their resolution of singularities.

2.1.1. *Qe schemes.* The class of quasi-excellent schemes was introduced by Grothendieck as the natural class where problems related to resolution of singularities behave well. The name "quasi-excellent" is perhaps not very elegant (it was not introduced by Grothendieck), and we feel it harmless to refer to them as *qe schemes*.

First recall that regular morphisms are a generalization of smooth morphisms in situations of morphisms which are not necessarily of finite type. Following [Gro67, IV<sub>2</sub>, 6.8.1] a morphism of schemes  $f: Y \rightarrow X$  is said to be *regular* if

- the morphism  $f$  is flat and
- all geometric fibers of  $f: Y \rightarrow X$  are regular.

A locally noetherian scheme  $X$  is a *qe scheme* if the following two conditions hold:

- for any scheme  $Y$  of finite type over  $X$ , the regular locus  $Y_{\text{reg}}$  is open; and
- For any point  $x \in X$ , the completion morphism  $\text{Spec } \hat{\mathcal{O}}_{X,x} \rightarrow \text{Spec } \mathcal{O}_{X,x}$  is regular.

It is a known, but nontrivial fact, that a scheme  $Y$  of finite type over a qe scheme is also a qe scheme, see, for example, [Mat80, 34.A]. A ring  $A$  is a qe ring if  $\text{Spec } A$  is a qe scheme.

Sec:resolution

2.1.2. *Resolution of singularities of qe schemes: characteristic 0.* Since [Gro67, IV<sub>2</sub>, 7.9.6], it is universally hoped that every qe scheme admits a good resolution of singularities. It is now known, see [Tem12, Theorem 1.2.1], that all qe *noetherian*  $\mathbb{Q}$ -schemes admit resolution of singularities, and this resolution is given by a sequence of blowings up with nonsingular centers, which is functorial in regular morphisms of qe schemes. Note that here we only require the morphisms to be regular, not necessarily surjective. Furthermore, functoriality of this resolution implies that one also gets a functorial way to resolve an arbitrary qe scheme (locally noetherian but not necessarily noetherian) by a morphism  $f_X: X' \rightarrow X$ . In general there is no canonical factorization of  $f_X$  into a composition of blowings up with regular centers, but  $f_X$  can be realized as an infinite composition whose restrictions onto noetherian open subschemes of  $X$  are finite, see [Tem12, Theorem 5.3.2].

2.1.3. *Positive and mixed characteristics hypothesis.* In Theorem 1.3.3 (3), the precise hypothetical statement we need about resolutions is the following analogue of [Tem12, Theorem 1.2.1]:

Hyp:resolution

**Hypothetical Statement 2.1.4.** (1) **FUNCTORIAL RESOLUTION:** *Each noetherian qe  $\mathbb{F}_p$  scheme (respectively,  $\mathbb{Z}$ -scheme)  $X$  of dimension  $\leq d + 1$  admits an ideal sheaf  $J_X$  whose blowing up  $\tilde{X} \rightarrow X$  is a resolution of singularities, in a manner functorial for regular surjective morphisms  $X' \rightarrow X$ , so that  $\tilde{X}' = X' \times_X \tilde{X}$ .*

(2) **EQUIVARIANCE:** *Moreover, if  $G$  is a smooth group scheme of arbitrary dimension,  $a: G \times X \rightarrow X$  an action and  $p_X: G \times X \rightarrow X$  the projection, then  $J_X$  is  $G$ -equivariant:  $a^{-1}J_X = p_X^{-1}J_X$ .*

In mixed characteristics we will also need:

(3) **FUNCTORIALITY OF TOROIDAL CHARTS:** *For any monoidal chart  $j: \hat{X}_p \hookrightarrow Y := \text{Spec } C[[M]]$  in the sense of [AT15b, Lemma 2.3.9 and Theorem 2.3.11], we have  $j^{-1}J_Y = J_{\hat{X}_p}$  with  $J_Y$  a monomial ideal.*

The statement here, with a single resolving ideal sheaf  $J_X$ , differs from [Tem12, Theorem 1.2.1], which gives a sequence of blowings up. We will see below that a sequence of blowings up can be transformed, functorially under *surjective* regular morphisms, into a single ideal sheaf.

We note that the equivariance statement (2) in dimension  $(d + 1)$  follows from statement (1) in dimension  $(d + 1) + \dim G$ , but here we wish to only make assumptions up to dimension  $d + 1$ . It is conceivable that a version of (2) sufficient for our needs follows from (1) by taking slices, but we will not pursue this question.

Let us say that a scheme  $X$  is *locally monoidal* if locally it admits a logarithmic structure making it to a logarithmically regular scheme. It is expected that there should exist a canonical resolution of such schemes of combinatorial nature, which is, in particular, independent of the characteristics. Our Statement (3) asserts such independence in mixed characteristics; in pure characteristics it is a consequence of equivariance. It is analogous to Hypothetical Statement 2.2.4(3) below. Similarly to Hypothetical Statement 2.2.4, proving Statements (1)–(3) for locally monoidal schemes is expected to be easier than the general case. For example, it is proved in [IT14, Theorem 3.4.9] for logarithmically regular schemes (with a single logarithmic structure), but known functoriality [IT14, Theorem 3.4.15] is not enough to extend

it to locally monoidal schemes. In addition, very recently Buonerba resolved certain locally monoidal varieties in [Buo15].

Sec:principalization

**2.2. Principalization.** In addition to resolution of singularities, we will need a version of functorial principalization of coherent ideal sheaves on qe regular schemes *with boundaries*.

We consider the category of triples  $(X, I, D)$  where  $X$  is a noetherian regular qe scheme,  $I$  is a coherent ideal sheaf,  $D \subset X$  a normal crossings divisor, and arrows are regular morphisms  $f: X_1 \rightarrow X_2$  such that  $f^{-1}I_2 = I_1$  and  $f^{-1}D_2 = D_1$ . Again we only require the morphisms to be regular, not necessarily surjective. A principalization of  $I$  is a sequence of blowings up of regular centers  $\phi_X: \tilde{X} \rightarrow X$ , which are the identity on the locus where  $I$  is the unit ideal, such that  $\phi_X^{-1}I$  is generated locally at every point by a single monomial in regular parameters. The principalization is compatible with  $D$  if at each stage  $X_i$  of the sequence forming  $\phi_X: \tilde{X} \rightarrow X$ , the inverse image  $D_i \subset X_i$  of  $D$  is a normal crossings divisor, and the new center of blowing up has normal crossings with  $D_i$ . Principalizations compatible with  $D$  form a category again, and functorial principalization provides a functor from triples  $(X, I, D)$  to principalizations  $\phi_X: \tilde{X} \rightarrow X$ . As we do not require the morphism  $f$  to be surjective, we have to use the equivalence relation mentioned in Section 1.5. However, we will only apply the result in the context of surjective morphism, so this equivalence will not figure in any of our applications.

Functorial principalization of ideal sheaves for *varieties* over a field of characteristic 0 is known, see [BM97, Sections 11,13], [Kol07, Theorem 3.26]; it is in fact compatible with an a-priori given normal crossings divisor  $D \subset X$ . The second author is in the process of writing a general functorial principalization of ideal sheaves on noetherian regular qe  $\mathbb{Q}$ -schemes with the methods of [Tem09]. Until that becomes available we prefer not to use it. In the present paper we only need principalization for ideals which are locally given as monomial ideals, in particular the ideal is locally the pullback of a toric ideal through a morphism to a toric variety. These belong to a class for which functorial principalization can be obtained using the case of varieties and methods of [IT14, Theorem 2.4.1, p. 95], as follows.

princhyp

**Hypothetical Statement 2.2.1.** *Principalization exists for qe schemes.*<sup>1</sup>

←1

A triple  $(X, I, D)$  is said to be  $\mathbb{Q}$ -*absolute* if there exists an open covering  $\coprod U_\alpha \rightarrow X$ , regular  $\mathbb{Q}$ -varieties  $Z_\alpha$ , regular morphisms  $f_\alpha: U_\alpha \rightarrow Z_\alpha$ , ideal sheaves  $I_\alpha$  on  $Z_\alpha$  and divisors  $D_\alpha \subset Z_\alpha$  such that  $f_\alpha^{-1}I_\alpha = I|_{U_\alpha}$  and  $f_\alpha^{-1}D_\alpha = D|_{U_\alpha}$ . The collection of  $\mathbb{Q}$ -absolute triples forms a full subcategory of the category of triples. Functorial principalization of  $\mathbb{Q}$ -absolute triples  $(X, I, D)$  compatible with  $D$  is a functor from this subcategory to principalizations of the corresponding ideals.

The statement we need is the following:

absolute-principalization

**Proposition 2.2.2.** *There exists a functorial principalization  $\phi_X: \tilde{X} \rightarrow X$  compatible with  $D$  of  $\mathbb{Q}$ -absolute triples  $(X, I, D)$ .*

*Proof.* We may replace  $\coprod U_\alpha$  by a finite covering, since  $X$  is noetherian. We write  $U_{\alpha\beta} = U_\alpha \times_X U_\beta$ . Now, we will use the ideas from the proof of [IT14, Theorem 2.4.3].

<sup>1</sup>(Michael) Will add details later (dimensions, etc). Currently put this for referencing.

First we construct a principalization. For this it suffices to construct a principalization of  $\coprod(U_\alpha, I|_{U_\alpha}, D|_{U_\alpha})$  whose two pullbacks to the fiber product  $W := \coprod U_{\alpha\beta}$  coincide. The triple  $(Z, I_Z, D_Z) := \coprod(Z_\alpha, I_\alpha, D_\alpha)$  has a principalization compatible with  $D_\alpha$  coming from the principalization functor for  $\mathbb{Q}$ -varieties. This pulls back to a principalization of  $\coprod(U_\alpha, I|_{U_\alpha}, D|_{U_\alpha})$  and we need to show that the two pullbacks to  $W$  coincide. We have two regular morphisms  $f, g: W \rightarrow Z$ . By Popescu's theorem (see [Pop86] or [Spi99]),  $f$  is the limit of smooth morphisms  $f_\gamma: W_\gamma \rightarrow Z$ . By [Gro67, IV<sub>3</sub>, Proposition 8.13.1],  $g$  factors through a morphism  $g_\gamma: W_\gamma \rightarrow Z$  for a large enough  $\gamma$  and then [IT14, Proposition 2.4.3] implies that replacing  $W_\gamma$  by a neighborhood of the image of  $W$  we can achieve that  $g_\gamma$  is also smooth. Since the two pullbacks of  $I_Z$  and  $D_Z$  to  $W$  coincide, there is some  $\gamma$  such that the two pullbacks of  $I_Z$  and  $D_Z$  to  $W_\gamma$  coincide. It follows by functoriality of principalization for varieties that the two principalizations on  $W_\gamma$  coincide, and therefore they coincide on  $W$ , as required.

We now demonstrate that this principalization is functorial. Consider a regular surjective morphism  $f: (X_1, I_1, D_1) \rightarrow (X_2, I_2, D_2)$  with coverings  $\coprod U_{1\alpha}$  and  $\coprod U_{2\beta}$  and  $\mathbb{Q}$  varieties  $Z_{1\alpha}$  and  $Z_{2\beta}$ . Then composing  $U_{2\beta} \rightarrow Z_{2\beta}$  with  $f$  we get another covering  $\coprod f^{-1}U_{2\beta}$  with regular maps to  $Z_{2\beta}$ , so it is enough to show that the resulting principalizations on  $X_1$  coincide. We now write  $W = \coprod U_{1\alpha} \times_{X_1} f^{-1}U_{2\beta}$ , which maps to  $Z_1 = \coprod Z_{1\alpha}$  and  $Z_2 = \coprod Z_{2\beta}$ . By the same argument as earlier we have that  $W \rightarrow Z_1 \times Z_2$  is the limit of a family  $W_\gamma \rightarrow Z_1 \times Z_2$ , where the two maps  $W_\gamma \rightarrow Z_i$  are smooth. As above we conclude that the ideals and divisors coincide on some  $W_\gamma$  and the two principalizations coincide on  $W$  and therefore on  $X_1$ . ♣

**2.2.3. Positive and mixed characteristics hypothesis.** In Theorem 1.3.3 (3), the precise hypothetical setup and statement we need about principalization is the following. A triple  $(X, I, D)$  with  $X$  regular,  $I$  an ideal sheaf and  $D$  a normal crossings divisor is said to be *locally monoidal* if there is an open covering  $\coprod U_\alpha \rightarrow X$ , logarithmically regular structures  $(U_\alpha, M_\alpha)$  such that  $D$  is part of the toroidal divisor, and monoid ideals  $I_\alpha \subset M_\alpha$  such that  $I_{U_\alpha}$  is generated by the image of  $I_\alpha$  under  $M_\alpha \rightarrow \mathcal{O}_{U_\alpha}$ .

Hyp:principalization

**Hypothetical Statement 2.2.4.** (1) *Each locally monoidal  $\mathbb{F}_p$ -triple (respectively,  $\mathbb{Z}$ -triple)  $(X, I, D)$  of dimension  $\leq d$  admits a principalization  $\tilde{X} \rightarrow X$  compatible with  $D$ , in a manner functorial for regular morphisms  $X' \rightarrow X$ .*

(2) *Moreover, if  $G$  is a smooth group scheme of arbitrary dimension,  $a: G \times X \rightarrow X$  an action and  $p_X: G \times X \rightarrow X$  the projection, and if  $I$  and  $D$  are equivariant:  $a^{-1}I = p_X^{-1}I$  and  $a^{-1}D = p_X^{-1}D$ , then  $\tilde{X} \rightarrow X$  is  $G$ -equivariant as well.*

Again in mixed characteristics we also need:

(3) **FUNCTORIALITY OF TOROIDAL CHARTS:** *For any monoidal chart  $j: \hat{X}_p \hookrightarrow Y := \text{Spec } C[[M]]$  in the sense of [AT15b, Lemma 2.3.9 and Theorem 2.3.11], we have  $\tilde{\hat{X}}_p = \hat{X}_p \times_Y \tilde{Y}$ , with  $\tilde{Y} \rightarrow Y$  a toric morphism.*

We remark that the results of [IT14, Section 3.1.14] suggest that this statement may be within reach: in that paper the local non-functorial problem is solved, and the problem reduces to making the process functorial even if one changes the logarithmic structure  $M_\alpha$  on  $U_\alpha$ .



ec:functorial-projective

**2.3. Functorial constructions.** In our method, it will be important to describe certain morphisms we will obtain as blowing up of a concrete ideal or an explicitly described projective morphism, since further constructions will depend on this data. Moreover, this should be done functorially with respect to surjective regular morphisms. In the current section we develop a few basic functorial constructions of this type.

There are few ways to describe a projective morphism: using Proj, using ample sheaves, or using projective fibrations, but each approach involves choices. Neither description is “more natural” than the others, and we will have to switch between them. Similarly to [Gro67, II] we choose the language of projective fibrations to be the basic one and we will show how other descriptions are canonically reduced to projective fibrations.

**2.3.1. Projective fibrations.** Let  $X$  be a scheme. For a coherent  $\mathcal{O}_X$ -module  $E$  consider the projective fibration  $\mathbb{P}(E) = \mathbb{P}_X(E) := \text{Proj}_X \text{Sym}^\bullet(E)$  associated with  $E$ . It has a canonical twisting sheaf  $\mathcal{O}_{\mathbb{P}(E)}(1)$ , and  $E \rightarrow \pi_* \mathcal{O}(1)$  is an isomorphism. This construction is functorial for all morphisms: if  $\phi: X' \rightarrow X$  is any morphism and  $E' = \phi^* E$  then  $\mathbb{P}_{X'}(E') = X' \times_X \mathbb{P}_X(E)$ , and  $\mathcal{O}_{\mathbb{P}(E')}(1)$  is the pullback of  $\mathcal{O}_{\mathbb{P}(E)}(1)$ .

Sec:projective-morphisms

**2.3.2. Projective morphisms.** By the usual definition [Gro67, II, 5.5.2], a morphism  $f: Y \rightarrow X$  is projective if it factors through a closed immersion  $i: Y \hookrightarrow \mathbb{P}_X(E)$  for a coherent  $\mathcal{O}_X$ -module  $E$ . In this paper, we will use the convention that by saying “ $f$  is projective” we fix  $E$  and  $i$ . In particular,  $Y$  acquires a canonical relatively very ample sheaf  $\mathcal{O}_Y(1) = \mathcal{O}_{\mathbb{P}(E)}(1)|_Y$ .

**2.3.3. Relation to Proj.** For a projective morphism  $f: Y \rightarrow X$  we also obtain a canonical description of  $Y$  as a Proj. Namely, if  $I_Y \subseteq \mathcal{O}_{\mathbb{P}(E)}$  denotes the ideal defining  $Y$  then  $Y = \text{Proj}_X A$ , where  $A^\bullet = \text{Sym}^\bullet(E)/I_B$  is a quasi-coherent  $\mathcal{O}_X$ -algebra with coherent graded components, generated over  $A^0 = \mathcal{O}_X$  by its degree-1 component  $A^1$ . Again this structure is functorial for all morphisms: if  $\phi: X' \rightarrow X$  is any morphism and  $A' = \phi^* A$  then  $\text{Proj}_{X'} A' = X' \times_X \text{Proj}_X A$ .

Conversely, if a graded  $\mathcal{O}_X$ -algebra  $A^\bullet$  has coherent components and is generated over  $A^0 = \mathcal{O}_X$  by  $A^1$  then  $\text{Sym}^\bullet(A^1) \twoheadrightarrow A^\bullet$  and we obtain a closed immersion  $i: \text{Proj}_X A \hookrightarrow \mathbb{P}_X(A^1)$ . Thus,  $Y = \text{Proj}_X A$  is projective over  $X$ , and the associated graded quasi-coherent algebra is  $A$  itself. This construction is also functorial for all morphisms.

**Remark 2.3.4.** We note that the construction of a projective morphism from Proj is right inverse to the construction of Proj from a projective morphism, but they are not inverse: going from a projective morphisms to Proj and back to a projective morphism one usually changes the projective fibration.

**Remark 2.3.5.** In this paper we use superscripts to denote degrees of homogeneous components of a graded object, as in  $A^i \subset A^\bullet$ . When considering weights of a given  $\mathbb{G}_m$ -action we will use subscripts. We hope this will not cause confusion.

**2.3.6. General Proj.** Consider now a general quasi-coherent graded  $\mathcal{O}_X$ -algebra with coherent graded components, which is only assumed to be generated over  $A_0 = \mathcal{O}_X$  in finitely many degrees. Writing  $A^{M^\bullet} = \bigoplus_j A^{Mj}$  for a positive integer

$M$ , we have a canonical isomorphism  $Y = \text{Proj}_X A^\bullet \simeq \text{Proj}_X A^{M^\bullet}$ . For a suitable  $M$  the algebra  $A^{M^\bullet}$  is generated in degree 1 by  $A^M$ . If we take the minimal  $M_0$  such that  $A^{M^\bullet}$  is generated in degree 1, then  $L$  is not functorial for all morphisms. Rather it is functorial for all *flat surjective* morphisms  $X' \rightarrow X$ : if  $A^{M^\bullet}$  is generated in degree 1 then  $(A')^{M^\bullet}$  is generated in degree 1, and the opposite is true whenever  $X' \rightarrow X$  is flat surjective; this follows since surjectivity of  $((A')^1)^{\otimes n} \rightarrow (A')^n$  implies surjectivity of  $(A^1)^{\otimes n} \rightarrow A^n$  by flat decent. Combining this construction with the previous one we obtain an interpretation of  $Y \rightarrow X$  as a projective morphism, and this construction is functorial for all flat surjective morphisms.

**Remark 2.3.7.** This construction applies to the following situation: assume  $f: Y \rightarrow X$  is a proper morphism of noetherian schemes and  $L$  is an  $f$ -ample sheaf. Then  $A^\bullet = \mathcal{O}_X \oplus \bigoplus_{k=1}^{\infty} f_*(L^k)$  is generated in finitely many degrees and  $Y = \text{Proj}_X A$ . Therefore,  $L$  gives rise to an interpretation of  $f$  as a projective morphism functorially for all surjective flat morphisms.

Sec:blowup-projective

2.3.8. *Blowings up.* An important variant is that of blowings up. Consider a coherent ideal sheaf  $I$  on  $X$ . The Rees algebra  $R_X(I) = \bigoplus_{k=0}^{\infty} I^k$  is generated in degree 1, and we define  $Bl_I(X) = \text{Proj}_X R_X(I)$ . In particular,  $Bl_I(X)$  is projective over  $X$  with the closed immersion  $Bl_I(X) \hookrightarrow \mathbb{P}_X(I)$ . If  $\phi: X' \rightarrow X$  is a morphism, then  $I^k \mathcal{O}_{X'} = (I \mathcal{O}_{X'})^k = (I')^k$  and  $\phi^*(I^k) \rightarrow I^k \mathcal{O}_{X'}$  is surjective, giving a canonical morphism  $\phi': Bl_{I'}(X') \rightarrow Bl_I(X)$  over  $\phi$ . Clearly  $(\phi')^* L = L'$ . So a blowing up is functorially projective. If moreover  $X' \rightarrow X$  is flat, then  $Bl_{I'}(X') = X' \times_X Bl_I(X)$ .

We will need an opposite construction, using a variant of [Har77, Theorem II.7.17] for regular schemes. Assume  $X$  is regular and  $f: Y \rightarrow X$  is a proper birational morphism with an ample sheaf  $L$  (e.g., if  $Y \rightarrow X$  is projective we can take  $L = \mathcal{O}_Y(1)$ ). Then after replacing  $L$  by a positive power which is functorial for flat surjective morphisms, we have that  $Y = \text{Proj}_X A^\bullet$ , where  $A^\bullet$  is generated over  $A_0 = \mathcal{O}_X$  by its degree-1 component, and  $A^k = f_* L^k$ .

Locally on  $X$ , write  $L^k$  as a fractional ideal on  $Y$ , giving it as a fractional ideal  $F_{L,k}$  on  $X$  since  $Y \rightarrow X$  is birational. Since  $A^\bullet$  is generated in degree 1, we have that  $F_{L,k} = F_{L,1}^k$  (see [Har77, Theorem II.7.17 Step 5]). Since  $X$  is factorial, there is a unique expression  $F_{L,1} = MI$ , where  $M$  is an invertible fractional ideal and  $I$  is an *ideal sheaf* without invertible factors. Explicitly,  $F_{L,1}^*$  is invertible, so we can write  $I = F_{L,1}^* F_{L,1}$  and  $M = F_{L,1}^{**}$ . It follows that  $F_{L,k} = M^k I^k$ . Note that while the construction is local on  $X$  and depends on an embedding of  $L$  in the fraction field, the ideal sheaf  $I$  glues canonically. Locally on  $X$  we have a canonical isomorphism  $Y \simeq Bl_I(X)$ , which evidently glues canonically. We have obtained that a projective birational morphism  $f: Y \rightarrow X$  with  $X$  regular is a blowing up, functorially for flat surjective morphisms  $X' \rightarrow X$  of regular schemes.

For future reference we record the following well known result that follows from the universal property of blowings up.

factorblowlem

**Lemma 2.3.9.** *If  $X$  is an integral scheme and a blowing up  $Y = Bl_I(X) \rightarrow X$  factors through a proper birational morphism  $Z \rightarrow X$  then  $Y = Bl_{I \mathcal{O}_Z}(Z)$ .*

Sec:sequence-projective

2.3.10. *Sequences of projective morphisms.* Now assume  $Z \xrightarrow{g} Y \xrightarrow{f} X$  is a sequence of projective morphisms of noetherian schemes, say  $Z \hookrightarrow \mathbb{P}_Y(F)$  and  $Y \hookrightarrow \mathbb{P}_X(E)$  for a coherent  $\mathcal{O}_Y$ -module  $F$  and a coherent  $\mathcal{O}_X$ -module  $E$ . For a large enough

$n$  the map  $f^*f_*(F(n)) \xrightarrow{\alpha} F(n)$  is surjective, hence  $\mathbb{P}_Y(F) = \mathbb{P}_Y(F(n))$  embeds into  $\mathbb{P}_X(E \otimes f_*F(n))$  and we obtain a closed immersion  $Z \hookrightarrow \mathbb{P}_X(E \otimes f_*F(n))$ . Choosing the minimal  $n$  such that  $\alpha$  is surjective we obtain a construction that realizes composition of projective morphisms as a projective morphism functorially for flat surjective morphisms  $X' \rightarrow X$ .

If  $X$  is regular we can combine this with the previous statements, so if  $Y_m \rightarrow \dots \rightarrow Y_1 \rightarrow X$  is a sequence of blowings up we have that  $Y_m \rightarrow X$  is a blowing up of an ideal sheaf, functorially for flat and surjective morphisms of regular schemes.

**Remark 2.3.11.** We will not use this, but blowings up can also be composed in terms of ideals. One can show that if  $X$  is normal then the composition of  $Y = Bl_I(X) \xrightarrow{f} X$  and  $Bl_J(Y) \rightarrow Y$  is of the form  $Bl_{f_*(f^{-1}(I^n)J)}(X) \rightarrow X$  for a large enough  $n$ .

### 3. FUNCTORIAL TOROIDAL FACTORIZATION

**3.1. Statement.** We follow the treatment of toroidal schemes in [AT15b, Section 2.3], in particular they carry logarithm structures in the Zariski topology. A toroidal ideal  $I$  on a toroidal scheme  $X$  with logarithmic structure  $M$  is the ideal generated by the image of a monomial ideal in  $M$  through  $M \rightarrow \mathcal{O}_X$ . We define a category  $\text{TorBl}_{\text{rs}}$  of toroidal blowings up, similar to  $\text{Bl}_{\text{rs}}$ :

- (1) An object is a birational transformation  $X_1 \rightarrow X_2$  where  $X_1, X_2$  are toroidal and regular, and  $X_1 \rightarrow X_2$  is given as the normalized blowing up of a toroidal ideal  $I \subset \mathcal{O}_{X_2}$ .
- (2) An arrow from  $X'_1 \rightarrow X'_2$  to  $X_1 \rightarrow X_2$  consists of a regular surjective morphism  $g: X'_2 \rightarrow X_2$ , such that  $U_{X_2} = g^{-1}U_{X'_2}$  and  $I' = I\mathcal{O}_{X'_2}$ .

We similarly define a *toroidal weak factorization*  $X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2$  of a toroidal blowing up  $X_1 \rightarrow X_2$ , where the schemes  $V_i$ , ideals  $J_i$  and centers  $Z_i$  are toroidal. These form *the regular surjective category*  $\text{TorFact}_{\text{rs}}$  of *toroidal weak factorizations* in a manner similar to the above.

p:toroidal-factorization

**Proposition 3.1.1.** *Let  $X_1 \rightarrow X_2$  be a toroidal morphism of toroidal schemes obtained by normalized blowing up a toroidal ideal. Then there is a toroidal weak factorization  $X_1 = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X_2$  in a functorial manner: there is a section  $\text{TorBl}_{\text{rs}} \rightarrow \text{TorFact}_{\text{rs}}$  of the forgetful functor  $\text{TorFact}_{\text{rs}} \rightarrow \text{Bl}_{\text{rs}}$ .*

**Remark 3.1.2.** Jarosław Włodarczyk informed us that one can prove a stronger result: a factorization procedure which is functorial for all regular strict morphisms  $g: X'_2 \rightarrow X_2$ , not required to be surjective. His proposed argument involves subtle modifications at the heart of the algorithm in [Wł09, Sections 4 and 5]. The proof we provide at the end of this section shows that *any* procedure for toric factorization gives rise to a functorial procedure.

**3.2. Cone complexes.** Before proving Proposition 3.1.1 we need to discuss a generalization of the polyhedral cone complexes with integral structure of [KKMSD73] which was introduced in [ACP12, 2.5] to accommodate any toroidal embedding in the sense of [KKMSD73], allowing for self intersections and monodromy. In this paper we only assign polyhedral cone complexes to Zariski toroidal schemes, without self intersections or monodromy, but the generalized polyhedral cone complexes are used as a combinatorial tool to achieve functoriality.

Fix a toroidal scheme  $X$ . Recall that the polyhedral complex of [KKMSD73] or the equivalent Kato fan of [Kat94] assigns a polyhedral cone  $\overline{\sigma}_Z$  with integral structure to each toroidal stratum  $Z \subset X$ ; each inclusion  $Z' \hookrightarrow Z \subset X$  gives rise to a linear map  $\nu : \sigma_Z \rightarrow \sigma_{Z'}$ , which identifies  $\sigma_Z$  as a face of  $\sigma_{Z'}$  in such a way that the integral structure on  $\sigma_Z$  is the restriction of the integral structure of  $\sigma_{Z'}$ : this is called a *face map*. We define  $\Sigma(X) = \varinjlim(\{\sigma_Z\}, \{\nu\})$  - it is similar to the fan of a toric variety, but is not embedded in a space  $N_{\mathbb{R}}$  and the intersection of two cones may be the union of faces rather than just one face.

A map of polyhedral cone complexes  $\varinjlim(\{\sigma'_i\}, \{\nu'_k\}) \rightarrow \varinjlim(\{\sigma_j\}, \{\nu_l\})$  is defined to be a collection of cone maps  $\sigma'_i \rightarrow \sigma_{j(i)}$  compatible with the face maps  $\nu'_k$  and  $\nu_k$ . A toroidal map  $X' \rightarrow X$  gives rise to a map of cone complexes; here are a few well known relationships:

- (1) A proper birational toroidal morphism gives rise to a subdivision, and there is an equivalence of categories between proper toroidal birational morphisms and subdivisions. Blowings up of ideals correspond to subdivisions determined by piecewise linear continuous integral functions which are convex on each cone; following [KKMSD73] we call these *projective subdivisions* (in the combinatorial literature they are *coherent subdivisions*).
- (2) A regular morphism  $g : X'_2 \rightarrow X_2$  such that  $U_{X_2} = g^{-1}U_{X'_2}$  gives rise to a map of complexes  $\Sigma(g) : \Sigma(X') \rightarrow \Sigma(X)$  where all the maps  $\sigma'_i \rightarrow \sigma_{j(i)}$  are face maps - this is called a *face map of complexes*.
- (3) If the map  $g : X'_2 \rightarrow X_2$  is also surjective then  $\Sigma(g)$  is surjective.
- (4) The scheme  $X$  is regular if and only if all the cones  $\sigma_i \subset \Sigma(X)$  are nonsingular in the usual toric sense.
- (5) If  $X$  is regular then the closure of a stratum is always regular (this would fail if we allowed self intersections); we call such subschemes *toroidal centers*.
- (6) The blowing up  $X' \rightarrow X$  of an irreducible toroidal center  $\overline{Z}$  on a *regular*  $X$  corresponds to the star subdivision  $\Sigma' \rightarrow \Sigma(X)$  at the barycenter of  $\sigma_Z$ . The blowing up  $X' \rightarrow X$  of any regular toroidal subscheme  $W$  corresponds to the simultaneous star subdivision  $\Sigma' \rightarrow \Sigma(X)$  at the barycenters of all the cones corresponding to the connected components of  $W$ .

Thus proposition 3.1.1 would follow if the projective subdivision  $\Sigma(X_1) \rightarrow \Sigma(X_2)$  can be factored as a composition of such simultaneous star subdivisions and their inverses, in such a way that the intermediate steps are projective subdivisions of  $\Sigma(X_2)$ , in a functorial manner with respect to surjective face maps. This will be our Lemma 3.5.1 below.

Morelli's  $\pi$ -desingularization lemma of fan cobordisms [Wlo03, Lemma 10.4.3] gives a non-functorial result in the case of fans; this was generalized in [AMR99] to polyhedral cone complexes. In [AKMW02] it is made functorial under *automorphisms*, which is not sufficient for our purposes here.

Consider the category whose objects are projective subdivisions  $\Sigma_1 \rightarrow \Sigma_2$  of nonsingular cone complexes given by a fixed piecewise linear continuous integral function  $f : \Sigma_2 \rightarrow \mathbb{R}$  convex on each cone and arrows  $(\Sigma'_2, f') \rightarrow (\Sigma_2, f)$  induced by surjective face maps  $h : \Sigma'_2 \rightarrow \Sigma_2$  with  $f' = f \circ h$ . Functoriality would be easily achieved if the connected component of any object  $\Sigma_1 \rightarrow \Sigma_2$  in this category had a final object, as we show below in Lemma 3.5.1. Indeed, this would mean that applying Morelli's lemma to the final object would induce a factorization for

the whole component, giving the result. Unfortunately final objects usually do not exist in the category of cone complexes. Our next goal is to enlarge this category so that final objects do exist, see Lemma 3.3.1 below.

**3.3. Generalized cone complexes and existence of final objects.** A generalized cone complex is given by any finite diagram  $(\{\sigma_j\}, \{\nu_l\})$  of cones and face maps. We allow for more than one face map  $\sigma_j \rightarrow \sigma_l$ , including non-trivial self-face maps  $\sigma_j \rightarrow \sigma_j$ . We think of a generalized cone complex  $\Sigma$  as a structure imposed on the topological space  $\underline{\Sigma} = \varinjlim(\{\sigma_j\}, \{\nu_l\})$ . Thus an arrow of generalized cone complexes  $(\{\sigma'_i\}, \{\nu'_k\}) \rightarrow (\{\sigma_j\}, \{\nu_l\})$  is given by compatible cone maps as above; an arrow is a face map if it is given by compatible face maps; and an arrow is declared to be an isomorphism if it is a face map inducing a bijection of sets  $\varinjlim(\{\sigma'_i\}, \{\nu'_k\}) \rightarrow \varinjlim(\{\sigma_j\}, \{\nu_l\})$ .

Cone complexes are a full subcategory of generalized cone complexes. They are distinguished by the property that, for any cones  $\tau, \sigma$  of  $\Sigma$  a face map  $\nu : \tau \rightarrow \sigma$  in  $\Sigma$  is unique if it exists. Thus proposition 3.1.1 would again follow if any projective subdivision  $\Sigma_1 \rightarrow \Sigma_2$  of *generalized* nonsingular cone complexes can be factored as a composition of simultaneous star subdivisions and their inverses, in a functorial manner with respect to surjective cone maps. The advantage of working with generalized cone complexes is the following:

**Lem:final**

**Lemma 3.3.1.** *The connected component of the projective subdivision  $\Sigma_1 \rightarrow \Sigma_2$  of generalized cone complexes in the category induced by surjective face maps  $\Sigma'_2 \rightarrow \Sigma_2$  has a final object.*

*Proof.* The projective subdivision  $\Sigma_1 \rightarrow \Sigma_2$  is induced by an implicit piecewise linear convex integral function  $f : \Sigma_2 \rightarrow \mathbb{R}$ . Write  $\Sigma_2 = (\{\sigma_j\}, \{\nu_l\})$ . Then  $\nu_l : \sigma_i \rightarrow \sigma_j$  has the property that  $f_{\sigma_i} = f_{\sigma_j} \circ \nu_l$ . Let  $\{\mu_k\}$  be the collection of all face maps  $\mu_k : \sigma_m \rightarrow \sigma_n$  with the property that  $f_{\sigma_m} = f_{\sigma_n} \circ \mu_k$ . Then  $\Delta := (\{\sigma_j\}, \{\mu_k\})$  is a generalized cone complex, the maps  $f_{\sigma_j}$  glue to give a piecewise linear integral function  $\tilde{f} : \Delta \rightarrow \mathbb{R}$ , and since  $\{\nu_l\} \subset \{\mu_k\}$  we have a map of diagrams  $g : \Sigma_2 \rightarrow \Delta$  such that  $f = \tilde{f} \circ g$ .

It is convenient to have another presentation of  $\Delta$ . Choose one representative  $\bar{\sigma}$  from each isomorphism class of cones in  $\Delta$ . Given two such representatives  $\bar{\tau}$  and  $\bar{\sigma}$ , consider all maps  $\bar{\nu}_l : \bar{\tau} \rightarrow \bar{\sigma}$  in  $\Delta$ . Clearly  $\bar{\Delta} = (\{\bar{\sigma}\}, \{\bar{\nu}_l\})$  maps as a subdiagram to  $\Delta$ , and the map is an isomorphism since it is clearly a bijection on set theoretic limits.

We claim that  $(\Delta, \tilde{f})$  is a final object in the component of  $(\Sigma_2, f)$  in the category of generalized cone complexes with piecewise linear integral function. For this it suffices to show that if  $(\Sigma'_2, f')$  is an object and  $h : \Sigma_2 \rightarrow \Sigma'_2$  is a surjective face map such that  $f' \circ h = f$  then  $g = g'' \circ h$  where  $g'' : \Sigma'_2 \rightarrow \Delta$  is a morphism so that  $f' = \tilde{f} \circ g''$ .

First, if we apply the construction of  $\Delta$  to  $\Sigma'_2$  we get a map  $g' : \Sigma'_2 \rightarrow \Delta'$  which sits in a commutative diagram

$$\begin{array}{ccc}
 \Sigma_2 & \xrightarrow{g} & \Delta \\
 \downarrow h & \searrow f & \downarrow \tilde{h} \\
 & & \mathbb{R} \\
 \Sigma'_2 & \xrightarrow{g'} & \Delta' \\
 & \nearrow f' & \downarrow \tilde{h}' \\
 & & \mathbb{R}
 \end{array}$$

On the other hand  $\bar{\Delta} \simeq \Delta$  and  $\bar{\Delta}' \simeq \Delta'$ , and the map  $\bar{\Delta} \rightarrow \bar{\Delta}'$  induced by  $\tilde{h}$  is an isomorphism of diagrams: since  $h$  is a surjective face map, any cone in  $\Sigma'_2$  is isomorphic to a cone of  $\Sigma_1$  via an isomorphism compatible with  $f$  and vice versa. So  $\tilde{h}$  gives a bijection between the isomorphism classes of cones, and the maps  $\bar{\nu}$  between cones are determined by the compatibility of the function  $\tilde{f} = \tilde{f}'$  on them. So  $\Delta \rightarrow \Delta'$  is an isomorphism, giving the requisite map of generalized complexes  $g'' = \tilde{h}^{-1} \circ g'$ . ♣

**3.4. Barycentric subdivisions and factorization for generalized cone complexes.** We proceed to extend the factorization of subdivisions of cone complexes to generalized cone complexes. We do it by a reduction step using barycentric subdivisions:

Lem:barycentric

- Lemma 3.4.1.** (1) ([ACP12, 2.5]) *The barycentric subdivision  $B(\Delta)$  of a generalized cone complex  $\Delta$  is a projective subdivision obtained by a sequence of simultaneous star subdivisions. If  $\Delta$  is nonsingular then the star subdivisions are smooth. The generalized cone complex  $B(\Delta)$  is in fact a cone complex.*
- (2) ([AMR99, Lemma 8.7]) *The barycentric subdivision  $B(\Delta)$  of a nonsingular cone complex  $\Delta$  is a projective subdivision obtained by a sequence of simultaneous smooth star subdivisions. The nonsingular cone complex  $B(\Delta)$  is in fact isomorphic to a fan.*

*Proof.* (1) Write  $\Delta = (\{\sigma_j\}, \{\mu_k\})$ . We need to show that if  $\tau_B, \sigma_B$  are cones in  $B(\Delta)$ , then a face map  $\tau_B \rightarrow \sigma_B$  in  $B(\Delta)$  is unique if it exists. Suppose the minimal cone containing the image of  $\tau_B$  is  $\tau$  and the corresponding cone for  $\sigma_B$  is  $\sigma$ . Then it suffices to show that the restriction to  $\tau_B$  of a face map  $\psi : \tau \rightarrow \sigma$  in  $\Delta$  carrying  $\tau_B$  into  $\sigma_B$  is unique if it exists. We can write  $\sigma_B = \langle b(\sigma_{i_1}), \dots, b(\sigma_{i_k}) \rangle$  uniquely as the cone generated by the barycenters  $b(\sigma_{i_r})$  of faces  $\sigma_{i_r}$  of  $\sigma$  of dimensions  $i_1 < \dots < i_k$ , and similarly  $\tau_B = \langle b(\tau_{j_1}), \dots, b(\tau_{j_l}) \rangle$ . So  $\psi$  must carry  $b(\tau_{j_s})$  to the barycenter of a cone of  $\sigma$  of dimension  $j_s$ , in other words  $\psi(b(\tau_{j_s})) = b(\sigma_{j_s})$ . Since  $\{b(\tau_{j_1}), \dots, b(\tau_{j_l})\}$  span  $\tau_B$  this means that the restriction of  $\psi$  is unique if it exists.


- (2) Consider the vector space  $V = \bigoplus_{\sigma \in \Delta} \mathbb{R}_\sigma$  with one basis element for each cone of  $\sigma$ . Assume  $\Delta$  is a cone complex. In [AMR99, Lemma 8.7] it is shown that  $B(\Delta)$  has a real embedding in  $V$ , and the image is the real support of a fan. The embedding is obtained by sending  $b(\sigma)$  to the unit

vector  $e_\sigma \in \mathbb{R}_\sigma \subset V$ . Here we assume that  $\Delta$  is nonsingular, and we need to check that the embedding gives an isomorphism of cone complexes, namely that the integral structures coincide. Note that the lattice in any cone  $\langle b(\sigma_{i_1}), \dots, b(\sigma_{i_k}) \rangle$  in  $B(\Delta)$  is generated by the elements  $b(\sigma_{i_1}), \dots, b(\sigma_{i_k})$ . The image of this lattice in  $V$  is precisely generated by  $e(\sigma_{i_1}), \dots, e(\sigma_{i_k})$ , and coincides with the intersection of the cone  $\langle e(\sigma_{i_1}), \dots, e(\sigma_{i_k}) \rangle$  with  $\bigoplus_{\sigma \in \Delta} \mathbb{Z}_\sigma$ . So the image of  $B(\Delta)$  is indeed a fan, as required. ♣

generalized-factorization

**Lemma 3.4.2.** *Let  $\Delta$  be a nonsingular generalized cone complex and  $f : \Delta \rightarrow \mathbb{R}$  a piecewise linear function, convex and integral on each cone, such that the corresponding subdivision  $\Delta_1 \rightarrow \Delta$  is nonsingular. Then  $\Delta_1 \rightarrow \Delta$  admits a factorization into nonsingular star subdivisions and their inverses, with all intermediate steps projective over  $\Delta$ .*

*Proof.* By Lemma 3.4.1 we may replace  $\Delta_1$  by its second barycentric subdivision, so we may assume  $\Delta_1$  is isomorphic to a fan. The common subdivision of  $B(B(\Delta_1))$  and  $B(B(\Delta))$  is a projective subdivision of  $B(B(\Delta_1))$ , so there is a sequence of star subdivisions  $\Delta'_1 \rightarrow B(B(\Delta_1))$  such that  $\Delta'_1 \rightarrow \Delta$  factors through a projective subdivision  $\Delta'_1 \rightarrow \Delta' := B(B(\Delta))$ . Since  $\Delta'$  is isomorphic to a fan and  $\Delta'_1$  is a projective subdivision, Morelli's  $\pi$  desingularization lemma applies, see [Mor96] or [Wlo03, Lemma 10.4.3], giving a factorization by star subdivisions and their inverses, all projective over  $\Delta'$ . Combining these transformation, we obtain the desired factorization, with all steps projective over  $\Delta$ :



♣

### 3.5. Functoriality for generalized cone complexes.

factorization-functorial

**Lemma 3.5.1.** *The factorization in Lemma 3.4.2 can be made functorial for surjective face maps: we can associate to  $(\Delta, f)$  a factorization so that, given a surjective face map  $\phi : \Sigma \rightarrow \Delta$ , the factorization of  $(\Sigma, f \circ \phi)$  is the pullback of the factorization of  $(\Delta, f)$  along  $\phi$ .*

*Proof.* For each connected component of the category of pairs  $(\Delta, f)$  with face maps between them choose a final object  $(\tilde{\Delta}, \tilde{f})$ . By Lemma 3.4.2 there is a factorization  $\tilde{\Delta}_1 \dashrightarrow \dots \dashrightarrow \tilde{\Delta}$  of  $(\tilde{\Delta}, \tilde{f})$ . Given an arbitrary  $(\Delta, f)$  it has a morphism  $\psi_\Delta : \Delta \rightarrow \tilde{\Delta}$  to the final object  $(\tilde{\Delta}, \tilde{f})$ , so that  $f = \tilde{f} \circ \psi_\Delta$ . The pullback  $\Delta_1 \dashrightarrow \dots \dashrightarrow \Delta$  of  $\tilde{\Delta}_1 \dashrightarrow \dots \dashrightarrow \tilde{\Delta}$  along  $\psi_\Delta$  is a factorization of  $(\Delta, f)$ , and its pullback along  $\phi$  is simply the pullback  $\Sigma_1 \dashrightarrow \dots \dashrightarrow \Sigma$  along  $\psi_\Delta \circ \phi = \psi_\Sigma$  of  $\tilde{\Delta}_1 \dashrightarrow \dots \dashrightarrow \tilde{\Delta}$ , so the process is functorial. ♣

### 3.6. Functoriality for toroidal factorization.

*Proof of Proposition 3.1.1.* The toroidal morphism  $X_1 \rightarrow X_2$  corresponds to a subdivision  $\Sigma(X_1) \rightarrow \Sigma(X_2)$  induced by a piecewise linear function  $f : \Sigma(X_2) \rightarrow \mathbb{R}$  convex and integral on each cone. This is functorial: a surjective regular morphism  $X'_2 \rightarrow X_2$  gives rise to a surjective face map  $\phi : \Sigma(X'_2) \rightarrow \Sigma(X_2)$  such that  $X'_1 \rightarrow X'_2$  corresponds to  $f \circ \phi$ .

By Lemma 3.5.1 we have a factorization  $\Sigma(X_1) \dashrightarrow \dots \dashrightarrow \Sigma(X_2)$ , functorial for surjective face maps, into nonsingular star subdivisions and their inverses, with all intermediate steps functorially projective over  $\Sigma(X_2)$ . This gives rise to a toroidal factorization  $X_1 \dashrightarrow \dots \dashrightarrow X_2$  into blowings up and down, which is functorial for surjective regular morphisms, where the terms are functorially projective over  $X_2$ .  $\clubsuit$

## 4. BIRATIONAL COBORDISMS

A key tool in the factorization algorithm is the notion of *birational cobordism*, introduced in [Wlo00], where it is motivated by analogy with Morse theory. In this paper we adopt the approach of [AKMW02] which relies on Geometric Invariant Theory and variation of linearizations, see [BP90, Tha96, DH98].

Sec:GIT-E

**4.1. Geometric Invariant Theory of  $\mathbb{P}(E)$ .** Given a nonzero coherent sheaf  $E$  on  $X_2$ , the data of a  $\mathbb{G}_m$ -action  $\rho : \mathbb{G}_m \rightarrow \text{Aut} E$  on  $E$  is equivalent to the data of a  $\mathbb{Z}$ -grading  $E = \bigoplus_{a \in \mathbb{Z}} E_a$ , which is necessarily a finite sum:  $E = \bigoplus_{a=a_{\min}}^{a_{\max}} E_a$ . The homogeneous factor  $E_a$  is characterized by

$$\rho(t)v = t^a v \quad \forall v \in E_a.$$

Here and later we use the informal notation  $v \in E_a$  to indicate that  $v$  is a local section of  $E_a$ . Given such data, there is a resulting action of  $\mathbb{G}_m$  on  $\text{Sym}^\bullet(E)$  and a linearized action on  $\mathbb{P}(E) = \mathbb{P}_{X_2}(E)$ .

We require the following:

**Assumption 4.1.1.** *The sheaves  $E_{a_{\min}}$  and  $E_{a_{\max}}$  are everywhere nonzero, so  $\mathbb{P}(E_{a_{\min}}) \rightarrow X_2$  and  $\mathbb{P}(E_{a_{\max}}) \rightarrow X_2$  are surjective.*

Given an integer  $a$  viewed as a character of  $\mathbb{G}_m$ , we define a new action of  $\mathbb{G}_m$  on  $E$  by

$$\rho_a(t)v = t^{-a} \rho(t)(v).$$

This induces an action on  $\text{Sym}^\bullet(E)$  and on  $(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(1))$  which we also denote by  $\rho_a$ . Writing  $(\text{Sym}^\bullet(E))^{\rho_a}$  for the ring of invariants under this action, we denote

$$\mathbb{P}(E) //_a \mathbb{G}_m := \text{Proj}_{X_2}(\text{Sym}^\bullet(E))^{\rho_a}.$$

As customary, we unwind this as follows: we define the *unstable locus* of  $\rho_a$  to be the closed subscheme

Eq:unstable (1) 
$$\mathbb{P}(E)_a^{\text{un}} := \mathbb{P}\left(\bigoplus_{b < a} E_b\right) \sqcup \mathbb{P}\left(\bigoplus_{b > a} E_b\right),$$

and the *semistable locus* to be the complementary open

$$\mathbb{P}(E)_a^{\text{sst}} := \mathbb{P}(E) \setminus \mathbb{P}(E)_a^{\text{un}}.$$

We have the following well-known facts:



**Lem:GIT-E**

- Lemma 4.1.2.** (1) *The semistable locus  $\mathbb{P}(E)_a^{\text{sst}}$  is nonempty precisely when  $a_{\min} \leq a \leq a_{\max}$ .*
- (2) *Consider the rational map  $q_a : \mathbb{P}(E) \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  induced by the inclusion  $(\text{Sym}^\bullet(E))^{\rho_a} \subset (\text{Sym}^\bullet(E))$ . Then  $q_a$  restricts to an affine  $\mathbb{G}_m$ -invariant morphism  $\mathbb{P}(E)_a^{\text{sst}} \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  which is a submersive universal categorical quotient, thus  $\mathbb{P}(E) //_a \mathbb{G}_m = \mathbb{P}(E)_a^{\text{sst}} // \mathbb{G}_m$ .*
- (3) *For  $a_{\min} \leq a_1 < a_2 \leq a_{\max}$  we have  $\mathbb{P}(E)_{a_1}^{\text{sst}} \subset \mathbb{P}(E)_{a_2}^{\text{sst}}$  precisely when  $\bigoplus_{a=a_1}^{a_2-1} E_a = 0$ , and similarly  $\mathbb{P}(E)_{a_1}^{\text{sst}} \supset \mathbb{P}(E)_{a_2}^{\text{sst}}$  precisely when  $\bigoplus_{a=a_1+1}^{a_2} E_a = 0$ . In particular  $\mathbb{P}(E)_{a_1}^{\text{sst}} = \mathbb{P}(E)_{a_2}^{\text{sst}}$  precisely when  $\bigoplus_{a=a_1}^{a_2} E_a = 0$ .*
- (4) *If  $a_{\min} \leq a_1 < a_2 \leq a_{\max}$  and  $\bigoplus_{a=a_1}^{a_2-1} E_a = 0$ , then the inclusion  $\mathbb{P}(E)_{a_1}^{\text{sst}} \subset \mathbb{P}(E)_{a_2}^{\text{sst}}$  induces a projective morphism*

$$\mathbb{P}(E)_{a_1}^{\text{sst}} // \mathbb{G}_m \rightarrow \mathbb{P}(E)_{a_2}^{\text{sst}} // \mathbb{G}_m.$$

*Similarly if  $\bigoplus_{a=a_1+1}^{a_2} E_a = 0$  we have a projective morphism*

$$\mathbb{P}(E)_{a_1}^{\text{sst}} // \mathbb{G}_m \leftarrow \mathbb{P}(E)_{a_2}^{\text{sst}} // \mathbb{G}_m.$$

*Proof.* (1) We have  $a \leq a_{\max}$  if and only if  $\mathbb{P}(\bigoplus_{b < a} E_b) \neq \mathbb{P}(E)$ , and  $a_{\min} \leq a$  if and only if  $\mathbb{P}(\bigoplus_{b > a} E_b) \neq \mathbb{P}(E)$ .

- (2) (a) **AFFINE COVER OF THE QUOTIENT.** The scheme  $\mathbb{P}(E) //_a \mathbb{G}_m = \text{Proj}_{X_2}(\text{Sym}^\bullet(E))^{\rho_a}$  is covered by principal open sets

**Eq:D0**

$$(2) \quad D_f^0 := (\mathbb{P}(E) //_a \mathbb{G}_m) \setminus Z_{\mathbb{P}(E) //_a \mathbb{G}_m}(f)$$

associated to non-zero homogeneous invariant elements of the form  $f = \prod_{j=1}^s f_j$  where  $f_j \in E_{a+\delta_j}$  with  $\sum \delta_j = 0$ .

- (b) **COMMON ZERO LOCUS OF  $\{f\}$ .** We note that the common zero locus of elements of  $E_c$  is  $\mathbb{P}(E/E_c) = \mathbb{P}(\bigoplus_{b \neq c} E_b)$ . Now observe that any element  $f = \prod_{j=1}^s f_j$  as above has a factor  $f_j$  with  $\delta_j \geq 0$  and a factor  $f_j$  with  $\delta_j \leq 0$ . This means that  $f$  vanishes on  $\mathbb{P}(\bigoplus_{b < a} E_b)$  and on  $\mathbb{P}(\bigoplus_{b > a} E_b)$ , so  $f$  vanishes on  $\mathbb{P}(E)_a^{\text{un}}$ .

Conversely if  $x \notin \mathbb{P}(E)_a^{\text{un}}$  then we have some coordinates  $f_1 \in E_{a+\delta_1}$ ,  $\delta_1 \leq 0$  and  $f_2 \in E_{a+\delta_2}$ ,  $\delta_2 \geq 0$  which do not vanish:  $f_1(x) \neq 0 \neq f_2(x)$ . Taking any positive  $r, s$  so that  $r\delta_1 + s\delta_2 = 0$  we can form  $f = f_1^r f_2^s$ , and  $f(x) \neq 0$ . This implies that the common zero locus of the elements  $f = \prod_{j=1}^s f_j$  above in  $\mathbb{P}(E)$  is precisely  $\mathbb{P}(E)_a^{\text{un}}$ .

- (c) **COMPATIBLE AFFINE COVER OF  $\mathbb{P}(E)_a^{\text{sst}}$ .** It follows that  $\mathbb{P}(E)_a^{\text{sst}}$  is covered by principal open sets

**Eq:D**

$$(3) \quad D_f = \mathbb{P}(E) \setminus Z_{\mathbb{P}(E)}(f),$$

the inverse image of the affine open  $D_f^0$  of equation (2) is the affine open  $D_f$  of equation (3), and  $\mathbb{P}(E)_a^{\text{sst}} \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  is an affine morphism.

- (d) **COORDINATES AND INVARIANTS** The coordinate ring of  $D_f^0$  is the degree-zero component of  $(\text{Sym}^\bullet(E))^{\rho_a}[1/f]$ , which is the  $\rho_a$ -invariant summand of the degree-0 component of  $(\text{Sym}^\bullet(E))[1/f]$ . The latter is the coordinate ring of  $D_f$ . In particular,  $D_f^0 = D_f // \mathbb{G}_m$  is a submersive universal categorical quotient, see [AT15a, Lemma 4.2.6 and Corollary 4.2.11]. It follows from the definition (see [MFK94, Remark

5, p. 8]) that  $\mathbb{P}(E)_a^{\text{sst}} \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  is a submersive universal categorical quotient.

- (3) The situation is symmetric, so we only address the first statement. If  $\bigoplus_{a=a_1}^{a_2-1} E_a = 0$  then  $\mathbb{P}(\bigoplus_{b < a_2} E_b) = \mathbb{P}(\bigoplus_{b < a_1} E_b) \subset \mathbb{P}(E)_{a_1}^{\text{un}}$  and certainly  $\mathbb{P}(\bigoplus_{b > a_2} E_b) \subset \mathbb{P}(\bigoplus_{b > a_1} E_b) \subset \mathbb{P}(E)_{a_1}^{\text{un}}$ , so  $\mathbb{P}(E)_{a_1}^{\text{un}} \subset \mathbb{P}(E)_{a_2}^{\text{un}}$  as needed.

Conversely, if  $v \in \mathbb{P}(\bigoplus_{a=a_1}^{a_2-1} E_a)$  over  $x \in X_2$  and we take  $w \in \mathbb{P}(E_{a_{\min}})$  also over  $x$ , then either  $v \in \mathbb{P}(E_{a_1}) \subset \mathbb{P}(E)_{a_1}^{\text{sst}}$  or else  $(v+w) \in \mathbb{P}(E)_{a_1}^{\text{sst}}$ . In either case, if  $\bigoplus_{a=a_1}^{a_2-1} E_a \neq 0$  we have  $\mathbb{P}(E)_{a_1}^{\text{sst}} \not\subset \mathbb{P}(E)_{a_2}^{\text{sst}}$ , as needed.

- (4) The situation is symmetric, so we only address the first case, where  $a_{\min} \leq a_1 < a_2 \leq a_{\max}$  and  $\bigoplus_{a=a_1}^{a_2-1} E_a = 0$ , so that  $\mathbb{P}(E)_{a_1}^{\text{sst}} \subset \mathbb{P}(E)_{a_2}^{\text{sst}}$  by (3). Since  $\mathbb{P}(E)_{a_i}^{\text{sst}} \rightarrow \mathbb{P}(E) //_{a_i} \mathbb{G}_m$  are categorical quotients, we have a canonical morphism  $\varphi_{a_1/a_2}$  making the following diagram commutative:

$$\begin{array}{ccc} \mathbb{P}(E)_{a_1}^{\text{sst}} & \hookrightarrow & \mathbb{P}(E)_{a_2}^{\text{sst}} \\ \downarrow & & \downarrow \\ \mathbb{P}(E) //_{a_1} \mathbb{G}_m & \xrightarrow{\varphi_{a_1/a_2}} & \mathbb{P}(E) //_{a_2} \mathbb{G}_m. \end{array}$$

But  $\mathbb{P}(E) //_{a_i} \mathbb{G}_m$  are projective over  $X_2$ , hence  $\varphi_{a_1/a_2}$  is projective. ♣

This lemma gives the familiar “wall and chamber decomposition” of the interval  $[a_{\min}, a_{\max}]$  in the character lattice  $\mathbb{Z}$  into segments where the quotients  $\mathbb{P}(E)_{a_1}^{\text{sst}} //_{\mathbb{G}_m}$  are constant.

All the constructions above are compatible with arbitrary morphisms  $X'_2 \rightarrow X_2$ , except that the values of  $a_{\min}$  and  $a_{\max}$  and the ample sheaf for  $\phi_{a_1/a_2}$  are only compatible with surjective morphisms  $X'_2 \rightarrow X_2$ .

Rem:geometric-quotient

**Remark 4.1.3.** One can show that the quotient morphism  $\mathbb{P}(E)_a^{\text{sst}} \rightarrow \mathbb{P}(E)_a^{\text{sst}} //_{\mathbb{G}_m}$  is in fact universally submersive. If in addition  $E_a = 0$  it can be shown that the quotient morphism is a universal geometric quotient  $\mathbb{P}(E)_a^{\text{sst}} \rightarrow \mathbb{P}(E)_a^{\text{sst}} //_{\mathbb{G}_m}$ . These facts follow from [MFK94, Theorem 1.1 and Amplification 1.3], which are stated for schemes over a field in characteristic 0 but apply here since  $\mathbb{G}_m$  is a linearly reductive group-scheme over  $\mathbb{Z}$ . Since we do not need these facts, we will not provide a detailed proof, though we will use the notation  $\mathbb{P}(E)_a^{\text{sst}} //_{\mathbb{G}_m}$  when  $E_a = 0$ .

GIT

**4.2. Geometric Invariant Theory of  $B \subset \mathbb{P}(E)$ .** Continuing the discussion, let  $B \subset \mathbb{P}(E)$  be a closed reduced  $\mathbb{G}_m$ -stable subscheme. It is the zero locus of a homogeneous and  $\mathbb{G}_m$ -homogeneous ideal  $I_B \subset \text{Sym}^\bullet E$ . We define  $B_a^{\text{un}} := B \cap \mathbb{P}(E)_a^{\text{un}}$  and  $B_a^{\text{sst}} := B \cap \mathbb{P}(E)_a^{\text{sst}}$ . The image of  $q_a : B_a^{\text{sst}} \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  is denoted  $B //_a \mathbb{G}_m$ . We have canonically  $B //_a \mathbb{G}_m = \text{Proj}_{X_2} ((\text{Sym}^\bullet E / I_B)^{\rho_a})$ . We write  $a_{\min}(B) = \min\{a \mid B \cap \mathbb{P}(E_a) \neq \emptyset\}$  and similarly  $a_{\max}(B) = \max\{a \mid B \cap \mathbb{P}(E_a) \neq \emptyset\}$ . We deduce the analogous, still well-known, facts, which follow immediately from Lemma 4.1.2:

Lem:GIT-B

**Lemma 4.2.1.** (1) *The semistable locus  $B_a^{\text{sst}}$  is nonempty precisely when  $a_{\min}(B) \leq a \leq a_{\max}(B)$ .*

- (2) *The map  $q_a : B_a^{\text{sst}} \rightarrow \mathbb{P}(E) //_a \mathbb{G}_m$  is an affine  $\mathbb{G}_m$ -invariant morphism, inducing a categorical quotient  $B_a^{\text{sst}} \rightarrow B_a^{\text{sst}} //_{\mathbb{G}_m} = B //_a \mathbb{G}_m$ .*

- (3) For  $a_1 < a_2$  we have  $B_{a_1}^{\text{sst}} \subset B_{a_2}^{\text{sst}}$  precisely when  $B \cap \mathbb{P}(\oplus_{a=a_1}^{a_2-1} E_a) = \emptyset$ , and similarly  $B_{a_1}^{\text{sst}} \supset B_{a_2}^{\text{sst}}$  precisely when  $B \cap \mathbb{P}(\oplus_{a=a_1+1}^{a_2} E_a) = \emptyset$ . In particular  $B_{a_1}^{\text{sst}} = B_{a_2}^{\text{sst}}$  precisely when  $B \cap \mathbb{P}(\oplus_{a=a_1}^{a_2} E_a) = \emptyset$ .
- (4) If  $a_1 < a_2$  and  $B \cap \mathbb{P}(\oplus_{a=a_1}^{a_2-1} E_a) = \emptyset$ , then the inclusion  $B_{a_1}^{\text{sst}} \subset B_{a_2}^{\text{sst}}$  induces a projective morphism  $B_{a_1}^{\text{sst}} // \mathbb{G}_m \rightarrow B_{a_2}^{\text{sst}} // \mathbb{G}_m$ . Similarly if  $B \cap \mathbb{P}(\oplus_{a=a_1+1}^{a_2} E_a) = \emptyset$  we have a projective morphism  $B_{a_1}^{\text{sst}} // \mathbb{G}_m \leftarrow B_{a_2}^{\text{sst}} // \mathbb{G}_m$ .

This time we obtain a “wall and chamber decomposition” of the interval  $[a_{\min}(B), a_{\max}(B)]$ . We denote the “walls”, namely the values of  $a$  for which  $B \cap \mathbb{P}(E_a) \neq \emptyset$ , by  $a_{\min}(B) = a_0 < a_1 \cdots < a_m = a_{\max}(B)$ .

By replacing the embedding  $B \subset \mathbb{P}(E)$  by the Veronese re-embedding  $B \subset \mathbb{P}(\text{Sym}^2 E)$  we may, and will, assume

Ass: separate

**Assumption 4.2.2.**  $a_i + 1 < a_{i+1}$ .

We denote  $B_{a_i+}^{\text{sst}} = B_{a_{i+1}}^{\text{sst}}$  and  $B_{a_i-}^{\text{sst}} = B_{a_{i-1}}^{\text{sst}}$ , and note that  $B_{a_i+}^{\text{sst}} = B_{a_{i+1}-}^{\text{sst}}$ . Assumption 4.2.2 implies that now we always have projective morphisms  $\varphi_{a_i \pm}$ :

$$(4) \quad \begin{array}{c} B_{a_i-}^{\text{sst}} / \mathbb{G}_m \dashrightarrow B_{a_i+}^{\text{sst}} / \mathbb{G}_m = B_{a_{i+1}-}^{\text{sst}} / \mathbb{G}_m \dashrightarrow \cdots \\ \varphi_{a_i-} \searrow \quad \swarrow \varphi_{a_i+} \\ B_{a_i}^{\text{sst}} // \mathbb{G}_m \\ \searrow \quad \swarrow \\ B_{a_{i+1}}^{\text{sst}} // \mathbb{G}_m \end{array}$$

Finally, we will assume the following:

Ass: spread

**Assumption 4.2.3.** Each irreducible component of  $B$  meets both  $\mathbb{P}(E_{a_{\min}(B)})$  and  $\mathbb{P}(E_{a_{\max}(B)})$ .

Under this assumption the quotients  $B_a^{\text{sst}} // \mathbb{G}_m$  are all birational to each other, as long as  $a_{\min}(B) < a < a_{\max}(B)$ . For the extreme values we have isomorphisms  $B \cap \mathbb{P}(E_{a_{\min}(B)}) \rightarrow B_{a_{\min}(B)}^{\text{sst}} // \mathbb{G}_m$  and  $B \cap \mathbb{P}(E_{a_{\max}(B)}) \rightarrow B_{a_{\max}(B)}^{\text{sst}} // \mathbb{G}_m$ .

**Remark 4.2.4.** As in Remark 4.1.3, it can be shown that  $B_a^{\text{sst}} \rightarrow B_a^{\text{sst}} // \mathbb{G}_m$  is universally submersive, and if  $B \cap \mathbb{P}(E_a) = \emptyset$  we have a universal geometric quotient  $B_a^{\text{sst}} \rightarrow B_a^{\text{sst}} / \mathbb{G}_m$ .

Sec: def-cobordism

**4.3. Definition of a birational cobordism.** The notion of a birational cobordism for a blowing up we use in this paper extends the notion of *compactified relatively projective embedded birational cobordism* of [AKMW02, 2.4] by allowing a non-empty boundary. Ignoring the issue of the boundary, it is far more restrictive than the notion introduced in [Wto00].

Let  $\phi: X_1 \rightarrow X_2$  be an object of  $\text{Bl}_{\text{rs}}$ . A birational cobordism for  $\phi$  is a scheme  $B$  which is the blowing up of a  $\mathbb{G}_m$ -invariant ideal on  $\mathbb{P}_{X_2}^1$ , and embedded, in a manner satisfying Assumptions 4.2.2 and 4.2.3, as a  $\mathbb{G}_m$ -stable subscheme in  $\mathbb{P}(E)$  for a  $\mathbb{G}_m$ -sheaf  $E$  on  $X_2$ , such that

- (1)  $X'_1 = B_{a_0+}^{\text{sst}} / \mathbb{G}_m = B_{a_0}^{\text{sst}} // \mathbb{G}_m$  is obtained from  $X_1$  by principalizing  $D_1$ ,
- (2)  $X'_2 = B_{a_m-}^{\text{sst}} / \mathbb{G}_m = B_{a_m}^{\text{sst}} // \mathbb{G}_m$  is obtained from  $X_2$  by principalizing  $D_2$ ,  
and
- (3) the following diagram of rational maps commutes:

$$\begin{array}{ccccc}
B_{a_0}^{\text{sst}} & \xrightarrow{q_{a_0}} & X'_1 & \longrightarrow & X_1 \\
\downarrow \alpha & & & & \downarrow \phi \\
B_{a_m}^{\text{sst}} & \xrightarrow{q_{a_m}} & X'_2 & \longrightarrow & X_2
\end{array}$$

where  $\alpha$  is the birational map induced by the open dense inclusions

$$B_{a_0}^{\text{sst}} \subset B \supset B_{a_m}^{\text{sst}}.$$

The birational cobordism is said to respect the open set  $U \subset X_2$  if  $U$  is contained in the image of  $(B_{a_0+}^{\text{sst}} \cap B_{a_m-}^{\text{sst}})/\mathbb{G}_m$ . This happens whenever the ideal on  $\mathbb{P}_{X_2}^1$  whose blowing up is  $B$  restricts to the unit ideal on  $\mathbb{P}_U^1$ . We say that a birational cobordism  $B$  of  $\phi$  is *regular* if  $B$  is regular and the preimage  $D_B$  of  $D_2$  is a simple normal crossings divisor.

**4.4. Construction of regular birational cobordism.** We claim that one can associate a regular birational cobordism to any blowing up in  $\text{Bl}_{\text{rs}}$  functorially, and we formalize this claim as follows. There is an evident category  $\text{Cob}_{\text{rs}}$  of regular birational cobordisms of blowings up  $\phi: X_1 \rightarrow X_2$  in  $\text{Bl}_{\text{rs}}$ , with an evident forgetful functor  $\text{Cob}_{\text{rs}} \rightarrow \text{Bl}_{\text{rs}}$ . A morphism of regular birational cobordisms  $B' \rightarrow B$  is uniquely determined by a regular surjective morphism  $g: X'_2 \rightarrow X_2$ .

Prop: cobordism

**Proposition 4.4.1.** *The functor  $\text{Cob}_{\text{rs}} \rightarrow \text{Bl}_{\text{rs}}$  has a section  $\text{Bl}_{\text{rs}} \rightarrow \text{Cob}_{\text{rs}}$ .*

We provide a sketch of proof here, and more detail in Appendix A.

*Sketch of proof.* Following the construction of [AKMW02, Theorem 2.3.1], consider the blowing up of the ideal  $I \otimes \mathcal{O}_{\mathbb{P}_{X_2}^1} + I_{\{0\}}$ . This is a birational cobordism  $B_I$  for  $\phi$ , but it is singular. We apply canonical resolution of singularities to  $B_I$ , which is a functorial sequence of blowings up  $B^{\text{reg}} \rightarrow B$ : see Section 2.1.2, where in positive and mixed characteristic we may use Hypothetical Statement 2.1.4 since  $\dim B_I = \dim X_2 + 1$ . Then we principalize the preimage of  $D_2$  in  $B^{\text{reg}}$  by a blowing up sequence  $B \rightarrow B^{\text{reg}}$ , obtaining a regular birational cobordism  $(B, D_B)$  for  $\phi$ . The last step is non-trivial only when  $D_2 \neq \emptyset$  and then, in the positive and mixed characteristic, it involves Hypothetical Statement 2.2.1. ♣

## 5. FACTORING THE MAP

Throughout this section “functorial” means “functorial in  $X_1 \rightarrow X_2$  with respect to surjective regular morphisms”. By *total transform* of a divisor  $D \subset X$  under a (normalized) blowing up  $Bl_J(X) \rightarrow X$  we mean the union of the preimage of  $D$  and the total transform of  $J$ .

## 5.1. Blowing up the toric ideals.

5.1.1. *Initial factorization.* Proposition 4.4.1 provides a functorial birational cobordism  $(B, D_B)$  of  $\phi$ . Departing slightly from the notation of [AKMW02, Theorem 2.6.2], we write  $W_{i\pm} = B_{a_i\pm}^{\text{sst}}/\mathbb{G}_m$ , and  $W_i = B_{a_i}^{\text{sst}} // \mathbb{G}_m$ . Since  $W_{i+} \simeq W_{(i+1)-}$  we have a functorial factorization

Eq: wtd-fact

$$(5) \quad \begin{array}{ccccccc} & & W_{1-} & & W_{2-} & & W_{m-} \\ & \swarrow^{\varphi_{0+}} & & \swarrow^{\varphi_{1-}} \quad \searrow^{\varphi_{1+}} & & \swarrow^{\varphi_{(m-1)+}} & \searrow^{\varphi_{m-}} \\ X'_1 = W_0 & & & W_1 & & \dots & W_m = X'_2 \end{array}$$

with all terms functorially projective over  $X_2$ . Since the cobordism is compatible with  $U$ , the morphisms  $W_{i\pm} \rightarrow X_2$  and  $W_i \rightarrow X_2$  and hence also the morphisms  $\varphi_{i\pm}$  are isomorphisms on  $U$ . Note that since  $W_{m-1} \dashrightarrow W_m$  is a morphism it follows that  $\varphi_{(m-1)+}$  is an isomorphism, but this fact does not feature in our arguments. In general the terms  $W_i$  and  $W_{i\pm}$  in this factorization are singular, but we will use them to construct a non-singular factorization.

5.1.2. *Toric ideals.* Let  $D_i \subset W_i$ ,  $D_{i\pm} \subset W_{i\pm}$ ,  $D_{a_i} \subset B_{a_i}^{\text{sst}}$  and  $D_{a_i\pm} \subset B_{a_i\pm}^{\text{sst}}$  denote the preimages of  $D_2$ . We will show how main results of [AT15b] imply that since  $(W_i, D_i)$  is given as a quotient of  $(B_{a_i}^{\text{sst}}, D_{a_i})$ , it can be made toroidal by a canonical toric blowing up. Since  $B$  is regular and  $D_B$  is a simple normal crossings divisor,  $(B_{a_i}^{\text{sst}}, D_{a_i})$  is a toroidal scheme with a relatively affine  $\mathbb{G}_m$ -action. In [AT15b, Section 4.2] one functorially associates to  $(B_{a_i}^{\text{sst}}, D_{a_i})$  a  $\mathbb{G}_m$ -equivariant *toric ideal*  $J_i^B$  on  $B_{a_i}^{\text{sst}}$ . We define ideal sheaf  $J_i$  on  $W_i$  to be the  $\mathbb{G}_m$ -invariant part of  $J_i^B$ , and we call it the *invariant toric ideals*. By abuse of language, the ideal sheaves  $J_{i\pm} = J_i \mathcal{O}_{W_{i\pm}}$  will also be called invariant toric ideals.

**Theorem 5.1.3.** *For every  $1 \leq i \leq (m-1)$  the ideal sheaves  $J_i$  and  $J_{i\pm}$  are functorial and restrict to the unit ideal on  $U$ . Furthermore, let  $W_i^{\text{tor}} = Bl_{J_i} W_i$ <sup>2</sup> and  $W_{i\pm}^{\text{tor}} = Bl_{J_{i\pm}} W_{i\pm}$ , and denote by  $D_i^{\text{tor}} \subset W_i^{\text{tor}}$  and  $D_{i\pm}^{\text{tor}} \subset W_{i\pm}^{\text{tor}}$  the total transforms of  $D_i$  and  $D_{i\pm}$ , respectively. Then*

- (1)  $(W_i^{\text{tor}}, D_i^{\text{tor}})$  and  $(W_{i\pm}^{\text{tor}}, D_{i\pm}^{\text{tor}})$  are toroidal, and
- (2) the morphisms  $\varphi_{i\pm}$  induce toroidal morphisms

$$\varphi_{i\pm}^{\text{tor}} : (W_{i\pm}^{\text{tor}}, D_{i\pm}^{\text{tor}}) \rightarrow (W_i^{\text{tor}}, D_i^{\text{tor}})$$

that restrict to isomorphisms on  $U$ .

*Proof.* By [AT15b, Lemma 4.2.12]  $\mathbb{G}_m$  acts in a relatively affine way on  $B_{a_i}^{\text{tor}} := Bl_{J_i^B}(B_{a_i}^{\text{sst}})$ . Let  $D_{a_i}^{\text{tor}} \subset B_{a_i}^{\text{tor}}$  be the total transform of  $D_{a_i}$ , then by [AT15b, Theorem 1.1.2],  $(B_{a_i}^{\text{tor}}, D_{a_i}^{\text{tor}})$  is a toroidal scheme with toroidal action of  $\mathbb{G}_m$ , and

<sup>2</sup>(Michael) In fact this should be normalized blowing up. I don't change this now, since we may want to switch to blowings up in the end, using the normalization/saturation trick.

$W_i^{\text{tor}} = B_{a_i}^{\text{tor}} // \mathbb{G}_m$ . Note that  $D_i^{\text{tor}}$  is the image of  $D_{a_i}^{\text{tor}}$ , hence  $(W_i^{\text{tor}}, D_i^{\text{tor}})$  is toroidal by [AT15b, Theorem 1.1.3].

By [AT15b, Lemma 4.2.14],  $W_{i\pm}^{\text{tor}} = (B_{a_i}^{\text{tor}})_{\pm} // \mathbb{G}_m$ . Set  $(D_{a_i}^{\text{tor}})_{\pm} = D_{a_i}^{\text{tor}}|_{(B_{a_i}^{\text{tor}})_{\pm}}$ , then  $\mathbb{G}_m$  acts toroidally on  $((B_{a_i}^{\text{tor}})_{\pm}, (D_{a_i}^{\text{tor}})_{\pm})$  and hence the quotient  $(W_{i\pm}^{\text{tor}}, D_{i\pm}^{\text{tor}})$  is toroidal by [AT15b, Theorem 1.1.3]. Note also that  $\varphi_{i\pm}$  induce toroidal morphisms  $\varphi_{i\pm}^{\text{tor}}$  by [AT15b, Proposition 3.3.14].

3→ Functoriality of  $J_i^B$  implies that  $J_i$  and  $J_{i\pm}$  are functorial too. Note also that  $J_i^B$  is the unit ideal over  $U$ , as it is generated by monomials in semi-invariant coordinates along the fixed point set, and these coordinates are units over  $U$ . Since  $X_2$  is a qe scheme, normalization is finite. So  $B_{a_i}^{\text{tor}} \rightarrow B_{a_i}^{\text{sst}}$  is functorially projective.<sup>3</sup> ♣

We note that in general  $W_{i+}^{\text{tor}} \neq W_{(i+1)-}^{\text{tor}}$ . The steps  $W_{i-} \rightarrow W_i \leftarrow W_{i+}$  in the factorization (5) now look as follows:

Eq:tor-fact

$$(6) \quad \begin{array}{ccccc} & W_{i-}^{\text{tor}} & & W_{i+}^{\text{tor}} & \\ & \searrow \varphi_{i-}^{\text{tor}} & & \swarrow \varphi_{i+}^{\text{tor}} & \\ & & W_i^{\text{tor}} & & \\ & \swarrow \varphi_{i-} & \downarrow & \searrow \varphi_{i+} & \\ W_{(i-1)+} = W_{i-} & & W_i & & W_{i+} = W_{(i+1)-} \end{array}$$

**Remark 5.1.4.** In [AKMW02, Lemma 3.2.8] it is stated with a sketch of proof that the ideals  $J_i$  can be chosen so that  $\varphi_{i\pm}^{\text{tor}}$  are isomorphisms. We will not use this statement. We note however that this follows from [Tha96, Theorem 3.5]: if the  $l$ -toric ideal  $I_l$  generates all  $I_M, M \geq 1$  and also  $I_{-l}$  generates all  $I_{-M}, M \geq 1$ , then once  $l, -l \in S_i$ , the ample set of characters on  $B_{a_i}^{\text{sst}}$  used to determine  $J_i^B$  in [AT15b], then  $\varphi_{i\pm}^{\text{tor}}$  are isomorphisms. One can choose such  $l$  in a manner functorial for regular surjective morphisms.

Sec:Tying

## 5.2. Tying the maps together.

5.2.1. *Canonical resolution and principalization.* Extending [AKMW02, Section 4.2] to qe schemes with a boundary, we write  $W_{i\pm}^{\text{res}} \rightarrow W_{i\pm}$  for the composition of the canonical resolution of singularities  $W'_{i\pm} \rightarrow W_{i\pm}$  and the canonical principalization  $W_{i\pm}^{\text{res}} \rightarrow W'_{i\pm}$  of the preimage  $D'_{i\pm}$  of  $D_2$  in  $W'_{i\pm}$ , see Sections 2.1.2 and 2.2; in positive and mixed characteristics we require Hypothetical Statements 2.1.4 and 2.2.4. It is obtained by blowing up a functorial ideal on  $W_{i\pm}$  which is the unit ideal on  $U$ . By the construction,  $W_{i\pm}^{\text{res}}$  is regular and the preimage  $D_{i\pm}^{\text{res}}$  of  $D_2$  is a simple normal crossings divisor.

Note that the resolution and principalization processes are independent of the toroidal structures and hence coincide for  $(W_{(i-1)+}, D_{(i-1)+}) = (W_{i-}, D_{i-})$ , in other words  $(W_{(i-1)+}^{\text{res}}, D_{(i-1)+}^{\text{res}}) = (W_{i-}^{\text{res}}, D_{i-}^{\text{res}})$ . This provides a bridge between  $W_{(i-1)+}^{\text{tor}}$  and  $W_{i-}^{\text{tor}}$ .

<sup>3</sup>(Michael) There is a problem here: it is not clear how to make normalization functorially projective. Maybe will have to use that the normalized blow up along  $I$  is the blow up along something like  $(I^n)^{\text{nor}}$ . Anyway, this should be addressed in §2.4.

princrem

**Remark 5.2.2.** Since  $W_{1-} = X'_1$  is regular,  $X''_1 := W_{1-}^{\text{res}}$  is obtained from  $X'_1$  by principalization of  $D'_1$  and similarly  $X''_2 := W_{m-}^{\text{res}}$  is obtained from  $X'_2$  by principalization of  $D'_2$ . Both  $D'_1$  and  $D'_2$  are simple normal crossings divisors, so we could alternatively take  $W_{1-}^{\text{res}} = X'_1$  and  $W_{m-}^{\text{res}} = X'_m$ . Our choice helps to make notation uniform, though it results in a slightly longer factorization.

**Remark 5.2.3.** Note that the singularities requiring resolution in this step are far from general: it follows from Section 5.2.4 below that Zariski locally one can obtain a toroidal scheme from  $(W_{i\pm}, D_{i\pm})$  simply by enlarging the divisor  $D_{i\pm}$ . At least over an algebraically closed field they admit resolution of singularities, see [Wlo03, Theorem 8.3.2], and it seems reasonable to expect the same in general, and in a functorial manner.

torchart

5.2.4. *Local toroidal charts.* Our next aim is to connect  $W_{i\pm}^{\text{res}}$  and  $W_{i\pm}^{\text{tor}}$ . We will work locally on  $W_{i\pm}$ . For concreteness, fix a point  $x \in W_{i-}$  and consider the localization  $W_x := \text{Spec } \mathcal{O}_{W_{i-}, x}$ . We denote  $W_x^{\text{res}} = W_{i-}^{\text{res}} \times_{W_{i-}} W_x$  and similarly for  $W_x^{\text{tor}}$  and other  $W_{i-}$ -schemes we will introduce later. For shortness, we also write  $B_x = B_{a_{i-}}^{\text{sst}} \times_{W_{i-}} W_x$ ,  $D_{B_x} = D_{a_{i-}} \times_{W_{i-}} W_x$  and  $D_x = D_{i-} \times_{W_{i-}} W_x$ . By [AT15b, Proposition 3.4.1] there is an auxiliary toroidal structure  $\overline{D}_{B_x}$  on  $B_x$  that enlarges  $D_{B_x}$  and on which the action of  $\mathbb{G}_m$  is toroidal. Let  $\overline{D}_x$  be the image of  $\overline{D}_{B_x}$  in  $W_x$ .

For an fs torsion free monoid  $M$  we will use the notation  $\mathbf{A}_M = \text{Spec } \mathbb{Z}[M]$  and  $\overline{E}_M = \mathbf{A}_M \setminus \mathbf{A}_{M^{\text{gp}}}$ . By [AT15b, Corollary 3.2.11] there is a strongly equivariant toroidal chart  $(B_x, \overline{D}_x) \rightarrow (\mathbf{A}_{M_B}, \overline{E}_{M_B})$ . The  $\mathbb{G}_m$ -action induces a grading on  $M_B$ , and we set  $M = (M_B)_0$  and  $Y = \mathbf{A}_M$ . By [AT15b, Lemma 4.3.5] the toric ideal of  $B_x$  is the pullback of the toric ideal of  $\mathbf{A}_{M_B}$ , both ideals are toroidal and the toric blowing up of  $B_x$  is the base change of the toric blowing up of  $\mathbf{A}_{M_B}$ . Furthermore, taking invariants we obtain a local toroidal chart  $(W_x, \overline{D}_x) \rightarrow (Y, \overline{E}_M)$ , and again the invariant toric ideals  $J_{W_x} = J_{i-} \mathcal{O}_{W_x}$  and  $J_Y$  are toroidal and satisfy  $J_{W_x} = J_Y \mathcal{O}_{W_x}$ . Thus, writing  $Y^{\text{tor}} \rightarrow Y$  for the invariant toric blowing up of  $Y$ , we have  $W_x^{\text{tor}} = W_x \times_Y Y^{\text{tor}}$ . Notice also that  $W_{i\pm}$  and the ideal  $J_{i\pm}$  are locally monoidal.

The auxiliary toroidal structure was used to find the strictly equivariant chart and to establish compatibility of toric blowings up, since the latter are toroidal only with respect to the auxiliary structures. However, a posteriori the construction extends to  $(W_x, D_x)$  as follows: it is proved in [AT15b, 4.4.5] that  $D_{B_x}$  is the preimage of an equivariant toric divisor  $E_{M_B} \subseteq \overline{E}_{M_B}$ , and hence  $D_x$  is the preimage of a toric divisor  $E \subseteq \overline{E}_M$ , the image of  $E_{M_B}$  under  $\mathbf{A}_{M_B} \rightarrow \mathbf{A}_M$ . In particular,  $(Y, E)$  is a toroidal chart for  $(W_x, D_x)$ , and if  $E^{\text{tor}}$  and  $\overline{E}^{\text{tor}}$  denote the total transforms of  $E$  and  $\overline{E}$ , respectively, then  $(Y^{\text{tor}}, E^{\text{tor}})$  and  $(Y^{\text{tor}}, \overline{E}^{\text{tor}})$  are toroidal charts for  $(W_x^{\text{tor}}, D_x^{\text{tor}})$  and  $(W_x^{\text{tor}}, \overline{D}_x^{\text{tor}})$ , respectively.

5.2.5. *Principalization of toric ideals.* Consider the resolution of singularities  $Y' \rightarrow Y$  and let  $Y^{\text{res}} \rightarrow Y'$  be the principalization of the preimage of  $E$  in  $Y'$ . By Hypothetical Statements 2.1.4(3) and 2.2.4 resolution and principalization are compatible with toroidal charts, in particular,  $W_x^{\text{res}} = W_x \times_Y Y^{\text{res}}$  and hence  $Y^{\text{res}}$  is a toric chart for  $W_x^{\text{res}}$  and the ideal  $J_{i\pm} \mathcal{O}_{W_x^{\text{res}}} = J_Y \mathcal{O}_{W_x^{\text{res}}}$  comes from a toroidal ideal on  $Y^{\text{res}}$ . This proves that the ideal  $J_{i\pm}^{\text{res}} := J_{i\pm} \mathcal{O}_{W_{i\pm}^{\text{res}}}$  on  $W_{i\pm}^{\text{res}}$  is locally monoidal. Let  $W_{i\pm}^{\text{can}}$  denote the canonical principalization of  $J_{i\pm}^{\text{res}}$ , see Section 2.2. It is obtained by a functorial sequence of blowings up of nonsingular centers disjoint from  $U$  starting

from  $W_{i\pm}^{\text{res}}$ , see Proposition 2.2.2; in positive and mixed characteristics we require Hypothetical Statement 2.2.4.

By the universal property of blowing up, the maps  $W_{i\pm}^{\text{can}} \dashrightarrow W_{i\pm}^{\text{tor}}$  are morphisms. The map  $W_{i\pm}^{\text{can}} \rightarrow W_i$  is a composition of maps given functorially by blowing up ideals restricting to the unit ideal on  $U$ . By Section 2.3.10 the morphism  $W_{i\pm}^{\text{can}} \rightarrow W_i$  itself is given by blowing up a functorial ideal  $\tilde{J}_{i\pm}^{\text{can}}$  restricting to the unit ideal on  $U$ . So, by Lemma 2.3.9 the morphism  $W_{i\pm}^{\text{can}} \rightarrow W_{i\pm}^{\text{tor}}$  is given by blowing up the functorial ideal  $J_{i\pm}^{\text{can}} = \tilde{J}_{i\pm}^{\text{can}} \mathcal{O}_{W_{i\pm}^{\text{tor}}}$ . By  $D_{i\pm}^{\text{can}}$  we denote the total transform of  $D_{i\pm}^{\text{tor}}$ . In the local setting we also consider the auxiliary toroidal divisor  $\bar{D}_x^{\text{can}}$  which is the total transform of  $\bar{D}_x^{\text{tor}}$ . Diagram (6) now looks as follows:

Eq: can-fact

$$(7) \quad \begin{array}{ccccc} & & W_{i-}^{\text{can}} & & W_{i+}^{\text{can}} \\ & & \swarrow & & \swarrow \\ W_{(i-1)+}^{\text{res}} = W_{i-}^{\text{res}} & & W_{i-}^{\text{tor}} & & W_{i+}^{\text{tor}} \\ & & \swarrow & & \swarrow \\ W_{(i-1)+} & & W_i & & W_{i+} \\ & & \swarrow & & \swarrow \\ W_{(i-1)+} = W_{i-} & & W_i & & W_{i+} = W_{(i+1)-} \end{array}$$

(Note: The diagram above is a simplified representation of the complex commutative diagram in the image. The original diagram shows a network of nodes and arrows. The top row has  $W_{i-}^{\text{can}}$  and  $W_{i+}^{\text{can}}$ . Below them are  $W_{(i-1)+}^{\text{res}} = W_{i-}^{\text{res}}$  and  $W_{i+}^{\text{tor}}$ . Below those are  $W_{i-}^{\text{tor}}$  and  $W_{i+}^{\text{res}} = W_{(i+1)-}$ . Below those are  $W_{(i-1)+}$  and  $W_{i+}$ . The bottom row has  $W_{(i-1)+} = W_{i-}$  and  $W_{i+} = W_{(i+1)-}$ . Arrows connect these nodes, with labels  $\varphi_{i-}^{\text{tor}}$ ,  $\varphi_{i+}^{\text{tor}}$ ,  $\varphi_{i-}$ , and  $\varphi_{i+}$  indicating the maps between them.)

**Lemma 5.2.6.** *The ideal  $J_{i\pm}^{\text{can}}$  is toroidal. Thus,  $(W_{i\pm}^{\text{can}}, D_{i\pm}^{\text{can}}) \rightarrow (W_{i\pm}^{\text{tor}}, D_{i\pm}^{\text{tor}})$  is a functorial toroidal blowing up.*

*Proof.* STEP 1: REDUCTION TO TORIC CASE. Without loss of generality it suffices to deal with  $W_{i-}^{\text{can}} \rightarrow W_{i-}^{\text{tor}}$ , and we will work locally on  $W_{i-}$ . The blowing up  $W_x^{\text{can}} \rightarrow W_x^{\text{res}}$  is the canonical principalization of the toric ideal, which comes from  $Y$  through  $Y^{\text{res}}$ , so again by functoriality of 2.2.2 or Hypothetical Statement 2.2.4(3) we have  $W_x^{\text{can}} = W_x \times_Y Y^{\text{can}}$ , where the toric morphism  $Y^{\text{can}} \rightarrow Y^{\text{res}}$  is the principalization of  $J_Y \mathcal{O}_{Y^{\text{res}}}$ . By the functorial property of blowings up,  $Y^{\text{can}} \rightarrow Y$  factors through  $Y^{\text{tor}}$ . We have that  $Y^{\text{can}} = \text{Bl}_{\tilde{J}_Y^{\text{can}}}(Y)$  for a functorial ideal  $\tilde{J}_Y^{\text{can}}$  on  $Y$ , and by Lemma 2.3.9,  $Y^{\text{can}} = \text{Bl}_{J_Y^{\text{can}}}(Y)$ , where  $J_Y^{\text{can}} = \tilde{J}_Y^{\text{can}} \mathcal{O}_{Y^{\text{tor}}}$ . Let  $E^{\text{can}}$  and  $\bar{E}^{\text{can}}$  denote the total transforms of  $E^{\text{tor}}$  and  $\bar{E}^{\text{tor}}$ , respectively. Once again,  $(Y^{\text{can}}, E^{\text{can}})$  and  $(Y^{\text{can}}, \bar{E}^{\text{can}})$  are toroidal charts for  $(W_x^{\text{can}}, D_x^{\text{can}})$  and  $(W_x^{\text{can}}, \bar{D}_x^{\text{can}})$ . In the same fashion,  $J_x^{\text{can}} = J_Y^{\text{can}} \mathcal{O}_{W_x^{\text{can}}}$  and therefore it suffices to prove that the ideal  $J_Y^{\text{can}}$  is toroidal.

STEP 2: PROOF IN THE TORIC CASE. In [AKMW02, Proposition 4.2.1] it is shown that  $(Y^{\text{can}}, E^{\text{can}}) \rightarrow (Y^{\text{tor}}, E^{\text{tor}})$  is toroidal,<sup>4</sup> except that the ideal blown up is not shown to be toroidal. This can be shown as follows. In [AKMW02, Proposition 4.2.2] one constructs an action of  $\mathbb{G}_a^k$  on  $Y$ . One shows that the morphism  $Y^{\text{tor}} \rightarrow Y$  of charts is equivariant under this action, as well as the toric ideal; the scheme  $Y^{\text{tor}}$  is written as a product of  $\mathbb{G}_a^k$  with a toric scheme providing its toroidal structure. It suffices to show that the ideal defining the blowing up  $Y^{\text{can}} \rightarrow Y^{\text{tor}}$  is a  $\mathbb{G}_a^k$ -equivariant monomial ideal, since then its generating monomials are not divisible by the coordinates of the  $\mathbb{G}_a^k$  factor.

<sup>4</sup>(Michael) In the particular case when  $E = \emptyset$ .




Since the blowing up  $Y^{\text{res}} \rightarrow Y$  is the canonical resolution of singularities, the ideal defining this blowing up on a toric chart is monomial and  $\mathbb{G}_a^k$ -equivariant. Also the toric ideal on  $Y^{\text{res}}$  is monomial and  $\mathbb{G}_a^k$ -equivariant, therefore the same is true for the ideal defining its functorial principalization  $Y^{\text{can}} \rightarrow Y^{\text{res}}$ .  $\clubsuit$

The above lemma implies that the composition  $W_{i\pm}^{\text{can}} \rightarrow W_i^{\text{tor}}$  is a toroidal morphism given by blowing up a functorial toroidal ideal we denote by  $\bar{J}_{i\pm}^{\text{can}}$ . Let  $W'_i \rightarrow W_i^{\text{tor}}$  be the normalized blowing up of the product ideal  $\bar{J}_{i-}^{\text{can}} \bar{J}_{i+}^{\text{can}}$ , giving rise to toroidal morphisms  $W'_i \rightarrow W_{i\pm}^{\text{can}}$ . By [TT14, Theorem 3.4.9] there is a functorial toroidal resolution of singularities  $W_i^{\text{torres}} \rightarrow W'_i$ . This gives the following:


**Lemma 5.2.7.** *There is a toroidal nonsingular modification  $W_i^{\text{torres}} \rightarrow W_i^{\text{tor}}$  obtained by blowing up a functorial ideal, such that the maps  $W_i^{\text{torres}} \dashrightarrow W_{i\pm}^{\text{can}}$  are both toroidal morphisms.*

Note that these latter maps are again blowings up of the pullbacks of the ideal defining  $W_i^{\text{torres}} \rightarrow W_i^{\text{tor}}$ , which is functorial as well. Since the morphism is toroidal, it induces the identity on  $U$ , and the toroidal ideal blown up is the unit ideal on  $U$ .

We now have pieces of the diagram above looking as follows:



All maps are functorially the blowings up of ideals. The top diamond is at the same time toroidal, with maps given by blowings up of functorial toroidal ideals, so the toroidal structure is functorial in  $X_1 \rightarrow X_2$ . By Proposition 3.1.1, the two top maps  $W_i^{\text{torres}} \rightarrow W_{i\pm}^{\text{can}}$  have a functorial toroidal weak factorization; since it is toroidal it induces isomorphisms on  $U$ . This gives a factorization of the top diamond of the diagram above as follows:



Note that  $W_{1-}^{\text{res}} = X_1''$  and  $W_{m-}^{\text{res}} = X_2''$  by Remark 5.2.2, and  $X_i'' \rightarrow X_i$  possess obvious factorizations. Putting these together we functorially obtain a diagram

$$\begin{array}{ccccccc}
X_1'' & & W_1^{\text{torres}} & & W_{m-1}^{\text{torres}} & & X_2'' \\
\downarrow & \searrow & \swarrow & \searrow & \swarrow & \searrow & \downarrow \\
(\text{Fact}_{\text{rs}}) & & (\text{Fact}_{\text{rs}}) & & (\text{Fact}_{\text{rs}}) & & (\text{Fact}_{\text{rs}}) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
X_1 & \rightarrow & W_{1-}^{\text{res}} & \xrightarrow{\varphi_1} & W_{2-}^{\text{res}} & \xrightarrow{\varphi_2} & \dots & \xrightarrow{\varphi_{m-1}} & W_{(m-1)-}^{\text{res}} & \xrightarrow{\varphi_m} & W_{m-}^{\text{res}} & \rightarrow & X_2.
\end{array}$$

Note that  $W_i$  are given by blowing up of functorial ideals on  $X_2$ , and that  $W_{i\pm}^{\text{res}}$  are obtained by blowing up functorial ideals on  $W_i$ , all restricting to the identity on  $U$ . Similarly, the terms appearing in the diagonal arrows are given by blowing up of functorial ideals on  $W_{i\pm}^{\text{res}}$ . By the result of Section 2.3.10 all terms appearing are obtained by blowing up of functorial ideals on  $X_2$  restricting to the unit ideal on  $U$ . In case  $X_i \setminus U$  are normal crossings divisors, we have guarantees that the same holds for  $W_{i\pm}^{\text{res}}$ . It follows that the same holds for all terms in the sequence forming  $W_{i\pm}^{\text{can}} \rightarrow W_{i\pm}^{\text{res}}$  by the properties of canonical principalization, and for the terms in a factorization of  $W_i^{\text{torres}} \rightarrow W_{i\pm}^{\text{can}}$  since these are all nonsingular toroidal schemes. Renaming all these terms  $V_i, i = 1, \dots, l$ , Theorem 1.3.3 follows. ♣

## 6. EXTENDING THE FACTORIZATION TO OTHER CATEGORIES

In this section we use the factorization for schemes to construct an analogous factorization for blowings up of formal schemes, complex and non-archimedean analytic spaces, and stacks. We follow the general outline of the argument in [Tem12, Sections 5.1–5.2], though we decided to elaborate more details related to the relative GAGA issues. In fact, for this construction to work one only needs to have a reasonable comparison theory between algebraic blow ups and their analytifications, but some of these results do not seem to be covered by the literature, especially in the complex analytic case.

Sec:stacks

6.1. **Stacks.** Once functorial factorization for schemes is established it extends to stacks straightforwardly.

6.1.1. *Basic notions.* Our terminology concerning stacks follows that of [Tem12, §5.1]. In particular, by a *stack* we mean an Artin stack  $\mathfrak{X}$  and  $\mathfrak{X}$  is *qe* (respectively, *regular*) if it admits a smooth covering  $W \rightarrow \mathfrak{X}$  with  $W$  a qe (respectively, a regular) scheme. The definition of blowing up along a closed subscheme is compatible with flat morphisms and hence extends to stacks. We define the regular surjective category of blowings up of stacks  $\text{Bl}_{\text{rs}}^{\text{st}}$  and the regular surjective category of weak factorizations of blowings up of stacks  $\text{Fact}_{\text{rs}}^{\text{st}}$  as in definitions 1.3.1 and 1.3.2.

6.1.2. *Factorization for stacks.* We are now in position to extend the factorization to stacks.

Th:stacks

**Theorem 6.1.3.** *There is a functor  $\text{Bl}_{\text{rs}}^{\text{st}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}^{\text{st}}(\text{char} = 0)$  from the regular surjective category of blowings up  $f: \mathfrak{X}' \rightarrow \mathfrak{X}$  in characteristic zero to the regular surjective category of factorizations*

$$\mathfrak{X}' = \mathfrak{X}_0 \dashrightarrow \mathfrak{X}_1 \dashrightarrow \dots \dashrightarrow \mathfrak{X}_{l-1} \dashrightarrow \mathfrak{X}_l = \mathfrak{X},$$

*in characteristic zero such that the composite*

$$\text{Bl}_{\text{rs}}^{\text{st}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}^{\text{st}}(\text{char} = 0) \rightarrow \text{Bl}_{\text{rs}}^{\text{st}}(\text{char} = 0)$$

is the identity. The same holds in positive and mixed characteristics if Hypothetical Statements 2.1.4 and 2.2.4 hold true.

*Proof.* Choose a smooth covering of  $\mathfrak{X}$  by a qe scheme  $W$ . Then  $W$  and  $R = W \times_{\mathfrak{X}} W$  are regular qe schemes and the projections  $p_{1,2}: R \rightrightarrows W$  are surjective and smooth. The pullbacks  $W' \rightarrow W$  and  $R' \rightarrow R$  of  $\mathfrak{X}' \rightarrow \mathfrak{X}$  are objects of  $\text{Bl}_{\text{rs}}$ , hence Theorem 1.3.3 provides their regular factorizations  $(W_{\bullet})$  and  $(R_{\bullet})$ . By the functoriality, these factorizations are compatible with both  $p_1$  and  $p_2$ . Since both pullbacks of the factorization  $(W_{\bullet})$  to  $R$  coincide, flat descent implies that  $(W_{\bullet})$  comes from a factorization  $(\mathfrak{X}_{\bullet})$  of  $\mathfrak{X}' \rightarrow \mathfrak{X}$ .

To see that the factorization  $(\mathfrak{X}_{\bullet})$  is independent of a smooth covering  $W \rightarrow \mathfrak{X}$  we note that any smooth covering  $W' \rightarrow \mathfrak{X}$  that factors through  $W$  induces the same factorization of  $\mathfrak{X}' \rightarrow \mathfrak{X}$ , as follows from the functoriality of factorization with respect to the morphism  $W' \rightarrow W$ .

Finally, assume that  $(\mathfrak{Y}' \rightarrow \mathfrak{Y}) \rightarrow (\mathfrak{X}' \rightarrow \mathfrak{X})$  is a morphism in  $\text{Bl}_{\text{rs}}^{\text{st}}$ . Then there exist smooth coverings by qe schemes  $W \rightarrow \mathfrak{X}$  and  $T \rightarrow \mathfrak{Y}$  such that the morphism  $\mathfrak{Y} \rightarrow \mathfrak{X}$  lifts to a regular surjective morphism  $T \rightarrow W$ . It then follows easily from the functoriality of factorization with respect to  $T \rightarrow W$  that the factorization for stacks we constructed is compatible with  $\mathfrak{Y} \rightarrow \mathfrak{X}$ . Thus, the factorization for stacks is functorial.  $\clubsuit$

## 6.2. Geometric spaces.

6.2.1. *Categories.* We will work with the geometric spaces of the following four classes, that will simply be called *spaces*:

- (1) qe formal schemes as defined in [Tem12, Section 2.4.3],
- (2) semianalytic germs of complex analytic spaces, see Appendix B,
- (3)  $k$ -analytic spaces of Berkovich for a complete non-Archimedean field  $k$ , see [Ber93, Section 1],
- (3') rigid  $k$ -analytic spaces, where  $k$  is as above and non-trivially valued.

To make notation uniform, the category of all such spaces will be denoted  $\mathfrak{Sp}$  in each of the four cases.

**Remark 6.2.2.** (i) The case (3') is added for the sake of completeness. It is essentially included in (3) because the category of qcqs (i.e. quasi-compact and quasi-separated) rigid spaces is equivalent to the category of compact strictly analytic Berkovich spaces, and all our arguments will be "local enough".

(ii) Probably, there exist other contexts where our methods apply, e.g. semialgebraic geometry. We do not explore this direction here, but we will deal with the above four cases in a uniform way that should make it simpler for the interested reader to extend our results to other possible settings.

6.2.3. *Affinoid spaces.* We say that a space  $X$  is *affinoid* if it is of the following type:

- (1)  $X = \text{Spf}(A)$  is affine,
- (2)  $(\mathcal{X}, X)$  is an affinoid germ of a complex analytic space, see Section B.6
- (3)  $X = \mathcal{M}(A)$  is an affinoid  $k$ -analytic space,
- (3')  $X = \text{Sp}(A)$  is an affinoid rigid space over  $k$ .

6.2.4. *Admissible affinoid coverings.* To simplify the discussion we consider only affinoid coverings  $X = \cup_{i \in I} X_i$  of a qcqs space by its affinoid domains. Such a covering is called *admissible* if it possesses a finite refinement. Here is the main property of admissible coverings, which may fail for non-admissible ones (e.g. the covering of a germ  $(\mathcal{X}, X)$  by one-pointed subgerms  $(\mathcal{X}, x)$  with  $x \in X$ ).

**Lem:acyclicity**

**Lemma 6.2.5.** *Assume that  $X = \cup_{i \in I} X_i$  is an admissible covering of an affinoid space. Then for any coherent  $\mathcal{O}_X$ -module  $\mathcal{F}$  the Čech complex*

$$0 \rightarrow \mathcal{F}(X) \rightarrow \prod_i \mathcal{F}(X_i) \rightarrow \prod_{i,j} \mathcal{F}(X_i \cap X_j) \rightarrow \dots$$

*is acyclic.*

*Proof.* For formal schemes this is classical, and for non-archimedean geometry this is Tate's Acyclicity Theorem and its extension to Berkovich spaces. It remains to deal with complex germs. It suffices to deal with the case of finite coverings, and then we can replace the direct products with direct sums. Choosing a small enough representative  $\mathcal{X}$  of  $X$  we can assume that  $\mathcal{X}$  is Hausdorff. Choose families of Stein domains  $V_0 \supset V_1 \dots$  and  $V_{0i} \supset V_{1i} \dots$  for each  $i \in I$  such that  $X = \cap_{n=0}^{\infty} V_n$  and  $X_i = \cap_{n=0}^{\infty} V_{ni}$ . For each  $n \in \mathbb{N}$  the union  $\cup_{i \in I} V_{ni}$  is a neighborhood of  $X$  and hence it contains some  $V_m$ . Let  $m = m(n)$  be the minimal number for which the latter happens. The intersections  $U_{ni} = V_m \cap V_{ni}$  are Stein domains since  $\mathcal{X}$  is Hausdorff, hence  $V_m$  is covered by Stein domains  $U_{ni}$  and we obtain the acyclic Čech complex

$$0 \rightarrow \mathcal{F}(V_m) \rightarrow \oplus_i \mathcal{F}(U_{ni}) \rightarrow \oplus_{i,j} \mathcal{F}(U_{ni} \cap U_{nj}) \rightarrow \dots$$

Since  $\lim_{n \rightarrow \infty} m(n) = \infty$  and  $X_i = \cap_n U_{ni}$ , passing to the limit on  $n$  we obtain the sequence from the formulation of the Lemma. It remains to use that the filtered colimit is an exact functor.  $\clubsuit$

6.2.6. *Regular spaces.* Each category of spaces possesses a natural notion of regular spaces, see [Tem12, Section 5.2.2]. In fact, a space  $X$  is regular if it possesses an admissible affinoid covering  $X = \cup_i X_i$  such that the rings  $A_i = \mathcal{O}_X(X_i)$  are regular. In particular, it follows from Lemma B.6.1 that a germ of analytic space  $(\mathcal{X}, X)$  is regular if and only if  $\mathcal{X}$  is smooth in a neighborhood of  $X$ .

By  $\mathfrak{Sp}_{\text{reg}}$  we denote the full subcategory of  $\mathfrak{Sp}$  consisting of quasi-compact regular objects, and we do not impose any separatedness assumption.

6.2.7. *Smooth and regular morphisms.* Also, the category  $\mathfrak{Sp}$  has a natural notion of smooth morphisms. In cases (1), (2) and (3') this is the classical notion (with the obvious adjustment in (2)) and in (3) this is the notion of quasi-smooth morphisms as defined in [Duc13, Section 4].

In cases (2), (3) and (3') any morphism is of finite type, so we identify the notions of smooth and regular morphisms. Regular morphisms of qc formal schemes were defined in [Tem12, 2.4.12]: a morphism  $f: Y \rightarrow X$  is called *regular* if it admits an open covering of the form  $f_i: \text{Spf}(B_i) \rightarrow \text{Spf}(A_i)$  such that the homomorphisms  $A_i \rightarrow B_i$  are regular.

**Lem:regmor**

**Lemma 6.2.8.** *If  $Y \rightarrow X$  is a regular morphism of affinoid spaces in  $\mathfrak{Sp}$  then the homomorphism  $\mathcal{O}_X(X) \rightarrow \mathcal{O}_Y(Y)$  is regular.*

*Proof.* Case (1) is covered by [Tem12, Lemma 2.4.6]. Case (3), and hence also case (3'), follows from [Duc13, Proposition 4.5.1], [Duc09, Theorem 3.3] and the fact that for any affinoid space  $Z = \mathcal{M}(C)$  the map  $Z \rightarrow \text{Spec}(C)$  is surjective by [Ber93, Proposition 2.1.1]. Case (2) is dealt with similarly using that if  $Z$  is an affinoid germ,  $z \in Z$  and  $f: Z \rightarrow T = \text{Spec}(\mathcal{O}_Z(Z))$  is the natural map then  $f(Z)$  is the set of all closed points and the homomorphism  $\mathcal{O}_{T,f(z)} \rightarrow \mathcal{O}_{Z,z}$  is regular by Lemma B.6.1. ♣

**6.3. Relative GAGA.** Assume that  $X$  is an affinoid space,  $A = \mathcal{O}_X(X)$  and  $\mathcal{X} = \text{Spec } A$ . Relative GAGA relates the theory of  $\mathcal{X}$ -schemes and  $X$ -spaces.

**6.3.1. Analytification functor.** There exists an analytification/formal completion functor from  $\mathcal{X}$ -schemes of finite type to  $X$ -spaces. For uniformity, we will usually call this functor *analytification* and denote  $\mathcal{Y} \mapsto Y = \mathcal{Y}^{\text{an}}$ . It is constructed as follows:

- (i) The analytification of  $\mathbb{A}_{\mathcal{X}}^n$  is  $\mathbb{A}_X^n$ .
- (ii) If  $\mathcal{Y}$  is  $\mathcal{X}$ -affine, say  $\mathcal{Y} = \text{Spec } B$  with  $B = A[t_1, \dots, t_n]/(f_1, \dots, f_m)$ , then  $\mathcal{Y}^{\text{an}}$  is the vanishing locus of  $f_1, \dots, f_m$  in  $\mathbb{A}_X^n$ . It is easily seen to be independent of the  $A$ -presentation of  $B$ .
- (iii) The construction in (ii) is compatible with localizations, so in general one covers  $\mathcal{Y}$  by  $\mathcal{X}$ -affine schemes  $\mathcal{Y}_i$  and glues  $\mathcal{Y}^{\text{an}}$  from  $\mathcal{Y}_i^{\text{an}}$ .

**6.3.2. The analytification map.** There exist natural *analytification maps*  $\pi_{\mathcal{Y}}: \mathcal{Y}^{\text{an}} \rightarrow \mathcal{Y}$  which can be constructed through the steps (i)–(iii), or directly (ii) and (iii). Let us describe them in the affine case  $\mathcal{Y} = \text{Spec } B$ :

- (1) The map is  $\text{Spf } B \hookrightarrow \text{Spec } B$ . It is injective and the image is the set of open prime ideals of  $B$ .
- (2),(3') The map  $\mathcal{Y}^{\text{an}} \rightarrow \mathcal{Y}$  is injective and its image is the set of maximal ideals of  $B$ .
- (3) The map  $\mathcal{Y}^{\text{an}} \rightarrow \mathcal{Y}$  is surjective, see [Ber93, Proposition 2.6.2].

**6.3.3. Sheaves.** The analytification functor also extends to coherent sheaves: for any  $\mathcal{X}$ -scheme  $\mathcal{Y}$  of finite type there exists an analytification functor  $\text{Coh}(\mathcal{Y}) \rightarrow \text{Coh}(\mathcal{Y}^{\text{an}})$  given by  $\mathcal{F}^{\text{an}} = \pi_{\mathcal{Y}}^* \mathcal{F}$ .

Sec:Gagaproperties

**6.3.4. Properties.** For each  $\mathcal{X}$ -proper scheme  $\mathcal{Y}$  the analytification functor  $\text{Coh}(\mathcal{Y}) \xrightarrow{\sim} \text{Coh}(Y)$  is an equivalence of categories. In particular, the analytification functor induces an equivalence between the categories of projective  $\mathcal{X}$ -schemes and  $X$ -spaces. The references are:

- (1) Grothendieck's Existence Theorem, [Gro67, III<sub>1</sub>, 5.1.4].
- (2) Theorem C.1.1 below.
- (3) The analytification was introduced in [Ber93, Section 2.6], and comparison of coherent sheaves can be found in [Poi10, Theorem A.1].
- (3') Köpf's theorem, see [Köp74, Sections 5 and 6] and [Con06, Example 3.2.6].

**6.3.5. Analytification and regularity.** Various properties are respected by analytification, but for our needs we only need to study the situation with regularity.

Prop:gagareg

**Proposition 6.3.6.** *Assume that  $X$  is an affinoid space with  $A = \mathcal{O}_X(X)$ ,  $\mathcal{X} = \text{Spec}(A)$ , and  $\mathcal{Y}$  is an  $\mathcal{X}$ -scheme of finite type with  $Y = \mathcal{Y}^{\text{an}}$ , then*

- (i) If  $\mathcal{Y}$  is regular then  $Y$  is regular.
- (ii) Conversely, assume that  $Y$  is regular, then
  - (a) in cases (2), (3) and (3'),  $\mathcal{Y}$  is regular,
  - (b) in case (1) assume also that  $\mathcal{Y}$  is  $\mathcal{X}$ -proper, then  $\mathcal{Y}$  is regular.

*Proof.* Note that case (3') follows from (3) since a qcqs rigid space can be enhanced to an analytic space, and the regularity is preserved. We will study cases (1), (2) and (3) separately, but let us first make a general remark. The claims (i) and (ii)(a) are local on  $\mathcal{Y}$ , so we can assume that  $\mathcal{Y} = \text{Spec } B$  for a finitely generated  $A$ -algebra  $B$  in these cases.

Case (1). In this case,  $A$  is an  $I$ -adic ring and  $X = \text{Spf } A$ . Since  $A$  is qe,  $B$  is qe and so the  $I$ -adic completion homomorphism  $B \rightarrow \widehat{B}$  is regular. This implies (i) since if  $B$  is regular then  $\widehat{B}$  is regular, and so  $\text{Spf } \widehat{B}$  is regular.

Let us prove (ii). Since  $A$  is  $I$ -adic,  $I$  is contained in the Jacobson radical of  $A$  (see [AM69, Proposition 10.15(iv)]), and so any point of  $\mathcal{X}$  has a specialization in  $\mathcal{X}_s := V(I)$ . By the properness of  $f: \mathcal{Y} \rightarrow \mathcal{X}$ , any point of  $\mathcal{Y}$  has a specialization in  $\mathcal{Y}_s := f^{-1}(\mathcal{X}_s)$ , hence it suffices to prove the following claim: if  $\mathcal{Y}$  is of finite type over  $\mathcal{X}$  and  $Y$  is regular, then  $\mathcal{Y}$  is regular at any point  $y \in \mathcal{Y}_s$ .

The latter claim is local around  $y$ , hence we can assume, again, that  $\mathcal{Y} = \text{Spec } B$ . Let  $m \subset B$  be the ideal corresponding to  $y$ , then the  $m$ -adic completion  $B \rightarrow \widehat{B}_m$  factors through the  $I$ -adic completion  $B \rightarrow \widehat{B}$ , and so  $\widehat{B}_m$  is the completion of  $\widehat{B}$  along  $m\widehat{B}$ . Since  $X$  is qe,  $\widehat{B}$  is qe and so  $\widehat{B} \rightarrow \widehat{B}_m$  is regular. By our assumption  $\widehat{B}$  is regular, hence  $\widehat{B}_m$  is regular too. The homomorphism  $B_m \rightarrow \widehat{B}_m$  is faithfully flat, hence  $B_m$  is regular and we win.

Case (3). In this case,  $A$  is  $k$ -affinoid and  $X = \mathcal{M}(A)$ . Consider a point  $y \in Y$  and set  $\mathbf{y} = \pi_{\mathcal{Y}}(y) \in \mathcal{Y}$ . By [Duc13, (1.3.7.2)],  $\mathcal{Y}$  is regular at  $\mathbf{y}$  if and only if  $Y$  is regular at  $y$ . Since  $\pi_{\mathcal{Y}}$  is surjective this implies that  $\mathcal{Y}$  is regular if and only if  $Y$  is so.

Case (2). If  $y \in Y$  and  $\mathbf{y} = \pi_{\mathcal{Y}}(y)$  then it follows easily from Lemma B.6.1 that the homomorphism  $f_{\mathbf{y}}: \mathcal{O}_{\mathcal{Y}, \mathbf{y}} \rightarrow \mathcal{O}_{Y, y}$  induces an isomorphism of the completions. A local ring is regular if and only if its completion is regular, hence  $\mathcal{O}_{\mathcal{Y}, \mathbf{y}}$  is regular if and only if  $\mathcal{O}_{Y, y}$  is so. Since the image of  $\pi_{\mathcal{Y}}$  contains all closed points, we obtain that  $Y$  is regular if and only if  $\mathcal{Y}$  is regular.  $\clubsuit$

#### 6.4. The factorization theorem.

6.4.1. *Blowings up.* Each of the categories  $\mathfrak{Sp}$  has a natural notion of blowings up  $f: X' \rightarrow X$  along ideals (e.g., see [Tem12, Section 2.4.4] and [Tem12, Section 5.1.2]). In fact,  $Bl_I(X)$  can be described as follows: if  $Y \subset X$  is an affinoid domain,  $\mathcal{Y} = \text{Spec}(\mathcal{O}_X(Y))$  and  $\mathcal{I} \subset \mathcal{O}_{\mathcal{Y}}$  is induced by  $I$ , then the restriction of  $f$  onto  $Y$  is the analytification of the blowing up  $Bl_{\mathcal{I}}(\mathcal{Y}) \rightarrow \mathcal{Y}$ . We will only consider blowings up with nowhere-dense centers.

6.4.2. *Weak factorization.* By a weak factorization of  $X_1 \rightarrow X_2$  we mean a diagram

$$X_1 = V_0 \xrightarrow{-\phi_1} V_1 \xrightarrow{-\phi_2} \dots \xrightarrow{-\phi_{l-1}} V_{l-1} \xrightarrow{-\phi_l} V_l = X_2$$

along with subspaces  $Z_i$  and ideal sheaves  $J_i$  satisfying conditions (1-5) of Section 1.2, where in (2) and (4) the word “scheme” is replaced with “space”. For brevity of notation, such a datum will be denoted  $(V_\bullet, \phi_\bullet, Z_\bullet)$ .

We define the regular surjective category of blowings up  $\text{Bl}_{\text{rs}}^{\text{Sp}}$  in  $\text{Sp}$  and the regular surjective category of weak factorizations  $\text{Fact}_{\text{rs}}^{\text{Sp}}$  on  $\text{Sp}$  as in definitions 1.3.1 and 1.3.2. By definition, these categories are fibred over the category of regular spaces with regular morphisms, and the fibers over a regular space  $X$  will be denoted  $\text{Bl}_{\text{rs}}(X)$  and  $\text{Fact}_{\text{rs}}(X)$ . Thus,  $\text{Bl}_{\text{rs}}(X)$  is the set of blowings up  $X' \rightarrow X$  with regular  $X$  and  $\text{Fact}_{\text{rs}}(X)$  is the set of all regular factorizations of blowings up of  $X$ .

Lem:gagafact

**Lemma 6.4.3.** *Let  $X$  be an affinoid space,  $A = \mathcal{O}_X(X)$  and  $\mathcal{X} = \text{Spec } A$ . Then the analytification functor  $\mathcal{Y} \mapsto \mathcal{Y}^{\text{an}}$  induces bijections  $\text{Bl}_{\text{rs}}(X) \xrightarrow{\sim} \text{Bl}_{\text{rs}}(\mathcal{X})$  and  $\text{Fact}_{\text{rs}}(X) \xrightarrow{\sim} \text{Fact}_{\text{rs}}(\mathcal{X})$ .*

*Proof.* By the relative GAGA, see Section 6.3.4, analytification induces a bijection between the blowings up  $X' \rightarrow X$  and  $\mathcal{X}' \rightarrow \mathcal{X}$ . By Proposition 6.3.6,  $X'$  is regular if and only if  $\mathcal{X}'$  is regular, hence  $\text{Bl}_{\text{rs}}(X) \xrightarrow{\sim} \text{Bl}_{\text{rs}}(\mathcal{X})$ . The second bijection is proved similarly, but this time one also relates regularity of the centers in the factorizations.  $\clubsuit$

6.4.4. *The main theorem.* We are now in position to prove the following analogue of Theorem 1.3.3.

Th:main-C

**Theorem 6.4.5.** *There is a functor  $\text{Bl}_{\text{rs}}^{\text{Sp}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}^{\text{Sp}}(\text{char} = 0)$  from the regular surjective category of blowings up  $f: X' \rightarrow X$  in characteristic zero to the regular surjective category of factorizations*

$$X' = V_0 \dashrightarrow V_1 \dashrightarrow \dots \dashrightarrow V_{l-1} \dashrightarrow V_l = X,$$

*in characteristic zero such that the composite*

$$\text{Bl}_{\text{rs}}^{\text{Sp}}(\text{char} = 0) \rightarrow \text{Fact}_{\text{rs}}^{\text{Sp}}(\text{char} = 0) \rightarrow \text{Bl}_{\text{rs}}^{\text{Sp}}(\text{char} = 0)$$

*is the identity. The same holds in positive and mixed characteristics if Hypothetical Statements 2.1.4 and 2.2.4 hold true.*

*Proof.* First, let us construct a factorization of  $f: X' \rightarrow X$ . Fix an admissible affinoid covering  $X = \cup_{i=1}^n X_i$  and set  $X'_i = X_i \times_X X'$ . The rings  $A_i = \mathcal{O}_X(X_i)$  are qe, see [Tem12, Section 5.2.3], so the scheme  $\mathcal{X} = \coprod_{i=1}^n \mathcal{X}_i$  with  $\mathcal{X}_i = \text{Spec}(A_i)$  is noetherian and qe. Let  $I$  be the ideal defining  $f$  and let  $I_i \subset A_i$  be its restrictions. Consider the blowings up  $F_i: \mathcal{X}'_i \rightarrow \mathcal{X}_i$  defined by  $I_i$ . The analytification of  $F_i$  is the restriction  $f_i$  of  $f$  over  $X_i$  by the relative GAGA, hence  $\mathcal{X}'_i$  is regular by Proposition 6.3.6(ii).

Set  $\mathcal{X}' = \coprod_{i=1}^n \mathcal{X}'_i$  and consider the factorization  $(\mathcal{V}_\bullet, \Phi_\bullet, \mathcal{Z}_\bullet)$  of the blow up  $F: \mathcal{X}' \rightarrow \mathcal{X}$ . For each  $i$ , it induces a factorization  $(\mathcal{V}_{i,\bullet}, \Phi_{i,\bullet}, \mathcal{Z}_{i,\bullet})$  of  $F_i: \mathcal{X}'_i \rightarrow \mathcal{X}_i$  and the analytification of the latter is a factorization of  $f_i: X'_i \rightarrow X_i$  that will be denoted  $(V_{i,\bullet}, \phi_{i,\bullet}, Z_{i,\bullet})$ .

We claim that the latter factorizations glue to a factorization of  $f$ . It suffices to prove that for any  $i, j$  and an affinoid domain  $Y \subset X_i \cap X_j$  the restrictions of  $(V_{i,\bullet}, \phi_{i,\bullet}, Z_{i,\bullet})$  and  $(V_{j,\bullet}, \phi_{j,\bullet}, Z_{j,\bullet})$  onto  $Y$  coincide. Set  $B = \mathcal{O}_X(Y)$  and  $\mathcal{Y} = \text{Spec}(B)$ , and let  $G: \mathcal{Y}' \rightarrow \mathcal{Y}$  be the blowing up along the ideal induced by  $I$ . In particular, the analytification  $g: Y' \rightarrow Y$  of  $G$  is the restriction of  $f$ . The regular

homomorphisms  $A_i \rightarrow B$  and  $A_j \rightarrow B$  induce regular morphisms  $h_i, h_j: \mathcal{Y} \rightarrow \mathcal{X}$  such that  $G$  is the pullback of  $F$  with respect to either of these morphisms. The factorizations of  $G$  induced from  $(\mathcal{V}_\bullet, \Phi_\bullet, \mathcal{Z}_\bullet)$  via  $h_i$  and  $h_j$  coincide by Lemma 6.4.6 below. It remains to note that the factorizations of  $g$  induced from the factorizations of  $f_i$  and  $f_j$  are the analytifications of these factorizations of  $G$ .

We have constructed a factorization of  $f$ . The same argument as was used to glue local factorizations to a global one shows that the construction is independent of the affinoid covering. Finally, compatibility of factorization with a regular morphism  $h: Y \rightarrow X$  is deduced in the same way from Lemma 6.2.8 and compatibility with regular morphisms of factorization for schemes. ♣

The following result is an analogue of [Tem12, Lemma 2.3.1].

Lem:func

**Lemma 6.4.6.** *Assume that  $\mathcal{F}: \text{Bl}_{\text{rs}} \rightarrow \text{Fact}_{\text{rs}}$  is a factorization functor,  $f: X' \rightarrow X$  and  $g: Y' \rightarrow Y$  are two blowings up with regular source and target and  $h_i: Y \rightarrow X$  with  $i = 1, 2$  are two regular morphisms such that  $h_i^*(f) = g$ . Then the pullbacks of  $\mathcal{F}(f)$  to a factorization of  $g$  via  $h_1$  and  $h_2$  coincide.*

*Proof.* Extend  $h_i$  to morphisms  $\phi_i: Y \amalg X \rightarrow X$  so that the map on  $X$  is the identity. Each  $\phi_i$  is a surjective regular morphism, hence the pullback of  $\mathcal{F}(f)$  to  $Y \amalg X$  via  $\phi_i$  coincides with the factorization of the blowing up  $Y' \amalg X' \rightarrow Y \amalg X$ . Restricting the latter onto  $Y$  coincides with  $h_i^*(\mathcal{F}(f))$ . ♣

**Remark 6.4.7.** (i) An analogue of Lemma 6.4.6 holds true in any category  $\mathfrak{Sp}$  and the above proof applies verbatim.

(ii) Although  $h_i^*(\mathcal{F}(f))$  coincide, they can differ from  $\mathcal{F}(g)$  when  $h_i$  are not surjective. See also [Tem12, Remark 2.3.2(ii)].

#### APPENDIX A. CONSTRUCTION OF A BIRATIONAL COBORDISM VIA DEFORMATION TO THE NORMAL CONE

App:cobordism

*Proof of Proposition 4.4.1.* We follow the construction of [AKMW02, Theorem 2.3.1] word for word, except we make it even more explicit and check functoriality.

STEP 1: *cobordism  $B_{\mathcal{O}}$  for trivial blowing up.* We start with

$$B_{\mathcal{O}} = \mathbb{P}_{X_2}^1 = \mathbb{P}(\mathcal{O}_{X_2} \cdot T_0 \oplus \mathcal{O}_{X_2} \cdot T_1) =: \mathbb{P}_{X_2}(E_{\mathcal{O}}),$$

with its projection  $\pi_0: B_{\mathcal{O}} \rightarrow X_2$ . Providing the generators  $T_0$  and  $T_1$  with  $\mathbb{G}_m$ -weights 0 and 1, the scheme  $B_{\mathcal{O}}$  is a birational cobordism for the identity morphism with the trivial ideal (1), with the standard action of  $\mathbb{G}_m$  linearized, except that it does not satisfy Assumption 4.2.2. But that may be achieved after the fact by taking the symmetric square. The construction is clearly functorial.

STEP 2A: *construction of a singular cobordism  $B_I$ .* Assume  $X_1$  is given as the blowing up of the ideal  $I$  on  $X_2$ . We blow up the  $\mathbb{G}_m$ -equivariant ideal  $I^B := I \otimes \mathcal{O}_{B_{\mathcal{O}}} + I_{\{0\}}$  on  $B_{\mathcal{O}}$ , where  $I_{\{0\}}$  is the defining ideal of  $\{0\} \times X_2$ . The ideal is clearly the unit ideal on  $\mathbb{P}_{\mathcal{O}}^1$ . This blowing up gives rise to a  $\mathbb{G}_m$ -scheme  $B_I$  and projective morphism  $\pi_I: B_I \rightarrow B_{\mathcal{O}}$ ; this is evidently functorial in  $\phi$ . The arguments of Section 2.3.10 show that  $\pi^{B_I/X_2} := \pi_0 \circ \pi_I: B_I \rightarrow X_2$  is projective, again in a functorial manner. In particular  $B_I \subset \mathbb{P}(E_I)$  for some functorial  $\mathbb{G}_m$ -sheaf  $E_I$ .

STEP 2B: *coordinates of  $B_I$ .* Let us make the construction of the previous step explicit: write  $F_I = \pi_{0*} I^B(1) = I \cdot U_0 \oplus \mathcal{O}_{X_2} \cdot U_1$  with  $U_0, U_1$  having corresponding



$\mathbb{G}_m$ -weights 0 and 1. Let

$$E_I = F_I \otimes E_{\mathcal{O}} = I \cdot U_0 T_0 \oplus (\mathcal{O}_{X_2} \cdot U_1 T_0 \oplus I \cdot U_0 T_1) \oplus \mathcal{O}_{X_2} \cdot U_1 T_1$$

with corresponding  $\mathbb{G}_m$ -weights 0, 1 and 2. Again it does not satisfy Assumption 4.2.2, but again that may be achieved after the fact by taking the symmetric square.

We have a surjection  $\pi_0^* F_I \rightarrow I^B(1)$  where the first coordinate sends  $f \cdot U_0 \mapsto f T_0$  and the second sends  $U_1 \mapsto T_1$ . We thus have  $\mathbb{G}_m$ -equivariant closed embeddings

$$\begin{aligned} B_I &= Bl_{I^B}(B_{\mathcal{O}}) = Bl_{I^B(1)}(B_{\mathcal{O}}) \subset \\ \mathbb{P}_{B_{\mathcal{O}}}(\pi_0^* F_I) &= \mathbb{P}_{X_2}(F_I) \times_{X_2} B_{\mathcal{O}} = \mathbb{P}_{X_2}(F_I) \times_{X_2} \mathbb{P}_{X_2}(E_{\mathcal{O}}) \subset \\ &\mathbb{P}_{X_2}(F_I \otimes E_{\mathcal{O}}) = \mathbb{P}_{X_2}(E_I), \end{aligned}$$

where  $Bl_{I^B(1)}(B_{\mathcal{O}})$  denotes the blowing up of the fractional ideal  $I^B(1)$  and the last inclusion is the Segre embedding.

We describe  $B_I = \text{Proj}_{X_2} A$  as follows. The algebra

$$A := \bigoplus_d (I^d \cdot T_0^{2d} \oplus I^{d-1} \cdot T_0^{2d-1} T_1 \oplus \cdots \oplus \mathcal{O}_{X_2} \cdot T_0 T_1^{2d-1} \oplus \mathcal{O}_{X_2} \cdot T_1^{2d}),$$

with terms  $I^{d-k} \cdot T_0^j T_1^k$  when  $j > k$  and  $\mathcal{O}_{X_2} \cdot T_0^j T_1^k$  when  $j \leq k$ , is a graded  $\mathbb{G}_m$ -weighted quotient  $\text{Sym}^{\bullet} E_I \twoheadrightarrow A$ , where we set  $U_j = T_j$  and map  $I^{\otimes d} \twoheadrightarrow I^d$ .

We note that  $B_I$  admits an equivariant projection morphisms  $B_I \rightarrow B_{\mathcal{O}} = \mathbb{P}_{X_2}(E_{\mathcal{O}})$  which is an isomorphism away from the divisor  $(T_1^2)$ , and an equivariant projection morphism  $B_I \rightarrow \mathbb{P}_{X_2}(F_I)$ , whose image is the closed subscheme we denote

$$\mathbb{P}_{X_2}(F_I)' := \text{Proj}_{X_2} \bigoplus_{n \geq 0} \left( \bigoplus_{j=0}^n I^j \right).$$

The morphism  $B_I \rightarrow \mathbb{P}_{X_2}(F_I)'$  is an isomorphism away from the zero section  $\text{Proj}_{X_2} \bigoplus_{n \geq 0} \mathcal{O}_{X_2} \subset \mathbb{P}_{X_2}(F_I)'$ , whose complement is the total space  $\text{Spec Sym}((IO_{X_1})^{-1})$  of the invertible sheaf  $IO_{X_1}$  on  $X_1$ .

STEP 2C: *stable and unstable loci for weight 1*. The homogeneous Cartier divisor  $(T_0 T_1)$  is the union of two regular subschemes:  $X_1 = \text{Proj}_{X_2} \bigoplus_{n \geq 0} (I^n \cdot T_0^{2n})$  which is the zero locus of  $(T_0 T_1, T_1^2)$ , and  $X_2 = \text{Proj}_{X_2} \bigoplus_{n \geq 0} (\mathcal{O}_{X_2} \cdot T_1^{2n})$  which is the zero locus of  $(T_0 T_1, I \cdot T_0^2)$ . Since the zero locus of the ‘‘irrelevant ideal’’  $(I \cdot T_0^2, T_0 T_1, T_1^2)$  is empty, these two subschemes are disjoint. In particular each is a regular Cartier divisor. It follows that both  $X_1$  and  $X_2$  lie in the regular locus  $B_I^{\text{reg}}$ , which is open since  $B_I$  is of finite type over the qe scheme  $X_2$ .

We have  $X_1 = B_I \cap \mathbb{P}_{X_2}((E_I)_0)$  and  $X_2 = B_I \cap \mathbb{P}_{X_2}((E_I)_2)$ , where the indices 0 and 2 denote the components with given  $\mathbb{G}_m$ -weight (the variable  $a$  in Section 4.2). Their union  $(T_0 T_1)$  is the unstable locus  $(B_I)_1^{\text{un}}$ . The complement is affine, explicitly

$$\begin{aligned} (B_I)_1^{\text{sst}} &= \text{Spec}_{X_2} A[(T_0 T_1)^{-1}]_{\text{degree}=0} \\ &= \text{Spec}_{X_2} \left( \cdots \oplus I^2 \left( \frac{T_0}{T_1} \right)^2 \oplus I \left( \frac{T_0}{T_1} \right) \oplus \mathcal{O}_{X_2} \oplus \mathcal{O}_{X_2} \left( \frac{T_1}{T_0} \right) \oplus \mathcal{O}_{X_2} \left( \frac{T_1}{T_0} \right)^2 \oplus \cdots \right). \end{aligned}$$

This scheme is in general singular, but the quotient is simpler:

$$(B_I)_1^{\text{sst}} // \mathbb{G}_m = \text{Spec}_{X_2} \mathcal{O}_{X_2} = X_2.$$

STEP 2D: *stable and unstable loci for weight 2*. The projective Cartier divisor  $(T_1^2)$  can be identified as

$$\begin{aligned} (B_I)_2^{\text{un}} &= \mathbb{P}_{X_2}(I \cdot T_0^2) \cup \mathbb{P}_{Z(I)}(I/I^2 \cdot T_0^2 \oplus \mathcal{O} \cdot T_0 T_1) \\ &= X_1 \quad \cup \quad C(Z(I)), \end{aligned}$$

where  $C(Z(I))$  is the normal cone. The complement is again affine, of the form

$$\begin{aligned} (B_I)_2^{\text{sst}} &= \text{Spec}_{X_2} A[T_1^{-1}]_{\text{degree}=0} \\ &= \text{Spec}_{X_2} \left( \cdots \oplus \mathcal{O}_{X_2} \left( \frac{T_0}{T_1} \right)^2 \oplus \mathcal{O}_{X_2} \left( \frac{T_0}{T_1} \right) \oplus \mathcal{O}_{X_2} \right) = \mathbb{A}_{X_2}^1. \end{aligned}$$

Thus,

$$(B_I)_2^{\text{sst}} // \mathbb{G}_m = \text{Spec}_{X_2} \mathcal{O}_{X_2} = X_2$$

and the morphism  $(B_I)_2^{\text{sst}} \rightarrow X_2$  is smooth. Another way to see this is to notice that the map  $B_I \rightarrow B_{\mathcal{O}}$  restricts to an open embedding on  $(B_I)_2^{\text{sst}}$ , and the image is the complement of  $\{0\} \times X_2$ .

STEP 2E: *stable and unstable loci for weight 0*. The projective zero locus of  $(I \cdot T_0)^2$  can be identified as

$$(B_I)_0^{\text{un}} = \mathbb{P}_{X_2}(\mathcal{O}_{X_2} \cdot T_1^2) \cup \mathbb{P}_{Z(I)}(\mathcal{O}_{X_2} \cdot T_0 T_1 \oplus \mathcal{O}_{X_2} \cdot T_1^2) = X_2 \cup \mathbb{P}_{Z(I)}^1.$$

The complement is not necessarily affine, as  $I$  is not necessarily principal. However, recalling the sheaf  $F_I$  from STEP 2B, the morphism  $(B_I)_0^{\text{sst}} \rightarrow \mathbb{P}_{X_2}(F_I)$  is an open embedding, whose image is the complement of the zero section. So  $(B_I)_0^{\text{sst}}$  is the total space of the invertible sheaf  $I\mathcal{O}_{X_1}$  on  $X_1$ . Thus,  $(B_I)_0^{\text{sst}} // \mathbb{G}_m = X_1$  and the morphism  $(B_I)_0^{\text{sst}} \rightarrow X_1$  is smooth.

STEP 3A: *resolving  $B_I$  and principalizing the preimage of  $D_2$* . We apply canonical resolution of singularities to  $B_I$ , which is a functorial  $\mathbb{G}_m$ -equivariant sequence of blowings up  $B^{\text{reg}} \rightarrow B_I$ : see Section 2.1.2, where in positive and mixed characteristic we may use Hypothetical Statement 2.1.4 since  $\dim B = \dim X_2 + 1$ . Next, we apply canonical principalization to the preimage of  $D_2$  in  $B^{\text{reg}}$ , obtaining a functorial  $\mathbb{G}_m$ -equivariant blowing up sequence  $B \rightarrow B^{\text{reg}}$  such that  $B$  is regular and the preimage  $D \subset B$  of  $D_2$  is an ordered simple normal crossings divisor: see Section 2.2, where in positive and mixed characteristic we use Hypothetical Statement 2.2.1. We obtain a sequence of blowings up  $B_{\mathcal{O}} \leftarrow B_I \cdots \leftarrow B_k = B^{\text{reg}} \leftarrow B$  with ideals  $A_i$  supported on the singular locus, which is included in the preimage of  $\mathbb{P}_{X_2}((E_I)_1) = \mathbb{P}_{X_2}(\mathcal{O}_{X_2} \cdot U_1 T_0 \oplus I \cdot U_0 T_1)$ , and the preimages of  $D_2$ .

STEP 3B: *embedding*. By the arguments of Section 2.3.10, the blowing up sequence  $B \rightarrow B_{\mathcal{O}}$  is functorially a single blowing up of an ideal  $J$ . Write  $\tilde{J} = J\mathcal{O}_{B_I}$  so that  $B = Bl_{\tilde{J}} B_I$ . There is a functorially defined integer  $d$  such that  $\tilde{J}(d)$  is globally generated on  $B_I$  relative to  $X_2$ . Using [Har77, II.7.10(b)] we have an equivariant embedding of  $B$  inside

$$\mathbb{P}_{X_2}(\tilde{E}) := \mathbb{P}_{X_2} \left( \pi_*^{B_I/X_2} \tilde{J}(d) \right).$$

Since  $J$  is the unit ideal on  $(B_I)_0^{\text{sst}}$  and  $(B_I)_2^{\text{sst}}$ , we have that  $\pi_*^{B_I/X_2} \tilde{J}(d) \hookrightarrow \text{Sym}^d(E_I)$  is an isomorphism on the components of  $\mathbb{G}_m$ -weight 0 or  $2d$ . Since

$\text{Sym}^d(E_I)$  has weights  $0, \dots, 2d$  which survive in the ring of  $B_I$  it follows that  $a_{\min}(B) = 0, a_{\max}(B) = 2d$ . Inspecting the description of unstable loci in Section 4.1, Equation (1) we note that  $B_0^{\text{sst}} = B \times_{B_I} (B_I)_0^{\text{sst}}$  and  $B_{2d}^{\text{sst}} = B \times_{B_I} (B_I)_2^{\text{sst}}$ .<sup>5</sup> ←5

STEP 3C:  $B$  is a cobordism for  $\phi$  that respects  $U$ . We have shown in steps 2d and 2e that the morphisms  $q_2: (B_I)_2^{\text{sst}} \rightarrow X_2$  and  $q_1: (B_I)_0^{\text{sst}} \rightarrow X_1$  are smooth. Functoriality of resolution and principalization with respect to  $q_i$  implies that, once restricted to  $(B_I)_2^{\text{sst}}$ , respectively  $(B_I)_0^{\text{sst}}$ , the morphism  $B^{\text{reg}} \rightarrow B_I$  is an isomorphism and the morphism  $B \rightarrow B_I$  is the pullback of the principalization  $X'_2 \rightarrow X_2$  of  $D_2$ , respectively  $X'_1 \rightarrow X_1$  of  $D_1$ . It follows that  $B \times_{B_I} (B_I)_2^{\text{sst}} \cong \mathbb{G}_m = X'_2$  and  $B \times_{B_I} (B_I)_0^{\text{sst}} \cong \mathbb{G}_m = X'_1$  and hence  $B$  is a cobordism for  $\phi$ . Also, we note that  $B \cap \mathbb{P}(\tilde{E}_0) = X'_1$  and  $B \cap \mathbb{P}(\tilde{E}_{2d}) = X'_2$ , so Assumption 4.2.3 applies.

To show that  $B$  is compatible with  $U$  it suffices to show that all blowings up in the sequence  $B \rightarrow B_{\mathcal{O}}$  are trivial over  $\mathbb{P}_U^1$ . This is so for the blowing up  $B_I \rightarrow \mathbb{P}_{X_2}^1$  because  $I + I_{\{0\}}$  is the unit ideal on  $\mathbb{P}_U^1$ . This is so for the blowing up sequence  $B^{\text{reg}} \rightarrow B_I$  because  $\mathbb{P}_U^1$  is regular, and this is so the blowing up sequence  $B \rightarrow B^{\text{reg}}$  because  $\mathbb{P}_U^1$  is disjoint from the preimage of  $D_2$ .



App:germs

APPENDIX B. GERMS OF COMPLEX ANALYTIC SPACES

In this section we use germs to extend the category of complex analytic spaces to include certain Stein compacts. This will be used later to establish a tight connection between the scheme theory and complex analytic geometry. In particular, this is needed to develop a relative GAGA theory.

**B.1. Semianalytic sets.** We follow the setup of Frisch [Fri67]. A subset  $X$  of an analytic space  $\mathcal{X}$  is called *semianalytic* if its local germs belong to the minimal class of germs, stable under finite unions and complements, generated by inequalities of the form  $f(x) < 0$  for real analytic  $f$ , see [Fri67, p. 120]. It is called a *Stein* if  $X$  has a fundamental system of neighborhood of Stein subspaces of  $\mathcal{X}$ , see [Fri67, p. 123].

**B.2. The category of germs.** A *germ of a complex analytic space* (or, simply, a germ) is a pair  $(\mathcal{X}, X)$  consisting of an analytic space  $\mathcal{X}$  and a semianalytic subset  $X \subset \mathcal{X}$ . We call  $X$  the *support* of  $(\mathcal{X}, X)$  and we call  $\mathcal{X}$  a *representative* of  $(\mathcal{X}, X)$ . Sometimes, we will use the shorter notation  $X = (\mathcal{X}, X)$ .

A morphism  $\phi: (\mathcal{X}, X) \rightarrow (\mathcal{Y}, Y)$  consists of a neighborhood  $\mathcal{X}'$  of  $X$  and an analytic map  $f: \mathcal{X}' \rightarrow \mathcal{Y}$  taking  $X$  to  $Y$ . We say that  $f$  is a representative of  $\phi$ . Note that a morphism  $(\mathcal{X}, X) \rightarrow (\mathcal{Y}, Y)$  is an isomorphism if it induces a bijection of  $X$  and  $Y$  and an isomorphism of their neighborhoods.

We identify an analytic space  $X$  with the germ  $(X, X)$ . In particular, the category of analytic spaces becomes a full subcategory of the category of germs.

**B.3. The structure sheaf.** Given a germ  $(\mathcal{X}, X)$  we provide its support with the *structure sheaf*  $\mathcal{O}_X := \mathcal{O}_{\mathcal{X}}|_X = i^* \mathcal{O}_{\mathcal{X}}$ , where  $i: X \hookrightarrow \mathcal{X}$  is the embedding. In particular, we obtain a functor  $\mathcal{F}: (\mathcal{X}, X) \mapsto (X, \mathcal{O}_X)$  from the category of germs to the category of locally ringed spaces.

<sup>5</sup>(Dan) One could explain more - do you think we need to?

**Remark B.3.1.** We do not aim to develop a complete theory of semianalytic germs, so we do not study the natural question whether  $\mathcal{F}$  is fully faithful.

**B.4. Closed polydiscs and convergent power series.** Consider an analytic affine space  $\mathcal{X} = \mathbb{A}_{\mathbb{C}}^n$  with coordinates  $t_1, \dots, t_n$ . For any tuple  $r$  of numbers  $r_1, \dots, r_n \in [0, \infty)$ , by the closed polydisc  $D = D_r$  of radius  $r$  we mean the subset of  $\mathcal{X}$  given by the inequalities  $|t_i| \leq r_i$ . Note that  $r_i$  can be zero. By  $\mathbb{C}\{t_1, \dots, t_n\}_r^\dagger$  we denote the ring of overconvergent series in  $t_1, \dots, t_n$  of radius  $r$ . It is a noetherian regular excellent ring of dimension  $n$ , see [Mat80, Theorem 102].

Lem:polydisc

**Lemma B.4.1.** *Let  $D = D_r \subset \mathcal{X} = \mathbb{A}_{\mathbb{C}}^n$  be a polydisc and  $A = \mathcal{O}_{\mathcal{X}}(D) = \Gamma(\mathcal{O}_D)$ . Then,*

(i)  $\mathbb{C}\{t_1, \dots, t_n\}_r^\dagger \xrightarrow{\sim} A$ .

(ii)  $\Gamma(D, \cdot)$  induces an equivalence between the categories of coherent  $\mathcal{O}_D$ -modules and finitely generated  $A$ -modules, and higher cohomology of coherent  $\mathcal{O}_D$ -modules vanish.

(iii) For any  $a \in D$  the ideal  $m_a = (t_1 - a_1, \dots, t_n - a_n) \subset A$  is maximal, and any maximal ideal of  $A$  is of this form.

(iv) The completion of  $A$  along  $m_a$  is  $\mathbb{C}[[t_1 - a_1, \dots, t_n - a_n]]$ .

*Proof.* The first claim is a classical result of analysis of several complex variables. Assertion (ii) follows from the fact that  $D$  is the intersection of open polydiscs containing it, and the latter are Stein spaces. Assertion (iv) follows easily from (iii), so we will only prove (iii).

For any  $f \in A$  the quotient

$$g_1 = (f(t_1, \dots, t_n) - f(a_1, t_2, \dots, t_n)) / (t_1 - a_1)$$

lies in  $A$ , so  $f = (t_1 - a_1)g_1 + f_1(t_2, \dots, t_n)$  with  $f_1 = f(a_1, t_2, \dots, t_n)$ . Applying the same argument to  $t_2$  and  $f_1$ , etc., we will obtain in the end a representation  $f = f(a_1, \dots, a_n) + \sum_{i=1}^n (t_i - a_i)g_i$ . In particular,  $A/m_a = \mathbb{C}$  and hence  $m_a$  is maximal.

Conversely, assume that  $m \subset A$  is maximal. The norm  $\|f\| = \max_{x \in D} |f(x)|$  on  $A$  induces a norm on the field  $\kappa = A/m$ , hence the completion  $K = \hat{\kappa}$  is a Banach  $\mathbb{C}$ -field. Thus,  $K = \mathbb{C}$  by Gel'fand-Mazur theorem, and we obtain that  $t_i - a_i \in m$  for some  $a_i \in \mathbb{C}$ . Finally,  $|a_i| \leq r_i$  as otherwise  $t_i - a_i \in A^\times$ . ♣

**B.5. Classes of morphisms.** Let  $\phi: (\mathcal{Y}, Y) \rightarrow (\mathcal{X}, X)$  be a morphism of germs. We say that  $\phi$  is *without boundary* if there exists a representative  $f: \mathcal{Y}' \rightarrow \mathcal{X}$  such that  $Y = f^{-1}(X)$ . Let  $P$  be one of the following properties: smooth, open immersion, closed immersion. We say that  $\phi$  is  $P$  if it is without boundary and has a representative which is  $P$ . We say that  $\phi$  is an *embedding of a subdomain* (resp. *quasi-smooth*) if it possesses a representative which is an open immersion (resp. smooth).

**Remark B.5.1.** The above terminology is chosen to match its non-archimedean analogue as much as possible.

Sec:affgerm

**B.6. Affinoid germs.** A germ  $X$  is called *affinoid* if it admits a closed immersion into a germ of the form  $(\mathbb{C}^n, D)$  where  $D$  is a closed polydisc. Such a germ is controlled by the ring  $\mathcal{O}_X(X)$  very tightly.

Lem:affgerm

**Lemma B.6.1.** *Assume that  $X$  is an affinoid germ and let  $A = \mathcal{O}_X(X)$  and  $f: (X, \mathcal{O}_X) \rightarrow Y = \text{Spec}(A)$  the corresponding map of locally ringed spaces. Then,*

(i)  *$A$  is a quotient of a ring  $\mathbb{C}\{t_1, \dots, t_n\}_r^\dagger$ ; in particular it is an excellent noetherian ring.*

(ii)  *$\Gamma(X, \cdot)$  induces an equivalence between the categories of coherent  $\mathcal{O}_X$ -modules and finitely generated  $A$ -modules, and higher cohomology of coherent  $\mathcal{O}_X$ -modules vanish.*

(iii)  *$f$  establishes a bijection between  $X$  and the closed points of  $Y$ .*

(iv) *For any point  $x \in X$  with  $y = f(x)$  the homomorphism  $\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$  is regular and its completion  $\widehat{\mathcal{O}}_{Y,y} \rightarrow \widehat{\mathcal{O}}_{X,x}$  is an isomorphism.*

*Proof.* In the case of a closed polydisc the assertion was proved in Lemma B.4.1. In general, we fix a closed embedding  $i: X \hookrightarrow D$  into a closed polydisc. So,  $\mathcal{O}_X$  becomes a coherent  $\mathcal{O}_D$ -algebra such that the homomorphism  $\phi: \mathcal{O}_D \rightarrow \mathcal{O}_X$  is surjective, and then all assertions except the first half of (iv) follow easily from the case of a polydisc. For example,  $\Gamma(X, \mathcal{O}_X)$  is a quotient of  $\Gamma(D, \mathcal{O}_D)$  since  $H^1(D, \text{Ker}\phi) = 0$ , thereby proving (i).

The only new assertion is that  $\phi: \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$  is regular. This follows from the facts that  $\widehat{\phi}$  is an isomorphisms and the local ring  $\mathcal{O}_{Y,y}$  is excellent (since it is a localization of the excellent ring  $A$ ). ♣

APPENDIX C. THE COMPLEX RELATIVE GAGA THEOREM

Sec:relGAGA

**C.1. Statement of the theorem.** Let  $(\mathcal{X}, X)$  be an affinoid germ as in Appendix B with ring of global analytic functions  $A$ , and  $r \geq 0$  an integer. Set  $\mathbb{P}_X^r = \mathbb{C}\mathbb{P}^r \times X$  and endow it with a locally ringed space structure using the sheaf  $\mathcal{O}_{\mathbb{P}_X^r} = \mathcal{O}_{\mathbb{P}^r} |_{\mathbb{P}_X^r}$ . We have a germ  $(\mathbb{P}_X^r, \mathbb{P}_X^r)$  and a morphism of locally ringed spaces  $h: \mathbb{P}_X^r \rightarrow \mathbb{P}_A^r$ . The aim of this appendix is to prove the following extension of Lemma B.6.1:

Th:GAGA

**Theorem C.1.1** (Serre's Théorème 3). *Let  $(\mathcal{X}, X)$  be an affinoid germ with ring of global analytic functions  $A$ , and  $r \geq 0$  an integer. Then the pullback functor  $h^*: \text{Coh}(\mathbb{P}_A^r) \rightarrow \text{Coh}(\mathbb{P}_X^r)$  is an equivalence which induces isomorphisms on cohomology groups.*

Since  $(\mathcal{X}, X)$  is closed in  $(\mathbb{C}^n, D)$  it suffices to consider the case  $(\mathcal{X}, X) = (\mathbb{C}^n, D)$ . So from now on we make this assumption, and write  $A$  for the ring of holomorphic functions on  $X = D$ .

We follow the steps of Serre's original proof [Ser56, §3] in some detail, to alleviate our skepticism that this generalization might actually work. See also [Ked09], which sketches Serre's proof. One difficulty is that we do not know if  $D \times \mathbb{C}^r$  is Stein in the sense of [Fri67] or [GR04]. The problem is that if  $\{D_i\}$  are the open polydiscs containing  $D$  then  $\{D_i \times \mathbb{C}^r\}$  do not form a *fundamental* family of neighborhoods of  $D \times \mathbb{C}^r$ , while functions on  $D \times \mathbb{C}^r$  are only guaranteed to extend to some member of a fundamental family of neighborhoods. This is circumvented in Lemma C.2.2, which is the only point where we differ from the original arguments.

C.2. Cohomology.

**Proposition C.2.1** (Serre's Théorème 1). *Let  $\mathcal{F}$  be a coherent sheaf on  $\mathbb{P}_A^r$ . The homomorphism  $h^*: H^i(\mathbb{P}_A^r, \mathcal{F}) \rightarrow H^i(\mathbb{P}_D^r, h^*\mathcal{F})$  is an isomorphism.*

Lem:GAGA-basic-sheaves

**Lemma C.2.2.** (1) We have  $H^i(\mathbb{P}_A^r, \mathcal{F}) = H^i(\mathbb{P}_D^r, h^*\mathcal{F}) = 0$  for  $i > r$  and all  $\mathcal{F}$ .  
 (2) The proposition holds for  $\mathcal{F} = \mathcal{O}_{\mathbb{P}_A^r}$  for all  $r \geq 0$ .

*Proof.* (1) For  $H^i(\mathbb{P}_A^r, \mathcal{F}) = 0$  use the standard Čech covering of  $\mathbb{P}_A^r$ , which has only  $r + 1$  elements. We need to show  $H^i(\mathbb{P}_D^r, h^*\mathcal{F}) = 0$ .

On the analytic side we mimic the standard argument for vanishing using Čech cocycles of a covering by closed polydiscs instead of affine spaces. Let  $h^*\mathcal{F} \rightarrow S^\bullet$  be the standard flabby resolution of  $h^*\mathcal{F}$  by discontinuous sections, so  $H^i(Y, h^*\mathcal{F}|_Y) = H^i(\Gamma(Y, S^\bullet))$  for any subset  $Y \subset \mathbb{P}_D^r$ . Let  $\mathbb{C}^r \simeq U_i \subset \mathbb{C}\mathbb{P}^r$  be the standard open sets and let  $D_i \subset U_i$  be the standard closed polydisc of fixed radius  $> 1$ . Set  $X_i = D \times D_i \subset \mathbb{P}_D^r$  and for each subset  $I \subset \{0, \dots, n\}$  let  $X_I = \bigcap_{i \in I} X_i$ . Then  $X_I$  are complex affinoids for  $I \neq \emptyset$ , hence  $H^i(X_I, h^*\mathcal{F}|_{X_I}) = 0 = H^i(\Gamma(X_I, S^\bullet))$  for  $i > 0$  and  $I \neq \emptyset$ .

On the other hand

$$\mathcal{C}^\bullet(\{X_i\}, S^j) = \left[ \bigoplus_{|I|=1} S^j_{X_I} \rightarrow \bigoplus_{|I|=2} S^j_{X_I} \rightarrow \dots \right]$$

is a flabby resolution of  $S^j$  so  $H^0(\Gamma(\mathbb{P}_D^r, \mathcal{C}^\bullet(\{X_i\}, S^j))) = \Gamma(\mathbb{P}_D^r, S^j)$  and for  $i > 0$  we have  $H^i(\Gamma(\mathbb{P}_D^r, \mathcal{C}^\bullet(\{X_i\}, S^j))) = 0$ .

Consider the double complex  $C^{p,q} = \bigoplus_{|I|=p} \Gamma(X_I, S^q)$  and its two edges  $\Gamma(\mathbb{P}^r, S^\bullet)$  and  $\check{C}^p = \bigoplus_{|I|=p} \Gamma(X_I, h^*\mathcal{F})$ . We obtain that

$$H^i(\mathbb{P}_D^r, h^*\mathcal{F}) = H^i(\Gamma(\mathbb{P}^r, S^\bullet)) = \mathbb{H}^i(C^{\bullet,\bullet}) = H^i(\check{C}^\bullet).$$

The latter is trivial in degrees  $> r$ .

(2) We have that  $\Gamma(\mathcal{O}_{\mathbb{P}_A^r}) = A$  and  $H^i(\mathcal{O}_{\mathbb{P}_A^r}) = 0$  for  $i > 0$  by [Har77, Theorem III.5.1]. It suffices to show that  $\pi_*\mathcal{O}_{\mathbb{P}_D^r} = \mathcal{O}_D$  and  $R^i\pi_*\mathcal{O}_{\mathbb{P}_D^r} = 0$  for  $i > 0$  where  $\pi: \mathbb{P}_D^r \rightarrow D$  is the projection, since  $D$  is Stein. For this note that  $\mathcal{O}_{\mathbb{P}_D^r} = j_r^{-1}\mathcal{O}_{\mathbb{P}_{\mathbb{C}\mathbb{P}^n}^r}$ , where  $j_r: \mathbb{P}_D^r \rightarrow \mathbb{P}_{\mathbb{C}\mathbb{P}^n}^r$  is the inclusion:

$$\begin{array}{ccc} \mathbb{P}_D^r & \xrightarrow{j_r} & \mathbb{P}_{\mathbb{C}\mathbb{P}^n}^r \\ \pi \downarrow & & \downarrow \varpi \\ D & \xrightarrow{j_0} & \mathbb{C}\mathbb{P}^n. \end{array}$$

By the topological proper push-forward theorem [Ive86, Corollary VII.1.5] we have

$$R^i\pi_*\mathcal{O}_{\mathbb{P}_D^r} = j_0^{-1}R^i\varpi_*\mathcal{O}_{\mathbb{P}_{\mathbb{C}\mathbb{P}^n}^r},$$

and the result follows from Serre's original GAGA theorems.  $\clubsuit$

allrlem

**Lemma C.2.3.** The proposition holds for  $\mathcal{F} = \mathcal{O}_{\mathbb{P}_A^r}(n)$  for all  $r \geq 0$  and all integers  $n$ .

*Proof.* Induction identical to [Ser56, section 13 Lemme 5]: the result holds for  $r = 0$  since  $D$  is Stein. Supposing it holds for  $r - 1$  and all  $n$ , we have the exact sequence  $0 \rightarrow \mathcal{O}_{\mathbb{P}_D^r}(n-1) \rightarrow \mathcal{O}_{\mathbb{P}_D^r}(n) \rightarrow \mathcal{O}_{\mathbb{P}_D^{r-1}}(n) \rightarrow 0$  and the corresponding sequence for

$\mathbb{P}_A^r$ . We obtain a canonical homomorphism of long exact sequences

$$\begin{array}{ccccccc} H^{i-1}(\mathbb{P}_A^{r-1}, \mathcal{O}(n)) & \longrightarrow & H^i(\mathbb{P}_A^r, \mathcal{O}(n-1)) & \longrightarrow & H^i(\mathbb{P}_A^r, \mathcal{O}(n)) & \longrightarrow & H^i(\mathbb{P}_A^{r-1}, \mathcal{O}(n)) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^{i-1}(\mathbb{P}_D^{r-1}, \mathcal{O}(n)) & \longrightarrow & H^i(\mathbb{P}_D^r, \mathcal{O}(n-1)) & \longrightarrow & H^i(\mathbb{P}_D^r, \mathcal{O}(n)) & \longrightarrow & H^i(\mathbb{P}_D^{r-1}, \mathcal{O}(n)). \end{array}$$

The vertical arrows on the right and left are isomorphisms by the inductive assumption. It follows that the result holds for  $r$  and  $\mathcal{O}(n-1)$  if and only if it holds for  $\mathcal{O}(n)$ . Since we have proven that it holds for  $\mathcal{O}$ , it holds for all  $n$ .  $\clubsuit$

*Proof of the proposition.* The proof is identical to Serre's Théorème 1. We apply descending induction on  $i$  for all coherent  $\mathbb{P}_A^r$  modules  $\mathcal{F}$ . The case of  $i > r$  is proved by the lemma. Since  $\mathcal{F}$  is coherent there is an epimorphism  $\mathcal{E} \rightarrow \mathcal{F}$  with  $\mathcal{E} = \bigoplus_{i=1}^n \mathcal{O}_{\mathbb{P}_A^r}(-k_i)$ . Denoting by  $\mathcal{G}$  the kernel,  $\mathcal{G}$  is coherent and we have a short exact sequence

$$0 \rightarrow \mathcal{G} \rightarrow \mathcal{E} \rightarrow \mathcal{F} \rightarrow 0.$$

Since the map  $h$  is flat we have an exact sequence

$$0 \rightarrow h^*\mathcal{G} \rightarrow h^*\mathcal{E} \rightarrow h^*\mathcal{F} \rightarrow 0.$$

In the commutative diagram of cohomologies with exact rows

$$\begin{array}{ccccccc} H^i(\mathbb{P}_A^r, \mathcal{E}) & \longrightarrow & H^i(\mathbb{P}_A^r, \mathcal{F}) & \longrightarrow & H^{i+1}(\mathbb{P}_A^r, \mathcal{G}) & \longrightarrow & H^{i+1}(\mathbb{P}_A^r, \mathcal{E}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^i(\mathbb{P}_D^r, h^*\mathcal{E}) & \longrightarrow & H^i(\mathbb{P}_D^r, h^*\mathcal{F}) & \longrightarrow & H^{i+1}(\mathbb{P}_D^r, h^*\mathcal{G}) & \longrightarrow & H^{i+1}(\mathbb{P}_D^r, h^*\mathcal{E}) \end{array}$$

the vertical arrows on the left and right are isomorphisms by Lemma C.2.3. By the induction hypothesis  $H^{i+1}(\mathbb{P}_A^r, \mathcal{G}) \rightarrow H^{i+1}(\mathbb{P}_D^r, h^*\mathcal{G})$  is an isomorphism as well. By the five lemma the result holds for  $H^i(\mathbb{P}_A^r, \mathcal{F}) \rightarrow H^i(\mathbb{P}_D^r, h^*\mathcal{F})$  as required.  $\clubsuit$

### C.3. Homomorphisms.

**Proposition C.3.1** (Serre's Théorème 2). *For any coherent  $\mathbb{P}_A^r$ -modules  $\mathcal{F}, \mathcal{G}$  the natural homomorphism*

$$\mathrm{Hom}_{\mathbb{P}_A^r}(\mathcal{F}, \mathcal{G}) \rightarrow \mathrm{Hom}_{\mathbb{P}_D^r}(h^*\mathcal{F}, h^*\mathcal{G})$$

*is an isomorphism. In particular the functor  $h^*$  is fully faithful.*

**Lemma C.3.2.** *The sheaf homomorphism*

$$h^*\mathcal{H}om_{\mathbb{P}_A^r}(\mathcal{F}, \mathcal{G}) \rightarrow \mathcal{H}om_{\mathbb{P}_D^r}(h^*\mathcal{F}, h^*\mathcal{G})$$

*is an isomorphism.*

*Proof.* This follows since  $\mathcal{O}_{\mathbb{P}_D^r}$  is a flat  $\mathcal{O}_{\mathbb{P}_A^r}$ -module. Indeed, for a closed point  $x \in \mathbb{P}_D^r$  corresponding to a point  $x' = h(x) \in \mathbb{P}_A^r$  we have

$$\begin{aligned} (h^*\mathcal{H}om_{\mathbb{P}_A^r}(\mathcal{F}, \mathcal{G}))_x &= \mathrm{Hom}_{\mathcal{O}_{x'}}(\mathcal{F}_{x'}, \mathcal{G}_{x'}) \otimes_{\mathcal{O}_{x'}} \mathcal{O}_x \\ &= \mathrm{Hom}_{\mathcal{O}_x}(\mathcal{F}_{x'} \otimes_{\mathcal{O}_{x'}} \mathcal{O}_x, \mathcal{G}_{x'} \otimes_{\mathcal{O}_{x'}} \mathcal{O}_x) \\ &= \mathcal{H}om_{\mathbb{P}_D^r}(h^*\mathcal{F}, h^*\mathcal{G})_x. \end{aligned}$$

$\clubsuit$

*Proof of the proposition.* By Serre's Théorème 1,  $h^*$  preserves cohomology of coherent sheaves. Taking  $H^0$  in the lemma the result follows. ♣

**C.4. The equivalence.** It remains to show:

**Proposition C.4.1.** *The functor  $h^*$  is essentially surjective.*

*Proof.* This is an inductive argument on  $r$  identical to Serre's Théorème 3 which we repeat below. The case  $r = 0$  follows from Lemma B.6.1. Assume the result is known for  $r - 1$  and let  $\mathcal{F}$  be a coherent sheaf on  $\mathbb{P}_D^r$ . By Lemma C.4.2 below there is an epimorphism  $\phi : \mathcal{O}(-n_0)^{k_0} \rightarrow \mathcal{F}$ , and applying this again to  $\text{Ker}(\phi)$  we get a resolution  $\mathcal{O}(-n_1)^{k_1} \xrightarrow{\psi} \mathcal{O}(-n_0)^{k_0} \rightarrow \mathcal{F} \rightarrow 0$ . By Serre's Théorème 2 the homomorphism  $\psi$  is the analytification of an algebraic sheaf homomorphism  $\psi'$ , so the cokernel  $\mathcal{F}$  of  $\psi$  is also the analytification of the cokernel of  $\psi'$ . ♣

Lem:global-generation

**Lemma C.4.2.** *Assume the proposition holds for  $r - 1$ . Then for any coherent sheaf  $\mathcal{F}$  on  $\mathbb{P}_D^r$  there is  $n_0$  so that  $\mathcal{F}(n)$  is globally generated whenever  $n > n_0$ .*

*Proof.* By compactness it suffices to show that global sections of  $\mathcal{F}(n)$  generate  $\mathcal{F}(n)_x$  for fixed  $x$ . By Nakayama it suffices to show that global sections of  $\mathcal{F}(n)$  generate the fiber  $\mathcal{F}(n)_x \otimes_{\mathcal{O}_{D,x}} \mathbb{C}_x$ .

Picking a hyperplane  $\mathbb{P}_D^{r-1} \simeq H \ni x$  we obtain an exact sequence  $0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_H \rightarrow 0$ , giving an exact sequence  $\mathcal{F}(-1) \xrightarrow{\varphi_1} \mathcal{F} \xrightarrow{\varphi_0} \mathcal{F}_H \rightarrow 0$ . Writing  $\mathcal{P}$  for  $\text{Ker}(\varphi_0) = \text{Im}(\varphi_1)$  we have two exact sequences

$$0 \rightarrow \mathcal{G} \rightarrow \mathcal{F}(-1) \rightarrow \mathcal{P} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathcal{P} \rightarrow \mathcal{F} \rightarrow \mathcal{F}_H \rightarrow 0,$$

noting that  $\mathcal{G}$  and  $\mathcal{F}_H$  are coherent sheaves on  $H$ . Twisting by  $\mathcal{O}(n)$  gives

$$0 \rightarrow \mathcal{G}(n) \rightarrow \mathcal{F}(n-1) \rightarrow \mathcal{P}(n) \rightarrow 0$$

and

$$0 \rightarrow \mathcal{P}(n) \rightarrow \mathcal{F}(n) \rightarrow \mathcal{F}_H(n) \rightarrow 0.$$

The long exact cohomology sequence gives

$$H^1(\mathbb{P}_D^r, \mathcal{F}(n-1)) \rightarrow H^1(\mathbb{P}_D^r, \mathcal{P}(n)) \rightarrow H^2(H, \mathcal{G}(n))$$

and

$$H^1(\mathbb{P}_D^r, \mathcal{P}(n)) \rightarrow H^1(\mathbb{P}_D^r, \mathcal{F}(n)) \rightarrow H^1(H, \mathcal{F}_H(n)).$$

By the assumption  $\mathcal{F}_H$  and  $\mathcal{G}$  are analytifications of algebraic sheaves, so for large  $n$  the terms on the right vanish by Serre's Théorème 1. It follows that  $\dim H^1(\mathbb{P}_D^r, \mathcal{F}(n))$  stabilizes for large  $n$ , and when it does the exact sequences above imply that  $H^1(\mathbb{P}_D^r, \mathcal{P}(n)) \rightarrow H^1(\mathbb{P}_D^r, \mathcal{F}(n))$  is bijective so  $H^0(\mathbb{P}_D^r, \mathcal{F}(n)) \rightarrow H^0(H, \mathcal{F}_H(n))$  is surjective. Since the result holds for analytifications of algebraic sheaves,  $\mathcal{F}_H(n)$  is globally generated for large  $n$ , implying that  $\mathcal{F}(n)_x \otimes_{\mathcal{O}_{D,x}} \mathbb{C}_x$  is generated by global sections, as needed. ♣



## REFERENCES

- ACP** [ACP12] D. Abramovich, L. Caporaso, and S. Payne, *The tropicalization of the moduli space of curves*, December 2012.
- AKMW** [AKMW02] Dan Abramovich, Kalle Karu, Kenji Matsuki, and Jarosław Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. **15** (2002), no. 3, 531–572 (electronic). MR 1896232 (2003c:14016)
- AM** [AM69] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969. MR 0242802 (39 #4129)
- bramovich-Matsuki-Rashid** [AMR99] Dan Abramovich, Kenji Matsuki, and Suliman Rashid, *A note on the factorization theorem of toric birational maps after Morelli and its toroidal extension*, Tohoku Math. J. (2) **51** (1999), no. 4, 489–537. MR 1725624 (2000i:14073)
- ATLuna** [AT15a] Dan Abramovich and Michael Temkin, *Luna’s Fundamental Lemma for diagonalizable groups*, May 2015, <http://arxiv.org/abs/1505.00754>.
- AT1** [AT15b] ———, *Torification of diagonalizable group actions on toroidal schemes*, manuscript in preparation, 2015.
- berihes** [Ber93] Vladimir G. Berkovich, *Étale cohomology for non-Archimedean analytic spaces*, Inst. Hautes Études Sci. Publ. Math. (1993), no. 78, 5–161 (1994). MR 1259429 (95c:14017)
- Borisov-Libgober** [BL05] Lev Borisov and Anatoly Libgober, *McKay correspondence for elliptic genera*, Ann. of Math. (2) **161** (2005), no. 3, 1521–1569. MR 2180406 (2008b:58030)
- Bierstone-Milman** [BM97] Edward Bierstone and Pierre D. Milman, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Invent. Math. **128** (1997), no. 2, 207–302. MR 1440306 (98e:14010)
- Brion-Procesi** [BP90] Michel Brion and Claudio Procesi, *Action d’un tore dans une variété projective*, Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989), Progr. Math., vol. 92, Birkhäuser Boston, Boston, MA, 1990, pp. 509–539. MR 1103602 (92m:14061)
- Buonerba** [Buo15] Federico Buonerba, *Functorial resolution of tame quotient singularities in positive characteristic*, November 2015, <http://arxiv.org/abs/1511.00550>.
- relativeample** [Con06] Brian Conrad, *Relative ampleness in rigid geometry*, Ann. Inst. Fourier (Grenoble) **56** (2006), no. 4, 1049–1126. MR 2266885 (2007h:14029)
- Dolgachev-Hu** [DH98] Igor V. Dolgachev and Yi Hu, *Variation of geometric invariant theory quotients*, Inst. Hautes Études Sci. Publ. Math. (1998), no. 87, 5–56, With an appendix by Nicolas Ressayre. MR 1659282 (2000b:14060)
- Ducros-excellent** [Duc09] Antoine Ducros, *Les espaces de Berkovich sont excellents*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 4, 1443–1552. MR 2566967 (2011a:14048)
- Ducros** [Duc13] A. Ducros, *Families of Berkovich spaces*, November 2013, <http://arxiv.org/abs/1212.0373>.
- Frisch** [Fri67] Jacques Frisch, *Points de platitude d’un morphisme d’espaces analytiques complexes*, Invent. Math. **4** (1967), 118–138. MR 0222336 (36 #5388)
- Grauert-Remmert-Steinspaces** [GR04] Hans Grauert and Reinhold Remmert, *Theory of Stein spaces*, Classics in Mathematics, Springer-Verlag, Berlin, 2004, Translated from the German by Alan Huckleberry, Reprint of the 1979 translation. MR 2029201 (2004k:32015)
- ega** [Gro67] A. Grothendieck, *Éléments de géométrie algébrique.*, Inst. Hautes Études Sci. Publ. Math. (1960-1967).
- Hartshorne** [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52. MR 0463157 (57 #3116)
- Hirflat** [Hir75] Heisuke Hironaka, *Flattening theorem in complex-analytic geometry*, Amer. J. Math. **97** (1975), 503–547. MR 0393556 (52 #14365)
- Illusie-Temkin** [IT14] Luc Illusie and Michael Temkin, *Exposé VIII. Gabber’s modification theorem (absolute case)*, Astérisque (2014), no. 363-364, 103–160, Travaux de Gabber sur l’uniformisation locale et la cohomologie étale des schémas quasi-excellents. MR 3329777
- Iversen** [Ive86] Birger Iversen, *Cohomology of sheaves*, Universitext, Springer-Verlag, Berlin, 1986. MR 842190 (87m:14013)

- Kato-toric** [Kat94] Kazuya Kato, *Toric singularities*, Amer. J. Math. **116** (1994), no. 5, 1073–1099. MR 1296725 (95g:14056)
- Kedlaya-GAGANotes** [Ked09] Kiran Kedlaya, *18.279: Algebraic geometry - GAGA*, MIT OpenCourseWare, [http://ocw.mit.edu/courses/mathematics/18-726-algebraic-geometry-spring-2009/lecture-notes/MIT18\\_726s09\\_lec22\\_gaga.pdf](http://ocw.mit.edu/courses/mathematics/18-726-algebraic-geometry-spring-2009/lecture-notes/MIT18_726s09_lec22_gaga.pdf), 2009.
- KKMS** [KKMSD73] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings. I*, Lecture Notes in Mathematics, Vol. 339, Springer-Verlag, Berlin, 1973. MR 0335518 (49 #299)
- Kollar** [Kol07] János Kollár, *Lectures on resolution of singularities*, Annals of Mathematics Studies, vol. 166, Princeton University Press, Princeton, NJ, 2007. MR 2289519 (2008f:14026)
- Kopf** [Köp74] Ursula Köpf, *Über eigentliche familien algebraischer varietäten über affinoiden räumen*, Schriftenreihe Math. Inst. Munster, 2. Serie. Heft 7, 1974.
- Matsumura** [Mat80] Hideyuki Matsumura, *Commutative algebra*, second ed., Mathematics Lecture Note Series, vol. 56, Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980. MR 575344 (82i:13003)
- GIT** [MFK94] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, third ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], vol. 34, Springer-Verlag, Berlin, 1994. MR 1304906 (95m:14012)
- Morelli** [Mor96] Robert Morelli, *The birational geometry of toric varieties*, J. Algebraic Geom. **5** (1996), no. 4, 751–782. MR 1486987 (99b:14056)
- Noohi** [Noo05] B. Noohi, *Foundations of Topological Stacks I*, ArXiv Mathematics e-prints (2005).
- Poineau** [Poi10] Jérôme Poineau, *Raccord sur les espaces de Berkovich*, Algebra Number Theory **4** (2010), no. 3, 297–334. MR 2602668 (2011g:12009)
- Popescu** [Pop86] Dorin Popescu, *General Néron desingularization and approximation*, Nagoya Math. J. **104** (1986), 85–115. MR 868439 (88a:14007)
- Porta-Yu** [PY14] M. Porta and T. Y. Yu, *Higher analytic stacks and GAGA theorems*, December 2014.
- Serre-GAGA** [Ser56] Jean-Pierre Serre, *Géométrie algébrique et géométrie analytique*, Ann. Inst. Fourier, Grenoble **6** (1955–1956), 1–42. MR 0082175 (18,511a)
- Simpson** [Sim96] C. Simpson, *Algebraic (geometric)  $\mathcal{S}n\mathcal{S}$ -stacks*, September 1996, p. 9014.
- Spivakovskiy** [Spi99] Mark Spivakovskiy, *A new proof of D. Popescu's theorem on smoothing of ring homomorphisms*, J. Amer. Math. Soc. **12** (1999), no. 2, 381–444. MR 1647069 (99j:13008)
- Temkin-embedded** [Tem09] Michael Temkin, *Functorial desingularization over  $\mathbb{Q}$ : boundaries and the embedded case*, December 2009, <http://arxiv.org/abs/0912.2570>.
- Temkin** [Tem12] ———, *Functorial desingularization of quasi-excellent schemes in characteristic zero: the nonembedded case*, Duke Math. J. **161** (2012), no. 11, 2207–2254. MR 2957701
- Thaddeus** [Tha96] Michael Thaddeus, *Geometric invariant theory and flips*, J. Amer. Math. Soc. **9** (1996), no. 3, 691–723. MR 1333296 (96m:14017)
- Ulirsch\_thesis** [Uli15] Martin Ulirsch, *Tropical geometry of logarithmic schemes*, Ph.D. thesis, Brown University, 2015, pp. viii+160.
- W-Cobordism** [Wło00] Jarosław Włodarczyk, *Birational cobordisms and factorization of birational maps*, J. Algebraic Geom. **9** (2000), no. 3, 425–449. MR 1752010 (2002d:14019)
- W-Factor** [Wło03] ———, *Toroidal varieties and the weak factorization theorem*, Invent. Math. **154** (2003), no. 2, 223–331. MR 2013783 (2004m:14113)
- W-ICM** [Wło06] ———, *Algebraic Morse theory and the weak factorization theorem*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 653–682. MR 2275616 (2008e:14016)
- W-Seattle** [Wło09] ———, *Simple constructive weak factorization*, Algebraic geometry—Seattle 2005. Part 2, Proc. Sympos. Pure Math., vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 957–1004. MR 2483958 (2009m:14014)
- Yu** [Yu14] T. Y. Yu, *Gromov compactness in non-archimedean analytic geometry*, January 2014, [arXiv:1401.6452](http://arxiv.org/abs/1401.6452).

DEPARTMENT OF MATHEMATICS, BOX 1917, BROWN UNIVERSITY, PROVIDENCE, RI, 02912,  
U.S.A

*E-mail address:* `abrmovic@math.brown.edu`

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIV'AT RAM,  
JERUSALEM, 91904, ISRAEL

*E-mail address:* `temkin@math.huji.ac.il`