


Conway's Nightmare: Brahmagupta and Butterflies

Richard Evan Schwartz *

February 6, 2022

A *cyclic quad* is a convex quadrilateral whose vertices all lie on the same circle. Equivalently, opposite interior angles sum to π . That is, if we let $\alpha, \beta, \gamma, \delta$ be the successive interior angles, then $\alpha + \gamma = \beta + \delta = \pi$.



Let A be the area of such a quadrilateral and let a, b, c, d be the side lengths. Let

$$B = \sqrt{(s-a)(s-b)(s-c)(s-d)}, \quad s = \frac{a+b+c+d}{2}. \quad (1)$$

Brahmagupta's formula says that $A = B$. I prefer to write

$$A^2 = B^2 = (s-a)(s-b)(s-c)(s-d). \quad (2)$$


Brahmagupta's formula goes back 1400 years. The historical paper [4] discusses how Brahmagupta himself may have proved it. John Conway long sought a simple and beautiful *geometric* proof of Brahmagupta's formula, like Sam Vandervelde's recent proof [7]. Is Brahmagupta's formula really a geometric result? I am not so sure. When generalized to polygons with more

*Supported by N.S.F. Grant DMS-2102802

sides, as in [5], the discussion turns decidedly and deeply algebraic. In any case, I am less interested in a geometric proof than I am in a proof that is short, conceptual, and without calculation.

My proof only depends on basic facts about polynomials and continuity, but I got the idea thinking about things in modern mathematics like flat cone surfaces, control theory, and ergodicity. (It also helps to have smart friends; see the acknowledgements at the end.) I call this proof “Conway’s nightmare” because Sam had been calling his proof “Conway’s dream” in an early draft of his paper. My proof probably would not have satisfied Conway. I will set the stage for the proof, give the proof, then discuss the mathematics that inspired it.

Setting the Stage: Let me explain my favorite proof of the Pythagorean Theorem. If we *fix the angles* of a right triangle T_c with sides a, b, c and hypotenuse c , then $\text{area}(T_c)$ is proportional to c^2 because a and b are both proportional to c . We might write this as $\text{area}(T_c) \propto c^2$.




Put another way, $\text{area}(T_c) = \lambda c^2$, where λ is a constant that depends on the angles. We can divide T_c into two smaller triangles T_a and T_b by dropping the altitude perpendicular to c . Each smaller triangle has the same angles as T_c , so the same λ works for all 3 triangles. Since $\text{area}(T_c) = \text{area}(T_a) + \text{area}(T_b)$ we have $\lambda c^2 = \lambda a^2 + \lambda b^2$. Cancelling λ gives the result.

In this proof, we considered what happens when we vary a right triangle in a special way: changing its size/position without changing the angles. Let us call this operation *morphing*. The name will be more apt when we apply it to cyclic quads. Noting how the relevant quantities associated to a right triangle change when we morph, we get the desired relation up to a constant that cancels out. This proves the Pythagorean Theorem one “angle-type” (a.k.a. *similarity class*) at a time.

My proof of Brahmagupta’s formula has the same flavor. There is one part (second paragraph) that examines how the relevant quantities vary when we morph. – i.e., vary without changing the angles. This analysis alone gives a useful partial result. The other part of the proof (first paragraph) combines the morphing result with an obvious result concerning another operation, *recutting*, to close the deal.

Let me recall two notions that arise in my proof. The *signed distance* between two points p, q on the same line ℓ is the dot product $(p - q) \cdot u$, where u is one of the two unit vectors parallel to ℓ . The *signed area* of a polygon is the sum of the areas of all the regions it bounds, weighted by the number of times the polygon winds around the points in each region. This quantity also has an algebraic expression, in terms of determinants. Both quantities require choices; this amounts to choosing signs for some example.

The Proof: Let $C = A^2/B^2$. Let X be the space of cyclic quads. To *morph* a quad is to replace it by one with the same angles. To *recut* a quad is to cut along a diagonal and reverse one triangle. Recutting preserves C . We claim morphing does too. From any point in X we can reach all nearby points by morphs and recuts. (Recut, morph, re-recut to perturb one pair of opposite angles; repeat using the other diagonal; morph one final time.) Hence C is constant on X . Since X has squares, $C = 1$.



Proof of Claim: Let $L(\rho, \sigma)$ be the space of lines $\ell_a, \ell_b, \ell_c, \ell_d$ with slopes $\rho, \sigma, -\rho, -\sigma$ and $\ell_a \cap \ell_c = (0, 0)$. Parametrize $L \cong \mathbf{R}^2$ by $(x, y) = \ell_b \cap \ell_d$. Let a, b, c, d be the signed distances between vertices of the associated quads, and let A be the signed area. Choose the signs so that $a, b, c, d, A > 0$ in a convex case. $B^2(x, y)$ is a degree 4 polynomial as $a(x, y), \dots, d(x, y)$ are linear. $A(x, y)$, a sum of determinants of linear functions, is a degree 2 polynomial. If $xy = 0$ the quads are *butterflies*, so $A = 0$; also $|a| = |c|$ and $|b| = |d|$ and $ab = -cd$, so 2 factors of B^2 vanish. Given their degrees, $A(x, y) \propto xy$ and $B^2(x, y) \propto (xy)^2$. Hence $C|_L$ is constant. Our claim follows: A (generic) quad and its morphs are all isometric to quads in the same L . ♠

Discussion: The main idea is that the roots, counted with multiplicity, determine a real polynomial up to constants provided that the number of roots equals the degree of the polynomial. A single evaluation then determines the constant. My proof can be summarized like this: The function $C = A^2/B^2$ is invariant under recutting. It is also invariant under morphing because, when analytically continued, A^2 and B^2 vanish to the same order on the set of butterflies and nowhere else. The recutting/morphing process spreads the constancy of C through X like a virus.

To make this idea work, we have to enlarge the space X so that it includes some nonconvex quads, especially butterflies. We'll explain it from another point of view here. First of all, let us modify X so that we consider cyclic quads modulo isometry. Call quads *cousins* if they are morphs of each other. We think of X as a fiber bundle, where the fibers are the cousin families. Each fiber is a convex cone in \mathbf{R}^2 . We create a new space X^* by replacing these cones by the copies of \mathbf{R}^2 which contain them. The space X^* is a plane bundle with the same base. The fibers are our L spaces.

The fact that A is a degree 2 polynomial on the fibers is a key idea of Bill Thurston's paper *Shapes of Polyhedra* [6]. In Thurston's work, he introduces local complex linear coordinates on the space of flat cone spheres with prescribed cone angles. Prescribing the cone angles is like restricting to a fiber. Thurston's coordinates are like my (x, y) coordinates. He shows the area of a flat cone surface with fixed cone angles is the diagonal part of a Hermitian form in his coordinates. There is also a real valued version of this theory which is even closer to my proof, exposted recently in the A.M.S. *Notices* [2] by Danny Calegari. The same ideas also arise in translation surfaces. The main point is that if you fix the slopes of the lines (or the cone angles, in Thurston's case), various algebraic functions are simplified and become linear.

Now we discard X^* and go back to X . Once we know that C is fiber-wise constant on X how do we get $C = 1$? Let me mention two alternate approaches first. Each fiber contains triangles (i.e. degenerate quads) and then we could deduce $C = 1$ by Heron's formula, a degenerate version of Brahmagupta's formula which is somewhat easier to prove. (There are other reductions of Brahmagupta to Heron, e.g. [1].) However, you would still need to prove Heron's formula. Better yet, Peter Doyle noticed that each fiber contains a (perhaps nonconvex) quad whose diagonals are perpendicular, and that for such quads Brahmagupta's formula can be verified with some clever but ultimately easy algebra. I'll leave this as a challenge.

My inspiration for the morphing/recutting proof came from control-theory flavored proofs of ergodicity. The prototypical example is E. Hopf's proof [3] that the geodesic flow on a hyperbolic surface is *ergodic*, meaning that any invariant (measurable) function is (almost everywhere) constant. This may seem far-fetched, but consider the picture.

The geodesic flow lives on a 3-manifold, the unit tangent bundle of the surface. This 3 manifold has 2 invariant codimension 1 foliations, the *stable foliation* and the *unstable foliation*. The first step in Hopf's proof is to use the expansion/contraction properties of the flow along the leaves of the foliations to establish the (almost everywhere) constancy on each leaf of the stable foliation and each leaf of the unstable foliation. The next step is to walk around, going from a stable leaf to an unstable leaf to a stable leaf, etc, to spread this constancy around over the whole 3-manifold. Our space X has 2 codimension 1 foliations, each consisting of quads sharing a pair of opposite interior angles. The method I gave, in the first paragraph of the proof, for moving around in X is the same kind of alternating walk through the leaves of these foliations.

I can't resist explaining 2 other ways I might have done the control theory part. The first approach minimizes "effort" and the second approach minimizes exposition length, but both make extra demands on the reader.

1. Using a finite sequence of morphs and recuts one can start with some point in X and reach an open subset of X . Hence C is constant on an open subset of X . Since X is connected and C is an analytic function, C is constant on X . Since X has squares, $C = 1$.
2. One can join an arbitrary cyclic quad to a square by a finite sequence of morphs and recuts. Hence $C = 1$.

To elaborate on the second approach, let me explain how to change an arbitrary cyclic quad to a square in an algorithmic way. To make the algorithm as clean as possible we note that both morphing and recutting extend to degenerate cyclic quads – i.e. triangles with one marked point. *Morph so as to maximize the intersection angle between the diagonals, recut along the longest diagonal, repeat until done.* The algorithm produces a finite number of marked triangles, then one or two quads with perpendicular diagonals, then a square. To appreciate this, you should restrict it to quads inscribed in the unit circle, and note how it increases the minimum length of the diagonals.

Acknowledgements: Peter Doyle rekindled my interest in Brahmagupta's formula by showing me Sam Vandervelde's proof. Then Peter went on to explain how one can rotate a supplementary quad so that it is a quad of $L(\rho, \sigma)$. Something about this rang a bell, and once I realized that this was just like Thurston's paper the rest fell into place. Peter's insight about the connection to $L(\rho, \sigma)$ was my main inspiration. After I explained my proof to Jeremy Kahn, he suggested the idea of taking the quotient by translation and working in \mathbf{R}^2 rather than \mathbf{R}^4 . This simplified the algebra. Danny Calegari, Dan Margalit, Javi Gomez-Serrano, and Joe Silverman all had helpful expository suggestions.

References

- [1] A. Hess, *A highway from Heron to Brahmagupta*, Forum Geometricorum 12 (2012) pp 121–192.
- [2] D. Calegari, *Pentagonum Pentagonorum*, A.M.S. Notices, Sept 2021
- [3] E. Hopf, *Ergodic theory and the geodesic flow on surfaces of constant negative curvature*, B. A. M. S. 77(6), (1971) pp 863-877
- [4] S. Kichenassamy, *Brahmagupta's derivation of the area of a cyclic quadrilateral*, Historica Mathematica, Elsevier, 2010, 37(1), pp 28–61
- [5] F. M. Maley, D. P. Robbins, J. Roskies, *On the Areas of Cyclic and Semicyclic Polygons*, arXiv: 0407300
- [6] W. Thurston, *Shapes of polyhedra and triangulations of the sphere*, Geometry and Topology Monographs, Volume 1: The Epstein Birthday Schrift, pp 511-549
- [7] S. Vandervelde, *In Search of Conway's Proof*, preprint 2022