

Logarithmic Geometry and Moduli

Dan Abramovich, Qile Chen, Danny Gillam, Yuhao Huang,
Martin Olsson, Matthew Satriano, and Shenghao Sun

ABSTRACT. We discuss the role played by logarithmic structures in the theory of moduli.

Contents

1	Introduction	1
2	Definitions and basic properties	4
3	Differentials, smoothness, and log smooth deformations	8
4	Log smooth curves and their moduli	14
5	D-semistability and log structures	19
6	Stacks of logarithmic structures	24
7	Log deformation theory in general	29
8	Rounding	34
9	Log de Rham and Hodge structures	39
10	The main component of moduli spaces	47
11	Twisted curves and log twisted curves	50
12	Log stable maps	55

1. Introduction

Logarithmic structures in algebraic geometry

It can be said that Logarithmic Geometry is concerned with a method of finding and using “hidden smoothness” in singular varieties. The original insight comes from consideration of de Rham cohomology. Since singular varieties naturally occur “at the boundary” of many moduli problems, logarithmic geometry was soon applied in the theory of moduli.

Foundations for this theory were first given by Kazuya Kato in [27], following ideas of Fontaine and Illusie. The main body of work on logarithmic geometry has been concerned with deep applications in the cohomological study of p -adic

2000 *Mathematics Subject Classification.* Primary 14A20; Secondary 14Dxx.

Key words and phrases. moduli, logarithmic structures.

and arithmetic schemes. This gave the theory an aura of “yet another extremely complicated theory”. The treatments of the theory are however quite accessible. We hope to convince the reader here that the theory is simple enough and useful enough to be considered by anybody interested in moduli of singular varieties, indeed enough to be included in a Handbook of Moduli.

Normal crossings and logarithmic smoothness

So what is the original insight? Let X be a nonsingular complex variety, S a curve with a point s and $f : X \rightarrow S$ a dominant morphism smooth away from s , in such a way that $f^{-1}s = X_s = Y_1 \cup \dots \cup Y_m$ is a reduced simple normal crossings divisor. Then of course $\Omega_{X/S} = \Omega_X/f^*\Omega_S$ fails to be locally free at the singular points of f . But consider instead the sheaves $\Omega_X(\log(X_s))$ of differential forms with at most logarithmic poles along the Y_i , and similarly $\Omega_S(\log(s))$. Then there is an injective sheaf homomorphism $f^*\Omega_S(\log(s)) \rightarrow \Omega_X(\log(X_s))$, and *the quotient sheaf $\Omega_X(\log(X_s))/\Omega_S(\log(s))$ is locally free.*

So in terms of logarithmic forms, *the morphism f is as good as a smooth morphism.*

There is much more to be said: first, this $\Omega_X(\log(X_s))/\Omega_S(\log(s))$ can be extended to a logarithmic de Rham complex, and its hypercohomology, while not recovering the cohomology of the singular fibers, does give rise to the limiting Hodge structure. So it is evidently worth considering.

Second, the picture is quite a bit more general, and can be applied to all toric and toroidal maps between toric varieties or toroidal embeddings (with a little caveat about the characteristic of the residue fields). So there is some flexibility in choosing $X \rightarrow S$.

The search for a structure

Since we are considering moduli, then as soon as we consider $X \rightarrow S$ as above we must also consider the normal crossings fiber $X_s \rightarrow \{s\}$. But what structure should we put on this variety? The notion of differentials with logarithmic poles along X_s is not in itself intrinsic to X_s . Also the normal crossings variety X_s is not in itself toric or toroidal, so a new structure is needed to incorporate it into the picture.

One is tempted to consider varieties which are assembled from nice variety by some sort of gluing, as normal crossings varieties are. But already normal crossings varieties do not give a satisfactory answer in general, because their deformation spaces have “bad” components. Here is a classical example: consider a smooth projective variety Z such that $\text{Pic}^0(Z)$ is nontrivial. Let L be a line bundle on Z and set $Y = \mathbb{P}(\mathcal{O} \oplus L)$, with zero section $Z \subset Y$. Let X be the blowing up of $Z \times 0 \subset Y \times \mathbb{A}^1$. We have a flat morphism $f : X \rightarrow \mathbb{A}^1$ with fiber $X_0 = f^{-1}(0) \simeq$

$Y \cup Y$, where the two copies of Y are glued with the zero section of one attached to the ∞ section of the other.

So clearly X_0 is a normal crossings variety with a nice smoothing to a copy of Y . But there are other deformations: the variety $Y \cup Y$ also deforms to $Y \cup Y'$ where $Y' = \mathbb{P}(\mathcal{O} \oplus L')$ and L' a deformation of the line bundle L . And it is not hard to see that $Y \cup Y'$ does not have a smoothing. Ideally one really does not want to see this deformation $Y \cup Y'$ in the picture - and ideally X_0 should have a natural structure whose deformation space excludes $Y \cup Y'$ automatically.

Such a structure was proposed by Friedman in [10], where the notion of *d-semistable varieties* was introduced. This structure is somewhat subtle, and while it solves the issue in this case, it is not quite as flexible as one could wish. As we will see in Section 5, logarithmic structures subsume d-semistability and do provide an appropriate flexibility.

Organization of this chapter

In this chapter we briefly describe logarithmic structures and indicate where they can be useful in the study of moduli spaces. Section 2 gives the basic definitions of logarithmic structures, and section 3 discusses logarithmic differentials and log smooth deformations, which are important in considering moduli spaces.

Section 4 gives the first example where logarithmic geometry fits well with moduli spaces: the moduli space of stable curves is the moduli space of log smooth curves. The issue of d-semistability does not arise since a nodal curve is automatically d-semistable. So the theory for curves is simple. Turning to higher dimensions, Section 5 shows how d-semistability can be described using logarithmic structures.

If one is to enlarge algebraic geometry to include logarithmic structures, the task of generalizing the techniques of algebraic geometry to logarithmic structure can certainly seem daunting. In section 6 we show how to encode logarithmic structure in terms of certain algebraic stacks. This allows us to reduce various constructions to the case of algebraic stacks. (One can argue that the theory of stacks is not simple either, but at least in the theory of moduli they have come to be accepted, with some exceptions [34].)

In section 7 we make use of logarithmic stacks to describe the complexes which govern deformations and obstructions for logarithmic structures even in the non-smooth case. This comes in handy later. For instance, even when studying moduli of log-smooth schemes, the moduli spaces tend to be singular, and their cotangent complexes are a necessary ingredients in constructing virtual fundamental classes.

Section 8 describes a beautiful construction, similar to polar coordinates, in which families of complex log smooth varieties give rise canonically to families of topological manifolds. Differential geometers have used polar coordinates on nodal

curves to “make space” for monodromy to act by Dehn twists. Rounding (using Ogus’s terminology) is a magnificent way to generalize this.

The immediate implications of logarithmic structures for De Rham cohomology and Hodge structures are described in Section 9.

We conclude by describing three applications, where logarithmic structures serve as the proverbial “magic powder” (term suggested by Kato and Ogus) to clarify or remove unwanted behavior from moduli spaces.

Section 10 describes a number of cases where the main irreducible component of a moduli space can be separated from other “unwanted” components by sprinkling the objects with a bit of logarithmic structures.

In Section 11 we introduce twisted curves, a central object of orbifold stable maps, and show how logarithmic structures give a palatable way to construct the moduli stack of twisted curves.

Section 12 gives background for the work of B. Kim, in which Jun Li’s moduli space of relative stable maps, with its obstruction theory and virtual fundamental class, is beautifully simplified using logarithmic structures.

Notation

Following the lead of Ogus [45], we try whenever possible to denote a logarithmic scheme by a regular letter (such as X) and the underlying scheme by \underline{X} . When this is impossible we write X for the underlying scheme and (X, \mathcal{M}_X) for a logarithmic scheme over it.

Acknowledgements

This chapter originated from lectures given by Olsson at the School and Workshop on Aspects of Moduli, June 15-28, 2008 at the De Giorgi Center at the Scuola Normale Superiore in Pisa, Italy. The material was revisited and expanded in our seminar during the Algebraic Geometry program at MSRI, 2009. We thank the De Giorgi Center, MSRI, their staff and program organizers for providing these opportunities. Thanks are due to Arthur Ogus and Phillip Griffiths, who lectured on two topics at the MSRI seminar. While no new material is intended here, we acknowledge that research by Abramovich, Gillam and Olsson is supported by the NSF. Finally thanks are due to Gavril Farkas and Ian Morrison for the invitation to write this chapter.

2. Definitions and basic properties

In this section we introduce the basic definitions of logarithmic geometry in the sense of [27]. Good introductions are given in [27] and [45]. Further technique is developed in [12].

Logarithmic structures

The basic definitions are as follows:

Definition 2.1. A *monoid* is a commutative semi-group with a unit. A morphism of monoids is required to preserve the unit element. We use Mon to denote the category of Monoids.

Definition 2.2. Let \underline{X} be a scheme. A *pre-logarithmic structure* on \underline{X} is a sheaf of monoids \mathcal{M}_X on the étale site $\underline{X}_{\text{ét}}$ combined with a morphism of sheaves of monoids: $\alpha : \mathcal{M}_X \rightarrow \mathcal{O}_{\underline{X}}$, called the structure morphism, where we view $\mathcal{O}_{\underline{X}}$ as a monoid under multiplication. A pre-log structure is called a *log structure* if $\alpha^{-1}(\mathcal{O}_{\underline{X}}^*) \cong \mathcal{O}_{\underline{X}}^*$ via α . The pair $(\underline{X}, \mathcal{M}_X)$ is called a *log scheme*, and will be denoted by X .

Note that, given a log structure \mathcal{M}_X on \underline{X} , we can view $\mathcal{O}_{\underline{X}}^*$ as a subsheaf of \mathcal{M}_X .

Definition 2.3. Given a log scheme X , the quotient sheaf $\overline{\mathcal{M}}_X = \mathcal{M}_X / \mathcal{O}_{\underline{X}}^*$ is called the *characteristic of the log structure* \mathcal{M}_X .

Definition 2.4. Let \mathcal{M} and \mathcal{N} be pre-log structures on \underline{X} . A *morphism* between them is a morphism $\mathcal{M} \rightarrow \mathcal{N}$ of sheaves of monoids which is compatible with the structure morphisms.

How should one think of such a beast? There are two extreme cases:

- (1) If an element $m \in \mathcal{M}$ has $\alpha(m) = x \neq 0$, one often thinks of m as some sort of partial data of a “branch of the logarithm of x ”. Evidently no data is added if x is invertible, but some is added otherwise. In particular, we will see later that m permits us to take the logarithmic differential dx/x of x .
- (2) If $\alpha(m) = 0$ it is often the case that m comes by restricting the log structure of an ambient space, and serves as the “ghost” of a logarithmic cotangent vector coming from that space. So the log structure “remembers” deformations that are lost when looking at the underlying scheme.

The log structure associated to a pre-log structure

We have a natural inclusion

$$i : (\text{log structures on } \underline{X}) \hookrightarrow (\text{pre-log structures on } \underline{X})$$

by viewing a log structure as a pre-log structure. We now construct a left adjoint.

Let $\alpha : \mathcal{M} \rightarrow \mathcal{O}_{\underline{X}}$ be a pre-log structure on \underline{X} . We define the *associated log structure* \mathcal{M}^a to be the push-out of

$$\begin{array}{ccc} \alpha^{-1}(\mathcal{O}_{\underline{X}}^*) & \longrightarrow & \mathcal{M} \\ \downarrow & & \\ \mathcal{O}_{\underline{X}}^* & & \end{array}$$

in the category of sheaves of monoids on $\underline{X}_{\acute{e}t}$, endowed with

$$\mathcal{M}^a \rightarrow \mathcal{O}_{\underline{X}} \quad (a, b) \mapsto \alpha(a)b \quad (a \in \mathcal{M}, b \in \mathcal{O}_{\underline{X}}^*).$$

In this way, we obtain a functor $a : (\text{pre-log structures on } \underline{X}) \rightarrow (\text{log structures on } \underline{X})$. From the universal property of push-out, any morphism of pre-log structure from a pre-log structure \mathcal{M} to a log structure on \underline{X} factor through \mathcal{M}^a uniquely.

Lemma 2.5. [45, 1.1.5] *The functor a is left adjoint to i .*

Example 2.6. The category of log structures on \underline{X} has an initial object, called the trivial log structure, given by the inclusion $\mathcal{O}_{\underline{X}}^* \hookrightarrow \mathcal{O}_{\underline{X}}$. It also has a final object, given by the identity map $\mathcal{O}_{\underline{X}} \rightarrow \mathcal{O}_{\underline{X}}$. Trivial log structures are quite useful as they make the category of schemes into a full subcategory of the category of log schemes (see Definition 2.9). The final object is rarely used since it is not fine, see definition 2.16.

Example 2.7. Let \underline{X} be a regular scheme, and $D \subset \underline{X}$ a divisor. We can define a log structure \mathcal{M} on \underline{X} associated to the divisor D as

$$\mathcal{M}(U) = \{g \in \mathcal{O}_{\underline{X}}(U) : g|_{U \setminus D} \in \mathcal{O}_{\underline{X}}^*(U \setminus D)\} \subset \mathcal{O}_{\underline{X}}(U).$$

The case where D is a normal crossings divisor is special - we will see later that it is *log smooth*.

Note that the concept of normal crossing is local in the étale topology. This is one reason we use the étale topology instead of the Zariski topology.

Example 2.8. Let P be a monoid, R a ring, and denote by $R[P]$ the monoid algebra. Let $\underline{X} = \text{Spec } R[P]$. Then \underline{X} has a canonical log structure associated to the canonical map $P \rightarrow R[P]$. We denote by $\text{Spec}(P \rightarrow R[P])$ the log scheme with underlying \underline{X} , and the canonical log structure.

The inverse image and the category of log schemes

Let $f : \underline{X} \rightarrow \underline{Y}$ be a morphism of schemes. Given a log structure \mathcal{M}_Y on \underline{Y} , we can define a log structure on \underline{X} , called the inverse image of \mathcal{M}_Y , to be the log structure associated to the pre-log structure $f^{-1}(\mathcal{M}_Y) \rightarrow f^{-1}(\mathcal{O}_Y) \rightarrow \mathcal{O}_{\underline{X}}$. This is usually denoted by $f^*(\mathcal{M}_Y)$. Using the inverse image of log structures, we can give the following definition.

Definition 2.9. A *morphism of log schemes* $X \rightarrow Y$ consists of a morphism of underlying schemes $f : \underline{X} \rightarrow \underline{Y}$, and a morphism $f^\flat : f^* \mathcal{M}_Y \rightarrow \mathcal{M}_X$ of log structures on \underline{X} .

We denote by $LSch$ the category of log schemes.

Example 2.10. In Example 2.8, the log structure on $\text{Spec}(P \rightarrow R[P])$ can be viewed as the inverse image of the log structure on $\text{Spec}(P \rightarrow \mathbb{Z}[P])$ via the canonical map $\text{Spec}(R[P]) \rightarrow \text{Spec}(\mathbb{Z}[P])$.

Example 2.11. Let k be a field, $\underline{Y} = \text{Spec } k[x_1, \dots, x_n]$, $D = V(x_1 \cdots x_r)$. Note that D is a normal crossing divisor in \underline{Y} . By example 2.7, we have a log structure \mathcal{M}_Y on \underline{Y} associated to the divisor D . In fact, \mathcal{M}_Y can be viewed as a subsheaf of $\mathcal{O}_{\underline{Y}}$ generated by $\mathcal{O}_{\underline{Y}}^*$ and $\{x_1, \dots, x_r\}$.

Consider the inclusion $j : p = \text{Spec } k \hookrightarrow \underline{Y}$ sending the point to the origin of \underline{Y} . Then $j^* \mathcal{M}_Y = k^* \oplus \mathbb{N}^r$, and the structure map $j^* \mathcal{M} \rightarrow \mathcal{O}_{\underline{X}}$ is given by $(a, n_1, \dots, n_r) \mapsto a \cdot 0^{n_1 + \dots + n_r}$, where we define $0^0 = 1$ and $0^n = 0$ if $n \neq 0$. Such point with the log structure above is called a *logarithmic point*; when $r = 1$ we call it the *standard logarithmic point*.

Charts of log structures

Definition 2.12. Let X be a log scheme, and P a monoid. A *chart* for \mathcal{M}_X is a morphism $P \rightarrow \Gamma(X, \mathcal{M}_X)$, such that the induced map of log structures $P^a \rightarrow \mathcal{M}_X$ is an isomorphism, where P^a is the log structure associated to the pre-log structure given by $P \rightarrow \Gamma(X, \mathcal{M}_X) \rightarrow \Gamma(X, \mathcal{O}_{\underline{X}})$.

In fact, a chart of \mathcal{M}_X is equivalent to a morphism

$$f : X \rightarrow \text{Spec}(P \rightarrow \mathbb{Z}[P]),$$

such that f^\flat is an isomorphism. In general, we have the following:

Lemma 2.13. [45, 1.1.9] *The morphism*

$$\text{Hom}_{LSch}(X, \text{Spec}(P \rightarrow \mathbb{Z}[P])) \rightarrow \text{Hom}_{Mon}(P, \Gamma(X, \mathcal{M}_X))$$

associating to f the composition

$$P \longrightarrow \Gamma(\underline{X}, P_X) \xrightarrow{\Gamma(f^\flat)} \Gamma(\underline{X}, \mathcal{M}_X)$$

is an isomorphism.

We can also consider charts for log morphisms.

Definition 2.14. Let $f : X \rightarrow Y$ be a morphism of log schemes. A *chart* for f is a triple $(P_X \rightarrow \mathcal{M}_X, Q_Y \rightarrow \mathcal{M}_Y, Q \rightarrow P)$ where P_X and Q_Y are the constant sheaves associated to the monoids P and Q , which satisfy the following conditions:

- (1) $P_X \rightarrow \mathcal{M}_X$ and $Q_Y \rightarrow \mathcal{M}_Y$ are charts of \mathcal{M}_X and \mathcal{M}_Y ;

- (2) the morphism of monoids $Q \rightarrow P$ makes the following diagram commutative:

$$\begin{array}{ccc} Q_X & \longrightarrow & P_X \\ \downarrow & & \downarrow \\ f^* \mathcal{M}_Y & \longrightarrow & \mathcal{M}_X. \end{array}$$

Fine log structures

Arbitrary log structures are too wild to manipulate; they are roughly analogous to arbitrary ringed spaces: both notions are useful for general constructions, but a narrower, more geometric category is desirable. In Definition 2.16 below we introduce the notion of fine log structures. Continuing the analogy above, these are well-behaved log structures analogous to noetherian schemes, in the sense that you can do geometry on them.

Given a monoid P , we can associate a group

$$P^{gp} := \{(a, b) \mid (a, b) \sim (c, d) \text{ if } \exists s \in P \text{ such that } s + a + d = s + b + c\}.$$

Note that any morphism from P to an abelian group factors through P^{gp} uniquely.

Definition 2.15. P is called *integral* if $P \rightarrow P^{gp}$ is injective. It is called *saturated* if it is integral and for any $p \in P^{gp}$, if $n \cdot p \in P$ for some positive integer n then $p \in P$.

Definition 2.16. A log scheme X is said to be *fine*, if étale locally there is a chart $P \rightarrow \mathcal{M}_X$ with P a finitely generated integral monoid. If moreover P can be chosen to be saturated, then X is called a *fine and saturated* (or *fs*) log structure. This is equivalent to saying that for every geometric point $\bar{x} \rightarrow \underline{X}$ the monoid $\overline{\mathcal{M}}_{X, \bar{x}}$ is saturated as in Definition 2.15. Finally if $P \simeq \mathbb{N}^k$ we say that the log structure is *locally free*.

In the following, we will focus on fine log schemes.

3. Differentials, smoothness, and log smooth deformations

The main reference in this section is [27].

Logarithmic differentials

In [14] Grothendieck defines a derivation as the difference of infinitesimal liftings of a section. We can do the same thing with logarithmic schemes. First, we need a concept of infinitesimal extension, which requires the following definition.

Definition 3.1. A morphism $f : X \rightarrow Y$ of log schemes is called *strict* if $f^\flat : f^* \mathcal{M}_Y \rightarrow \mathcal{M}_X$ is an isomorphism. It is called a *strict closed immersion*¹ if it is strict and the underlying map $\underline{X} \rightarrow \underline{Y}$ is a closed immersion in the usual sense.

¹The term used in [27] is *an exact closed immersion*.

Let us consider a commutative diagram of solid arrows of log schemes:

$$\begin{array}{ccc}
 T_0 & \xrightarrow{\phi} & X \\
 \downarrow J & \swarrow g_1 & \downarrow f \\
 T_1 & \xrightarrow{\psi} & Y
 \end{array}$$

where j is a strict closed immersion defined by an ideal J with $J^2 = 0$. Note that T_0 and T_1 have the same underlying topological space, and isomorphic étale sites. Then we have the following commutative diagram of sheaves of algebras:

$$\begin{array}{ccc}
 \mathcal{O}_{T_0} & \xleftarrow{\phi^{-1}} & \mathcal{O}_X \\
 \uparrow & \swarrow g_1^\# & \uparrow \\
 \mathcal{O}_{T_1} & \xleftarrow{\psi^{-1}} & \mathcal{O}_Y
 \end{array}$$

Then $g_1^\# - g_2^\#$ is a derivation $\partial_{g_1-g_2} : \phi^{-1}\mathcal{O}_X \rightarrow J$ in the usual sense. We also have a commutative diagram given by the log structures:

$$\begin{array}{ccc}
 \mathcal{M}_{T_0} & \xleftarrow{\phi^{-1}} & \mathcal{M}_X \\
 \uparrow & \swarrow g_1^b & \uparrow \\
 \mathcal{M}_{T_1} & \xleftarrow{\psi^{-1}} & \mathcal{M}_Y
 \end{array}$$

Note that we have an “exact sequence” of multiplicative monoids

$$\mathbf{1} \rightarrow (1 + J) \rightarrow \mathcal{M}_{T_1} \rightarrow \mathcal{M}_{T_0} \rightarrow \mathbf{1},$$

by which we mean that the group $1 + J$ acts freely on \mathcal{M}_{T_1} with quotient \mathcal{M}_{T_0} . Hence we obtain a morphism $D_{g_1-g_2} : \phi^{-1}\mathcal{M}_X \rightarrow J$ such that for every $m \in \phi^{-1}\mathcal{M}_X$ we have $(g_1^b - g_2^b)(m) = 1 + D_{g_1-g_2}(m)$. It is not hard to check that it is a monoid homomorphism: $D_{g_1-g_2}(m \cdot n) = D_{g_1-g_2}(m) + D_{g_1-g_2}(n)$ for any $m, n \in \phi^{-1}(\mathcal{M}_X)$. By the definition of log structures, we also have

- (1) $\alpha(m)D_{g_1-g_2}m = \partial_{g_1-g_2}(\alpha(m)), \forall m \in \phi^{-1}\mathcal{M}_X$;
- (2) $D_{g_1-g_2}|_{\psi^{-1}\mathcal{M}_Y} = 0$.

Remark 3.2. (1) Since the log structure contains all the invertible elements in the structure sheaf, the map $D_{g_1-g_2}$ determines $\partial_{g_1-g_2}$.

- (2) The above properties show that $D_{g_1-g_2}$ behaves like “ $d \log$ ”. This is one of the reasons for the name “logarithmic structure”.

Summarizing the above discussion gives the following definitions:

Definition 3.3. [44, Sec. 1.1],[45, Definition 1.1.1] Consider a morphism $f : X \rightarrow Y$ of fine log schemes. Let I be an \mathcal{O}_X -module. A *log derivation of X over Y to I*

is a pair (∂, D) where $\partial \in \mathcal{D}er_{\underline{Y}}(\underline{X}, I)$ and $D : \mathcal{M}_X \rightarrow I$ is an additive map such that the following conditions hold:

- (1) $D(ab) = D(a) + D(b)$ for $a, b \in \mathcal{M}_X$;
- (2) $\alpha(a)D(a) = \partial(\alpha(a))$, for $a \in \mathcal{M}_X$.
- (3) $D(a) = 0$, for $a \in f^{-1}\mathcal{M}_Y$.

The sheaf $\mathcal{D}er_Y(X, I)$ of log derivations of X over Y to I is the sheaf of germs of pairs (∂, D) . The sheaf $\mathcal{D}er_Y(X, \mathcal{O}_{\underline{X}})$ is usually denoted by $T_{X/Y}$, and is called *the logarithmic tangent sheaf of X over Y* .

As an analogue of differentials of usual schemes, we have the following result:

Proposition 3.4. [45, IV.1.1.6] *of Log differentials* There exists an $\mathcal{O}_{\underline{X}}$ -module $\Omega_{X/Y}^1$ with a universal derivations $(\partial, D) \in \mathcal{D}er_Y(X, \Omega_{X/Y}^1)$, such that for any $\mathcal{O}_{\underline{X}}$ -module I , the canonical map

$$\mathcal{H}om_{\mathcal{O}_{\underline{X}}}(\Omega_{X/Y}^1, I) \rightarrow \mathcal{D}er_Y(X, I), \quad u \mapsto (u \circ \partial, u \circ D)$$

is an isomorphism of $\mathcal{O}_{\underline{X}}$ -modules. In fact, we have the following construction:

$$\Omega_{X/Y}^1 = \Omega_{\underline{X}/\underline{Y}} \oplus (\mathcal{O}_{\underline{X}} \otimes_{\mathbb{Z}} \mathcal{M}_X^{gp}) / \mathcal{K}$$

where \mathcal{K} is the $\mathcal{O}_{\underline{X}}$ -module generated by local sections of the following forms:

- (1) $(d\alpha(a), 0) - (0, \alpha(a) \otimes a)$ with $a \in \mathcal{M}_X$;
- (2) $(0, 1 \otimes a)$ with $a \in \text{Im}(f^{-1}(\mathcal{M}_Y) \rightarrow \mathcal{M}_X)$.

The universal derivation (∂, D) is given by $\partial : \mathcal{O}_{\underline{X}} \xrightarrow{d} \Omega_{\underline{X}/\underline{Y}} \rightarrow \Omega_{X/Y}^1$ and $D : \mathcal{M}_X \rightarrow \mathcal{O}_{\underline{X}} \otimes_{\mathbb{Z}} \mathcal{M}_X^{gp} \rightarrow \Omega_{X/Y}^1$.

Definition 3.5. Given a morphism $f : X \rightarrow Y$ of log schemes, the $\mathcal{O}_{\underline{X}}$ -module $\Omega_{X/Y}^1$ is called *the sheaf of logarithmic differentials*. Sometimes we use the short notation Ω_f^1 for $\Omega_{X/Y}^1$.

Note that $\mathcal{H}om(\Omega_{X/Y}^1, \mathcal{O}_{\underline{X}}) \cong T_{X/Y}$.

Remark 3.6. If we consider only fine log structures, and assume that \underline{Y} is locally noetherian and \underline{X} locally of finite type over \underline{Y} , then both $\mathcal{D}er_Y(X, I)$ and $\Omega_{X/Y}^1$ in the definitions above are coherent sheaves. The proof of this can be found in [45, IV.1.1]

Example 3.7. Consider $R = k[x_1, \dots, x_n]/(x_1 \cdots x_r)$, where k is a field. Denote $\underline{X} = \text{Spec } R$. Let \mathcal{M}_X be the log structure on \underline{X} given by $\mathbb{N}^r \rightarrow R$, $e_i \mapsto x_i$, where e_i is the standard generator of the monoid \mathbb{N}^r . Let $Y = \text{Spec } (\mathbb{N} \rightarrow k)$ be the logarithmic point described in 2.11. Now we can define a morphism $f : X \rightarrow Y$ by the following diagram:

$$\begin{array}{ccc} \mathbb{N}^r & \longrightarrow & R \\ \Delta \uparrow & & \uparrow \\ \mathbb{N} & \longrightarrow & k \end{array}$$

where $\Delta : e \mapsto e_1 + \cdots + e_r$, and e is the standard generator of \mathbb{N} . Then it is easy to see that $\mathcal{D}er_Y(X, \mathcal{O}_{\underline{X}})$ is a free $\mathcal{O}_{\underline{X}}$ -module generated by

$$x_1 \frac{\partial}{\partial x_1}, \cdots, x_r \frac{\partial}{\partial x_r}, \frac{\partial}{\partial x_{r+1}}, \cdots, \frac{\partial}{\partial x_n},$$

with a relation $x_1 \frac{\partial}{\partial x_1} + \cdots + x_r \frac{\partial}{\partial x_r} = 0$. The sheaf Ω_f^1 is a free \mathcal{O}_X -module generated by the logarithmic differentials:

$$\frac{dx_1}{x_1}, \cdots, \frac{dx_r}{x_r}, dx_{r+1}, \cdots, dx_n$$

with a relation $\frac{dx_1}{x_1} + \cdots + \frac{dx_r}{x_r} = 0$.

Example 3.8. Let $h : Q \rightarrow P$ be a morphism of fine monoids. Denote $X = \text{Spec}(P \rightarrow \mathbb{Z}[P])$ and $Y = \text{Spec}(Q \rightarrow \mathbb{Z}[Q])$. Then we have a morphism $f : X \rightarrow Y$ induced by h . A direct calculation shows that $\Omega_f^1 = \mathcal{O}_{\underline{X}} \otimes \mathbf{Cok}(h^{gp})$. This can also be seen from the universal property of the sheaf of logarithmic differentials.

Logarithmic Smoothness

Let us go back to the following cartesian diagram of log schemes:

$$(3.9) \quad \begin{array}{ccc} T_0 & \xrightarrow{\phi} & X \\ J \downarrow j & & \downarrow f \\ T_1 & \xrightarrow{\psi} & Y \end{array}$$

where j is a strict closed immersion defined by J with $J^2 = 0$. As in the usual case, we can define log smoothness by the infinitesimal lifting property.

Definition 3.10. A morphism $f : X \rightarrow Y$ of fine log schemes is called *log smooth* (resp. *étale*) if the underlying morphism $\underline{X} \rightarrow \underline{Y}$ is locally of finite presentation and for any commutative diagram (3.9), étale locally on T_1 there exists a (resp. there exists a unique) morphism $g : T_1 \rightarrow X$ such that $\phi = g \circ j$ and $\psi = f \circ g$.

We have the following useful criterion for smoothness from [27, Theorem 3.5].

Theorem 3.11. (*K.Kato*) *Let $f : X \rightarrow Y$ be a morphism of fine log schemes. Assume we have a chart $Q \rightarrow \mathcal{M}_Y$, where Q is a finitely generated integral monoid. Then the following are equivalent:*

- (1) f is log smooth (resp. log étale);
- (2) étale locally on X , there exists a chart $(P_X \rightarrow \mathcal{M}_X, Q_Y \rightarrow \mathcal{M}_Y, Q \rightarrow P)$ extending the chart $Q_Y \rightarrow \mathcal{M}_Y$, satisfying the following properties.
 - (a) The kernel and the torsion part of the cokernel (resp. the kernel and the cokernel) of $Q^{gp} \rightarrow P^{gp}$ are finite groups of order invertible on X .

- (b) *The induced morphism from $\underline{X} \rightarrow \underline{Y} \times_{\text{Spec } \mathbb{Z}[Q]} \text{Spec } \mathbb{Z}[P]$ is étale in the classical sense.*

Remark 3.12. (1) We can require $Q^{gp} \rightarrow P^{gp}$ in (a) to be injective, and replace the requirement of $\underline{X} \rightarrow \underline{Y} \times_{\text{Spec } \mathbb{Z}[Q]} \text{Spec } \mathbb{Z}[P]$ be étale in (b) by requiring it to be smooth without changing the conclusion of the theorem 3.11.

- (2) The arrow in (b) shows that a log smooth morphism is “locally toric” relative to the base. Consider the case Y is a log scheme with underlying space given by $\text{Spec } \mathbb{C}$ with the trivial log structure, and $X = \text{Spec}(P \rightarrow \mathbb{C}[P])$ where P is a fine, saturated and torsion free monoid. Then \underline{X} is a toric variety with the action of $\text{Spec } \mathbb{C}[P^{gp}]$. According to the theorem, X is log smooth relative to Y , though the underlying space might be singular. These singularities are called toric singularities in [28]. This is closely related to the classical notion of toroidal embeddings, see [32].

Example 3.13. Using the theorem, we can check directly that the morphism f in example 3.7 is log smooth, but the underlying map has normal crossing singularities. We will see later that one of the major advantages of log structures is in dealing with the normal crossing singularities.

Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be morphisms of fine log schemes. Consider the sheaves of log differentials Ω_g^1 and $\Omega_{g \circ f}^1$, with their universal derivations (∂_g, D_g) and $(\partial_{g \circ f}, D_{g \circ f})$ respectively. We have a canonical map $f^* \Omega_g^1 \rightarrow \Omega_{g \circ f}^1$ induced by

$$f^*(\partial_g u) \mapsto \partial_{g \circ f} f^*(u) \quad \text{and} \quad f^*(D_g v) \mapsto D_{g \circ f} f^\flat(v),$$

where $u \in \mathcal{O}_Y$ and $v \in \mathcal{M}_Y$. Denote by (∂_f, D_f) the universal derivation associated to Ω_f^1 . Similarly, we have a canonical map $\Omega_{g \circ f}^1 \rightarrow \Omega_f^1$ induced by

$$\partial_{g \circ f} u' \mapsto \partial_f u' \quad \text{and} \quad D_{g \circ f} v' \mapsto D_f v',$$

where $u' \in \mathcal{O}_X$ and $v' \in \mathcal{M}_X$. The following proposition shows that log differentials behave like usual differentials, especially for log smooth morphisms.

Proposition 3.14. (1) *The sequence $f^* \Omega_g^1 \rightarrow \Omega_{g \circ f}^1 \rightarrow \Omega_f^1 \rightarrow 0$ is exact.*

- (2) *If f is log smooth, then Ω_f^1 is a locally free \mathcal{O}_X -module, and we have the following exact sequence: $0 \rightarrow f^* \Omega_g^1 \rightarrow \Omega_{g \circ f}^1 \rightarrow \Omega_f^1 \rightarrow 0$.*

- (3) *If $g \circ f$ is log smooth and the sequence in (2) is exact and splits locally, then f is log smooth.*

A proof can be found in [45, Chapter IV].

Logarithmic smooth deformation

Having discussed log smoothness, a natural thing to do is to develop log smooth deformations. In many cases, we would require this to be a flat deformation for the underlying space. Unfortunately log smoothness does not imply flatness, so we need the following definition.

Definition 3.15. A map of fine monoids $h : Q \rightarrow P$ is called *integral* if the induced map on monoid algebra $\mathbb{Z}[Q] \rightarrow \mathbb{Z}[P]$ is flat.

Definition 3.16. A morphism $f : X \rightarrow Y$ of fine log schemes is called *integral* if for every geometric point $\bar{x} \in X$, the map of characteristic monoids $h : f^{-1}(\overline{\mathcal{M}}_Y)_{\bar{x}} \rightarrow \overline{\mathcal{M}}_{X,\bar{x}}$ is integral.

Remark 3.17. (1) If f is integral, then étale locally we have a chart $(P_X \rightarrow \mathcal{M}_X, Q_Y \rightarrow \mathcal{M}_Y, Q \xrightarrow{h} P)$ such that h is integral.
 (2) If $h : Q \rightarrow P$ is integral of integral monoids, then for any integral monoid Q' , the push-out of $P \leftarrow Q \rightarrow Q'$ in the category of monoids is integral. Thus integral morphisms are stable under base change by integral log schemes.
 (3) Given a morphism $h : Q \rightarrow P$ of integral monoids, there is an explicit criterion, which looks complicated, but sometimes is useful for checking integrality of h directly: if $a_1, a_2 \in Q$, $b_1, b_2 \in P$ and $h(a_1)b_1 = h(a_2)b_2$, then there exist $a_3, a_4 \in Q$ and $b \in P$ such that $b_1 = h(a_3)b, b_2 = h(a_4)b$ and $a_1a_3 = a_2a_4$. This comes essentially from the equational criterion for flatness.

Now we have the following fact from [27, 4.5].

Proposition 3.18. *If f is a log smooth and integral morphism of fine log schemes, then f the underlying map is flat in the usual sense.*

Now let us consider the following deformation problem. We are given a log smooth integral morphism $f_0 : X_0 \rightarrow B_0$ of fine log schemes, and a strict closed immersion $j : B_0 \rightarrow B$ defined by an ideal J with $J^2 = 0$. We want to find a log smooth lifting $f : X \rightarrow B$ fitting in the following cartesian diagram:

$$\begin{array}{ccc} X_0 & \hookrightarrow & X \\ \downarrow & & \downarrow \\ B_0 & \hookrightarrow & B. \end{array}$$

Remark 3.19. Since f_0 is integral, and $\mathcal{M}_X/(1+J) \cong \mathcal{M}_{X_0}$, it is not hard to show that the lifting f is automatically integral and hence flat.

We have the following theorem for log smooth deformations.

Theorem 3.20. [27, 3.14] *With the notation as above, we have:*

- (1) *There is a canonical obstruction $\eta \in H^2(\underline{X}_0, T_{X_0/B_0} \otimes J)$ such that $\eta = 0$ if and only if there exists a log smooth lifting.*
- (2) *If $\eta = 0$, then the set of log smooth deformations form a torsor under $H^1(\underline{X}_0, T_{X_0/B_0} \otimes J)$.*

- (3) *The automorphism group of any deformation is given by $H^0(\underline{X}_0, T_{X_0/B_0} \otimes J)$.*

The theorem can be proved in a manner similar to the case of usual deformation theory as in [15, Exposé 3]. Another proof using the logarithmic cotangent complex can be found in [49, Thm 5.6], which we will discuss later.

4. Log smooth curves and their moduli

Now that we have reviewed some of the foundations, we can discuss the first application to the study of moduli spaces: F. Kato's interpretation of $\overline{\mathcal{M}}_{g,n}$ as a moduli space for log curves. The general philosophy is expressed by F. Kato in the introduction to [26]:

Philosophy. Since log smoothness includes some degenerating objects like semistable reductions, etc., the moduli space of log smooth objects should be already compactified, once its existence has been established.

Along the lines of this philosophy, to compactify $\mathcal{M}_{g,n}$, we want to introduce a notion of *log curve* which extends the notion of smooth curve. Following F. Kato, we do so after some preliminaries.

Relative characteristic sheaves

Recall from Definition 2.3 that the characteristic $\overline{\mathcal{M}}_X$ of a log scheme X is defined as $\mathcal{M}_X/\mathcal{O}_X^*$. In the study of log curves, the following relative notion of characteristic plays an important role.

Definition 4.1. Given a morphism $f : X \rightarrow Y$ of log schemes, the *relative characteristic* $\overline{\mathcal{M}}_{X/Y}$ is defined as the quotient $\mathcal{M}_X/\text{im}(f^*\mathcal{M}_Y \rightarrow \mathcal{M}_X)$ in the category of integral monoids.

Example 4.2. Let $f : X \rightarrow Y$ be the morphism from Example 3.7. Then the relative characteristic $\overline{\mathcal{M}}_{X/Y}$ is the cokernel in the category of integral monoids of the diagonal map $\Delta : \mathbb{N} \rightarrow \mathbb{N}^2$, which is \mathbb{Z} .

Lemma 4.3 ([26, Lemma 1.6]). *If $f : X \rightarrow Y$ is an integral morphism of fine log schemes, then $\overline{\mathcal{M}}_{X/Y, \bar{x}} = 0$ if and only if f is strict in an étale neighborhood of x .*

As the following example illustrates, the integrality assumption on f is necessary.

Example 4.4. Let P be the monoid on three generators x, y , and z subject to the relation $x + y = 2z$. We have an injection

$$i : P \longrightarrow \mathbb{N}^2$$

sending x to $(2, 0)$, y to $(1, 1)$, and z to $(0, 2)$. Let $X = \text{Spec } k[\mathbb{N}^2]$ and $Y = \text{Spec } k[P]$ with their canonical log structures. Then it follows from [27, Prop 3.4]

that the morphism of log schemes $f : X \rightarrow Y$ induced by i is log étale, but \underline{f} is not flat, and hence, by [27, Cor 4.5], f is not integral. It is easy to check that

$$\mathcal{M}_{X/Y,0} = \mathbb{N}^2/P \simeq \mathbb{Z}/2,$$

so $\overline{\mathcal{M}}_{X/Y,0} = 0$, but f is not strict.

Log curves

Definition 4.5. A *log curve* is a log smooth integral morphism $f : X \rightarrow S$ of fs log schemes such that the geometric fibers of \underline{f} are reduced connected 1-dimensional schemes.

We require f to be integral so that, by [27, Cor 4.5], f is flat. The reason for the fs assumption is to avoid cusps or worse singularities, as the following example shows.

Example 4.6. If $X = \text{Spec } k[\mathbb{N} - \{1\}]$ is given its canonical log structure and $S = \text{Spec } k$ is given the trivial log structure, then $X \rightarrow S$ is log smooth and integral; however, $\underline{X} = \text{Spec } k[x, y]/(y^2 - x^3)$ has a cusp.

It is a remarkable fact that by endowing our curves with log structures as in Definition 4.5, this is enough to control the singularities of the curve.

Theorem 4.7 ([26, Thm 1.3]). *If k is a separably closed field and $f : X \rightarrow S$ is a log curve with $\underline{S} = \text{Spec } k$, then \underline{X} has at worst nodal singularities. Moreover, if r_1, \dots, r_ℓ are the nodes of \underline{X} , then there exist smooth points s_1, \dots, s_n of \underline{X} such that*

$$\overline{\mathcal{M}}_{X/S} = \mathbb{Z}_{r_1} \oplus \dots \oplus \mathbb{Z}_{r_\ell} \oplus \mathbb{N}_{s_1} \oplus \dots \oplus \mathbb{N}_{s_n};$$

here M_x denotes the skyscraper sheaf for a monoid M supported at a point $x \in \underline{X}$.

The reader should think of the s_i in the above theorem as marked points. So we can already see how n -pointed curves emerge naturally from the log geometry perspective.

Example 4.8. Consider the closed subscheme \underline{X} of $\mathbb{P}_k^2 \times_k \mathbb{A}_k^1$ defined by $xz = ty$, where t is the coordinate of \mathbb{A}_k^1 and x, y, z are the coordinates of \mathbb{P}_k^2 . Then \underline{X} has a natural log structure \mathcal{M}_X . For example, on the locus where z is invertible, \underline{X} is given by $\text{Spec } k[P_z]$ with P_z a monoid on five generators a, b, c, c', u subject to the relations $c + c' = 0$ and $a + c = b + u$; here \mathcal{M}_X is given by the canonical log structure associated to P_z . Then the projection

$$X \longrightarrow \mathbb{A}_k^1$$

is a log curve, where X is given the log structure above and \mathbb{A}_k^1 is given the log structure defined by the divisor $t = 0$. We see that every fiber above $t \neq 0$ is isomorphic to \mathbb{P}_k^1 with log structure given by the divisor at 0 and ∞ ; the fiber above $t = 0$ is nodal. The n in Theorem 4.7 is equal to 2 for all geometric fibers.

Since our goal is to give a log geometric description of $\overline{\mathcal{M}}_{g,n}$, we would like to express the stability condition purely in terms of log geometry. The following proposition provides the key.

Proposition 4.9 ([26, Prop 1.13]). *With notation as in Theorem 4.7, there is a natural isomorphism*

$$\Omega_{X/S}^1 \longrightarrow \omega_{\underline{X}}(s_1 + \cdots + s_n),$$

where $\omega_{\underline{X}}$ is the dualizing sheaf of \underline{X} .

We therefore make the following definition.

Definition 4.10. Let $f : X \rightarrow S$ be a log curve and for all geometric points \bar{t} of \underline{S} , let $\ell(\bar{t})$ and $n(\bar{t})$ be such that

$$\overline{\mathcal{M}}_{X_{\bar{t}}/\bar{t}} = \mathbb{Z}_{r_1} \oplus \cdots \oplus \mathbb{Z}_{r_{\ell(\bar{t})}} \oplus \mathbb{N}_{s_1} \oplus \cdots \oplus \mathbb{N}_{s_{n(\bar{t})}}.$$

We say f is of type (g, n) if \underline{f} is proper, \underline{X} has genus g , and $n(\bar{t}) = n$ for all \bar{t} . We say f is stable of type of (g, n) if it is of type (g, n) and

$$H^0(\underline{X}_{\bar{t}}, T_{X_{\bar{t}}/\bar{t}}) = 0$$

for all geometric points \bar{t} of \underline{S} .

It is, in fact, true ([26, Prop 1.7]) that if $f : X \rightarrow S$ is a log curve of type (g, n) , then the s_i in each geometric fiber fit together to yield n sections σ_i of \underline{f} . It follows then that every stable log curve of type (g, n) is an n -pointed stable curve of genus g in the classical sense.

Log structures on stable curves

Having now shown that every log curve is naturally a pointed nodal curve, we shift gears and ask the following question: given a stable genus g curve $\underline{f} : \underline{X} \rightarrow \underline{S}$ with n marked points, how many log structures can we put on \underline{X} and \underline{S} so that the associated morphism of log schemes is a log curve with relative characteristic supported on our given n marked points? We begin by sketching the construction of a canonical such log structure.

Lemmas 2.1 and 2.2 of [26] show that to endow \underline{X} and \underline{S} with log structures as desired, it is enough to consider the case when $\underline{S} = \text{Spec } A$ and A is strict Henselian. For every node r_i of the closed fiber of \underline{f} , we can find an étale neighborhood U_i of the points specializing to r_i and a diagram

$$\begin{array}{ccc} U_i & \xrightarrow{\psi_i} & \text{Spec } A[x, y, t]/(xy - t) \\ \downarrow & & \downarrow \pi \\ S & \xrightarrow{\varphi_i} & \text{Spec } A[t] \end{array}$$

which is cartesian. Let $t_i \in A$ be the image of t under the morphism induced by φ_i . Endowing $\text{Spec } A[t]$ with the log structure associated to the morphism $\mathbb{N} \rightarrow A[t]$

sending 1 to t , and $\text{Spec } A[x, y, t]/(xy - t)$ with the log structure associated to $\mathbb{N}^2 \rightarrow A[x, y, t]/(xy - t)$ sending e_1 (resp. e_2) to x (resp. y), we see that (π, Δ) is a morphism of log schemes, where $\Delta : \mathbb{N} \rightarrow \mathbb{N}^2$ is the diagonal map. Pulling back these log structures under φ_i (resp. ψ_i), we obtain log structures \mathcal{L}_i (resp. \mathcal{M}'_i) on S (resp. U_i). Away from the points specializing to r_i , we define a log structure \mathcal{M}''_i as the pullback of \mathcal{L}_i . The log structures \mathcal{M}'_i and \mathcal{M}''_i glue to yield a log structure \mathcal{M}_i on X . Let \mathcal{N} be the log structure on X associated to the divisor defined by the marked points. We let

$$\mathcal{M}_X = \mathcal{M}_1 \oplus_{\mathcal{O}_X^*} \cdots \oplus_{\mathcal{O}_X^*} \mathcal{M}_\ell \oplus_{\mathcal{O}_X^*} \mathcal{N}$$

and

$$\mathcal{M}_S = \mathcal{L}_1 \oplus_{\mathcal{O}_S^*} \cdots \oplus_{\mathcal{O}_S^*} \mathcal{L}_\ell.$$

It is not difficult to see that with these definitions, we have endowed f with the structure of a log curve.

Moreover, a detailed analysis of the proof of Theorem 4.7 shows that this log structure we have just constructed is “minimal” among all possible log structures giving $\underline{X}/\underline{S}$ the structure of a log curve (see 1.8 and Thm 2.3 of [26]):

Theorem 4.11. *Let $\underline{X}/\underline{S}$ be a stable genus g curve with n marked points and let X/S be the log curve obtained by endowing $\underline{X}/\underline{S}$ with the canonical log structure above. If X'/S' is a log curve and $\underline{a} : \underline{S}' \rightarrow \underline{S}$ and $\underline{b} : \underline{X}' \rightarrow \underline{X}$ are morphisms such that $\underline{X}' \simeq \underline{X} \times_{\underline{S}} \underline{S}'$ and such that the divisors of marked points in \underline{X}' are sent scheme-theoretically to the divisors of marked points in \underline{X} , there are unique morphisms a and b of log schemes extending the morphisms \underline{a} and \underline{b} above such that*

$$\begin{array}{ccc} X' & \xrightarrow{b} & X \\ \downarrow & & \downarrow \\ S' & \xrightarrow{a} & S \end{array}$$

is cartesian in the category of fs log schemes.

Definition 4.12. A log curve X/S is called *basic* if it satisfies the universal property in Theorem 4.11.

Moduli

Before discussing moduli of log smooth curves, we begin with some generalities about log structures and stacks. Note that the definition of a log structure is not particular to schemes; indeed, Definition 2.2 makes sense for any ringed topos. We can therefore define log structures on the étale site of a Deligne–Mumford stack or the lisse-étale site of an Artin stack. The notions of fine and fs log structures carry over to this setting as well, so one can speak of fine (or fs) log algebraic stacks.

There is another equivalent way to talk about log structures on stacks; namely, if \mathfrak{X} is a stack over the category of schemes, and $\mathcal{M}_{\mathfrak{X}}$ is a log structure on \mathfrak{X} , then \mathfrak{X} can naturally be viewed as a stack over the category of log schemes. For concreteness, say \mathfrak{X} is a stack over the category of schemes with the étale topology. Then we obtain a category $\tilde{\mathfrak{X}}$ fibered over the category of log schemes by defining $\tilde{\mathfrak{X}}(S, \mathcal{M}_S)$ to be the category whose objects are morphisms

$$f : (S, \mathcal{M}_S) \rightarrow (\mathfrak{X}, \mathcal{M}_{\mathfrak{X}})$$

of log stacks and whose morphisms g from $f \in \tilde{\mathfrak{X}}(S, \mathcal{M}_S)$ to $f' \in \tilde{\mathfrak{X}}(S', \mathcal{M}_{S'})$ are given by diagrams

$$\begin{array}{ccc} (S, \mathcal{M}_S) & \xrightarrow{g} & (S', \mathcal{M}_{S'}) \\ & \searrow f & \swarrow f' \\ & (\mathfrak{X}, \mathcal{M}_{\mathfrak{X}}) & \end{array}$$

One checks that $\tilde{\mathfrak{X}}$ is a stack over the category of log schemes where coverings are given by surjective strict étale morphisms.

Conversely, given any stack \mathcal{Y} over the category of log schemes with the strict étale topology, we obtain a log stack $(\mathcal{Y}', \mathcal{M}_{\mathcal{Y}'})$ over the category of schemes with the étale topology by letting $\mathcal{Y}'(S)$ be the category of pairs (\mathcal{M}_S, ξ) where ξ is an object of $\mathcal{Y}(S, \mathcal{M}_S)$. The log structure $\mathcal{M}_{\mathcal{Y}'}$ is then defined by the following property: if $f : S \rightarrow \mathcal{Y}$ is a morphism which corresponds to the pair (\mathcal{M}_S, ξ) , then $f^* \mathcal{M}_{\mathcal{Y}'} = \mathcal{M}_S$.

It is however important to note that these *two procedures are not inverse to each other*. If we start with a logarithmic stack $(\mathfrak{X}, \mathcal{M}_{\mathfrak{X}})$, and take the result $(\tilde{\mathfrak{X}})'$ of the composite operation, we do not get \mathfrak{X} but rather the stack $LOG_{\mathfrak{X}}$ described in section 6. In order to recover the stack \mathfrak{X} over schemes from a stack $\tilde{\mathfrak{X}}$ over log schemes, it is necessary to distinguish objects similar to the basic log curves of Definition 4.12. Rather than launch into a premature categorical discussion, let us see how this works for log curves. The issue is revisited in sections 11 (with Olsson's terminology of *special log structures*) and 12 (with Kim's terminology of *minimal log structures*).

Let $\overline{\mathcal{M}}_{g,n}^{log}$ be the stack over the category of fs log schemes with the strict étale topology where $\overline{\mathcal{M}}_{g,n}^{log}(S, \mathcal{M}_S)$ is the category of stable log curves of type (g, n) over (S, \mathcal{M}_S) . Let $\mathcal{M}_{g,n}^{bas}$ be the substack of basic stable log curves of type (g, n) . By the above discussion, we obtain a log stack over the category of schemes with the étale topology, which we again denote by $(\mathcal{M}_{g,n}^{bas}, \mathcal{M}_{\mathcal{M}_{g,n}^{bas}})$.

Note that the Deligne-Mumford compactification $\overline{\mathcal{M}}_{g,n}$ carries a natural log structure $\mathcal{M}_{\overline{\mathcal{M}}_{g,n}}$ coming from the simple normal crossing divisor at the boundary. It follows from [7, §2] that $\mathcal{M}_{\overline{\mathcal{M}}_{g,n}}$ can also be described as the log structure which assigns to each stable curve $\underline{X}/\underline{S}$ the basic log structure obtained on \underline{S} . The

discussion following Definition 4.5 above shows that we have a natural morphism

$$F : (\mathcal{M}_{g,n}^{bas}, \mathcal{M}_{\mathcal{M}_{g,n}^{bas}}) \longrightarrow (\overline{\mathcal{M}}_{g,n}, \mathcal{M}_{\overline{\mathcal{M}}_{g,n}})$$

and from Theorem 4.11 we see

Theorem 4.13 ([26, §4]). *The morphism F is an equivalence of log stacks.*

4.14. Back to the big picture

We end by mentioning a type of converse to the philosophic principal mentioned at the beginning of this section. We have seen that since log smoothness includes degenerate objects, log geometry can naturally lead to compactifications; however, it is also generally true that we do not end up with “too many” degenerate objects.

Philosophy. Log geometry controls degenerations.

In higher dimensions, compactifications tend to have unwanted extra components. Log geometry helps to cut down on these components. Let us give some inkling of an idea as to why this should be true. Suppose \mathfrak{X} is an algebraic stack which is irreducible. Suppose we can find a proper algebraic stack $\tilde{\mathfrak{X}}$ with a fine log structure and an open immersion $i : \mathfrak{X} \rightarrow \tilde{\mathfrak{X}}$ such that \mathfrak{X} is the trivial locus of $\tilde{\mathfrak{X}}$.

As we now explain, log geometry provides us with a good method of trying to show that $\tilde{\mathfrak{X}}$ is irreducible as well. If k is separably closed and $x : \text{Spec } k \rightarrow \tilde{\mathfrak{X}}$ is a morphism, then pulling back the log structure on $\tilde{\mathfrak{X}}$ endows $\text{Spec } k$ with a fine log structure. By [27, Lemma 2.10], it follows that this is the log structure associated to a morphism of monoids $P \rightarrow k$, where P is fine. Hence, we have a strict closed immersion of log schemes $j : \text{Spec } k \rightarrow \text{Spec } k[[P]]$, where $\text{Spec } k[[P]]$ is given its canonical log structure. Note that the generic point $\text{Spec } K$ of $\text{Spec } k[[P]]$ carries the trivial log structure. Therefore, if x factors as a morphism of log stacks through j , then we automatically obtain a commutative diagram

$$\begin{array}{ccccc}
 \mathfrak{X} & \xrightarrow{\quad i \quad} & \tilde{\mathfrak{X}} & & \\
 \uparrow y & & \nearrow & & \uparrow x \\
 \text{Spec } K & \longrightarrow & \text{Spec } k[[P]] & \xleftarrow{\quad j \quad} & \text{Spec } k
 \end{array}$$

and hence x is the specialization of the point y of \mathfrak{X} . We see then that the log structure on $\text{Spec } k$ obtained from $\tilde{\mathfrak{X}}$ somehow serves as a compass telling us which way to look in order to find a family degenerating to our given point x of $\tilde{\mathfrak{X}}$.

5. D-semistability and log structures

The main references here are [10, 24, 47].

Convention. Throughout this section, every scheme is over an algebraically closed field. The notation X will be reserved for a *normal crossing variety*. By this we mean a variety for which every closed point $x \in X$ has an étale neighborhood $x \rightarrow U$ which admits an étale map $U \rightarrow \text{Spec} \frac{k[x_1, \dots, x_n]}{(x_1 \cdots x_r)} \rightarrow X$ such that x maps to the point with coordinates $(0, \dots, 0)$. By a standard neighborhood of $x \in X$, we mean an étale neighborhood of x as above. The notation \mathcal{M}_k denotes the log structure of the standard log point $\mathbb{N} \rightarrow k, n \mapsto 0^n$, see example 2.11.

Introduction

To study the geometry of a normal crossing variety X , e.g. to study the deformation theory of X , one would like to ask the following questions:

Question 5.1. Can we embed X into another variety $i : X \rightarrow \mathcal{X}$ as a normal crossing divisor?

Question 5.2. Can we find a semi-stable smoothing of X , i.e. embed $X \rightarrow \text{Spec} k$ in a flat family over a curve $\mathcal{X} \rightarrow C$, in such a way that there exists a diagram

$$\begin{array}{ccccc} X & \longrightarrow & \mathcal{X} & \longleftarrow & \mathcal{X}^* = \mathcal{X} \setminus X \\ \downarrow & & \downarrow & & \downarrow f^* \\ \text{Spec } k & \xrightarrow{0} & C & \longleftarrow & C^* = C \setminus 0 \end{array}$$

where \mathcal{X} is smooth, the squares are cartesian and f^* is smooth?

The answer to these questions are not always yes, because the existence of such maps would imply the existence of certain log structures on X , which in turn would imply intrinsic condition on X , so their existence is not guaranteed.

Example 5.3 (log structure of embedding type). If we can find an embedding as in Question (5.1) above, then $i^*(j_* \mathcal{O}_{\mathcal{X}^*}^\times) \rightarrow \mathcal{O}_X$ defines a log structure \mathcal{M}_X on X , which étale locally has a chart $\mathbb{N}^r \rightarrow \text{Spec} \frac{k[x_1, \dots, x_n]}{(x_1 \cdots x_r)}$ sending the element e_i in the standard basis of \mathbb{N}^r to x_i . This is called a *log structure of embedding type*.

Example 5.4 (log structure of semi-stable type). If we can find a semi-stable smoothing as in Question (5.2) above, then what we have in this case is not only a log structure of embedding type on X , but also a morphism (of sheaf of monoids) $f^b : f^* \mathcal{M}_k \rightarrow \mathcal{M}_X$, which makes X a log smooth variety over the standard log point $(\text{Spec } k, \mathcal{M}_k)$. Étale locally a chart for the log structure on X can be put in the form

$$\left(\text{Spec} \frac{k[x_1, \dots, x_n]}{(x_1 \cdots x_r)}, \mathbb{N}^r, e_i \mapsto x_i, i = 1, \dots, r \right).$$

Modulo the units in the monoid, the morphism of quotient monoids induced by $f^b : f^* \mathcal{M}_k \rightarrow \mathcal{M}_X$ is just the diagonal $\Delta : \mathbb{N} \rightarrow \mathbb{N}^r$. Such a pair $(\mathcal{M}_X, f^b : f^* \mathcal{M}_k \rightarrow \mathcal{M}_X)$ is called a *log structure of semi-stable type on X* .

Remark 5.5. The log structure \mathcal{M}_k on $\text{Spec } k$ can be defined by the pullback of the log structure $(\mathcal{M}_0 = j_* \mathcal{O}_{C^*}^\times \rightarrow \mathcal{O}_C)$ on C , i.e. the log structure defined by the divisor 0 of C . We have an isomorphism $\overline{\mathcal{M}}_k := \mathcal{M}_k/k^\times \cong (\mathcal{M}_0/\mathcal{O}_C^\times)_0 \cong \mathbb{N}$, where the second isomorphism assigns each function to its vanishing order at the point 0 in \mathbb{N} . This gives a geometric interpretation of the standard log point.

Concerning the existence of such log structure on the normal crossing variety X , we have the following theorems ([24], Sec.11):

Theorem 5.6. *Let X be a normal crossing variety over the spectrum of an algebraically closed field, then X can be equipped with a log structure of embedding type iff there exists a line bundle \mathcal{L} on X such that*

$$\mathcal{E}xt_X^1(\Omega_X^1, \mathcal{O}_X)|_D \cong \mathcal{L}|_D$$

where D is the non-smooth locus of X .

Remark 5.7. It is not hard to see $\mathcal{E}xt_X^1(\Omega_X^1, \mathcal{O}_X)|_D$ is a line bundle on D .

Definition 5.8. (see [10, Def. 1.13 and Prop. 2.3]) Let X and D as before. If $\mathcal{E}xt_X^1(\Omega_X^1, \mathcal{O}_X)|_D$ is a trivial on D , then we say that X is *d-semistable*.

Theorem 5.9 (d-semistability). *Let X be a normal crossing variety over the spectrum of an algebraically closed field, then X can be equipped with a log structure over the standard log point, such that the structure morphism is log smooth if and only if X is d-semistable.*

Generalization of these theorems can be found in [47], section 3.

Corollary 5.10. *If X has a semistable smoothing as in Question 5.2, then X is d-semistable.*

In fact, we can put a log structure on X such that it is log smooth over the standard log point by Example 5.4, so we can apply Theorem 5.9

Remark 5.11. Being d-semistable is not equivalent to having a semi-stably smoothing. See [54, Section 3] for counterexamples.

Example 5.12 (a normal crossing variety that is not d-semistable [10]). Let X be the subvariety of \mathbf{P}^3 defined by the product of 4 linear equations $f = L_1 L_2 L_3 L_4 = 0$. It is a normal crossing variety provided the four planes has no points in common. Then D is defined by the homogeneous ideal $(L_i L_j | 1 \leq i < j \leq 4)$, and it is not hard to calculate that $\mathcal{E}xt_X^1(\Omega_X^1, \mathcal{O}_X)|_D \cong \mathcal{O}_D(4)$, which is not trivial. So X is not d-semistable.

Corollary 5.13. *The X in Example 5.12 does not admit a semi-stably smoothing.*

Remark 5.14. If we put X in the 1-dimensional family \mathcal{X} defined by $f + tg = 0$ with parameter t , where g is a smooth quartic, then X is the fiber over $t = 0$ and for generic g , \mathcal{X} over $(t \neq 0)$ is smooth. But the whole space of this family is not smooth: in fact, for a generic g , this family is singular at the 24 points of $D \cap \{g = 0\}$. However, a single blowing up at such a point gives a $\mathbf{P}^1 \times \mathbf{P}^1$. Contract along either ruling gives back a family with parameter t which has one less singularity. If we do this process to all 24 points, we will get a family which is a semi-stable smoothing of \tilde{X} , the blowing-up of X at those 24 points. \tilde{X} is d-semistable. For details, see [10, Rem. 1.14].

Refined analysis of the existence of log structures

Continuing with the notation X, D above, let us analyze the situation. We want to break the job of finding a suitable log structure over the standard log point into 2 steps:

- (1) Put a log structure of embedding type on X .
- (2) See if it is possible to make that log structure semi-stable.

We will see that two related obstructions arise naturally, where the vanishing of the first corresponds to the first step, and the vanishing of the second, which means precisely being d-semistable, allows us to do the second step.

Since étale locally a log structure of embedding type always exists, let us consider the stack \mathcal{G} , which to each $U \in X_{\text{ét}}$ associates the groupoid of log structure of embedding type on U . Using Artin's approximation theorem one can show that any two elements of U are locally isomorphic, which means \mathcal{G} is a gerbe. Since $\text{Aut } \mathcal{G} \cong \mathcal{K}$, where \mathcal{K} is the kernel of the restriction map $\mathcal{O}_X^\times \rightarrow \mathcal{O}_D^\times$, we have:

Proposition 5.15. *There is an obstruction η in $H^2(X_{\text{ét}}, \mathcal{K})$ whose vanishing is equivalent to the existence of a log structure of embedding type on X .*

For the calculation of this obstruction, we state the following result (See [24], Sec.11 and [47], Sec.3):

Proposition 5.16. *In the long exact sequence of cohomology associated to the short exact sequence $1 \rightarrow \mathcal{K} \rightarrow \mathcal{O}_X^\times \rightarrow \mathcal{O}_D^\times \rightarrow 1$, the line bundle $\text{Ext}_X^1(\Omega_X^1, \mathcal{O}_X)|_D$ maps to $-\eta \in H^2(X_{\text{ét}}, \mathcal{K})$.*

Combining these results with the exactness of the long exact sequence, we get Theorem 5.6.

Remark 5.17. General theory tells us if $\eta = 0$, then the set of all log structure of embedding type on X is naturally a torsor under $H^1(X_{\text{ét}}, \mathcal{K})$. In general the set of all log structure of embedding type is only a pseudo torsor under this group.

Suppose $\eta = 0$, then we can put a log structure of embedding type on X , which maps to $(\text{Spec } k, k^\times)$. To go one step further, i.e., to make it a log structure of semistable type, we need a morphism of monoids $f^*\mathcal{M}_k \rightarrow \mathcal{M}_X$, such that in a standard neighborhood of $x \in X$, a chart of the morphism is given by the diagonal $\Delta : \mathbb{N} \rightarrow \mathbb{N}^r$, where r is the number of irreducible components passing through x .

Since $\mathcal{M}_k \cong \mathbb{N} \oplus k^\times$ (non-canonically), and the image of $k^\times \subset \mathcal{M}_k$ is determined by the underlying morphism of schemes. To give the morphism wanted from \mathcal{M}_k to \mathcal{M}_X , we only have to specify the image of an element in \mathcal{M}_k having vanishing order 1.

Now the question becomes a lifting problem for morphism of sheaves in monoids:

$$\begin{array}{ccc} & & \mathcal{M}_X \\ & \nearrow & \downarrow \beta \\ \mathbb{N} \cong f^{-1}\overline{\mathcal{M}}_k & \xrightarrow{\overline{f^b}} & \overline{\mathcal{M}}_X \end{array}$$

where étale locally $\overline{f^b}$ is the diagonal Δ . To lift $\overline{f^b}$ it is equivalent to lift the element $\overline{f^b}(1)$.

Consider the sheaf of all the possible local liftings of $\overline{f^b}(1)$, $T = \beta^{-1}(\overline{f^b}(1))$, then T is a torsor under \mathcal{O}_X^\times . To find a lifting of $\overline{f^b}(1)$, it is equivalent to find a global section of T , i.e. a trivialization of T .

It seems like we have got an obstruction of finding a log structure on X which is semi-stable in $H^1(X_{\text{ét}}, \mathcal{O}_X) = \text{Pic}(X)$. This is, however, not quite true. In fact, what we got is for each log scheme X with a log structure of embedding type, the obstruction of making it semi-stable. And our original question (on the existence of log structure of semi-stable type) allows some ambiguity of choosing the log structure of embedding type \mathcal{M}_X on X . As we said in remark 5.17, in this case the set of all log structure of embedding type on X is an $H^1(X_{\text{ét}}, \mathcal{K})$ -torsor, which implies:

Proposition 5.18. *If $\eta = 0$, i.e. there exists a log structure of embedding type on X . Then there is an obstruction for finding a log structure of semi-stable type on X , $\eta' \in H^1(X_{\text{ét}}, \mathcal{O}_X^\times)/H^1(X_{\text{ét}}, \mathcal{K})$, whose vanishing is equivalent to the existence of such a log structure.*

By the long exact sequence of cohomology, $H^1(X_{\text{ét}}, \mathcal{O}_X^\times)/H^1(X_{\text{ét}}, \mathcal{K})$ embeds into $H^1(D_{\text{ét}}, \mathcal{O}_D^\times) \cong \text{Pic}(D)$. For the calculation of η' as an element of $\text{Pic}(D)$, we state the following proposition (See [47]).

Proposition 5.19. *We have $-\eta' = [\mathcal{E}xt_X^1(\Omega_X^1, \mathcal{O}_X)|_D] \in \text{Pic}(D)$.*

Combining these two proposition, we get Theorem 5.9.

6. Stacks of logarithmic structures

The main reference of this section is [46].

A motivating example

Before introducing the stack LOG_S classifying fine log structures on schemes over a fine log scheme S , constructed by Olsson in [46], let us look at an example, which will give the local covers of LOG_S .

Definition 6.1. Let \underline{X} be a scheme and $r \geq 1$ an integer. A *Deligne-Faltings log structure of rank r on \underline{X}* (abbreviated as a *DF log structure of rank r*) is the following data:

- a sequence L_1, \dots, L_r of line bundles on \underline{X} , and
- a morphism $s_i : L_i \rightarrow \mathcal{O}_{\underline{X}}$ of line bundles, for each i .

Consider the following three categories fibered in groupoids over the category of schemes:

- (1) the category of triples $(\underline{X}, L, s : L \rightarrow \mathcal{O}_{\underline{X}})$ consisting of a scheme \underline{X} and a DF log structure of rank 1 on \underline{X} ;
- (2) the category of pairs $(X, \beta : \mathbb{N} \rightarrow \overline{\mathcal{M}}_X)$ consisting of a fine log scheme X and a morphism of sheaves of monoids β that étale locally lifts to a chart: $\tilde{\beta} : \mathbb{N} \rightarrow \mathcal{M}_X$;
- (3) the quotient stack $[\mathbb{A}^1/\mathbb{G}_m]$, where the quotient is formed with respect to the multiplication action of \mathbb{G}_m on \mathbb{A}^1 .

Lemma 6.2. ([27], complement 1). *These three categories fibered in groupoids are equivalent.*

Let us sketch the proof. Given a DF log structure $(L, s : L \rightarrow \mathcal{O}_{\underline{X}})$ of rank 1 on \underline{X} , define a sheaf of monoids \mathcal{M}' on \underline{X} to be

$$\coprod_{n \geq 0} \underline{\text{Isom}}(\mathcal{O}_{\underline{X}}, L^{\otimes n}),$$

the sheafification of the presheaf that takes U to $\coprod_{n \geq 0} \text{Isom}(\mathcal{O}_U, (L|_U)^{\otimes n})$. It comes with a natural morphism of sheaves of monoids $\mathcal{M}' \rightarrow \mathbb{N}$, where the monoid structure on \mathcal{M}' is induced by

$$(n, a : \mathcal{O} \rightarrow L^{\otimes n}) \cdot (m, b : \mathcal{O} \rightarrow L^{\otimes m}) = (n + m, a \otimes b).$$

The map $s : L \rightarrow \mathcal{O}_{\underline{X}}$ induces a morphism

$$\underline{\text{Isom}}(\mathcal{O}_{\underline{X}}, L^{\otimes n}) \xrightarrow{\otimes^s} \underline{\text{Hom}}(\mathcal{O}_{\underline{X}}, \mathcal{O}_{\underline{X}}) = \mathcal{O}_{\underline{X}}$$

of sheaves, hence giving a pre-log structure on \mathcal{M}' :

$$\mathcal{M}' \rightarrow \mathcal{O}_{\underline{X}}.$$

We take \mathcal{M}_X to be the log structure associated to this pre-log structure \mathcal{M}' . Note that $\mathcal{M}'/\mathcal{O}_X^* \cong \mathbb{N}$, and we define $\beta : \mathbb{N} \rightarrow \overline{\mathcal{M}}_X$ to be the composite

$$\beta : \mathbb{N} \cong \mathcal{M}'/\mathcal{O}_X^* \rightarrow \mathcal{M}_X/\mathcal{O}_X^*.$$

Locally the line bundle L is trivial, and one can choose a trivialization of L , which gives trivializations of all $L^{\otimes n}$. Sending $n \in \mathbb{N}$ to this trivialization defines a section $\mathbb{N} \rightarrow \mathcal{M}'$, and hence a section $\tilde{\beta} : \mathbb{N} \rightarrow \mathcal{M}' \rightarrow \mathcal{M}_X$. One can check that this is a chart.

Conversely, given a fine log structure $(\mathcal{M}_X, \mathcal{M}_X \xrightarrow{\alpha} \mathcal{O}_X)$ on \underline{X} with a morphism $\beta : \mathbb{N} \rightarrow \overline{\mathcal{M}}_X$ that étale locally lifts to a chart $\tilde{\beta} : \mathbb{N} \rightarrow \mathcal{M}_X$, we have a section $\beta(1)$ of $\overline{\mathcal{M}}_X$, and its inverse image under $\pi : \mathcal{M}_X \rightarrow \overline{\mathcal{M}}_X$ is an \mathcal{O}_X^* -torsor, which corresponds to a line bundle L . The composition

$$\pi^{-1}(\beta(1)) \subset \mathcal{M}_X \xrightarrow{\alpha} \mathcal{O}_X$$

gives a morphism of line bundles $s : L \rightarrow \mathcal{O}_X$.

Giving a morphism $\underline{X} \rightarrow [\mathbb{A}^1/\mathbb{G}_m]$ is equivalent to giving a \mathbb{G}_m -torsor (namely a line bundle L) with a \mathbb{G}_m -equivariant morphism to \mathbb{A}^1 :

$$\begin{array}{ccc} Y & \longrightarrow & \mathbb{A}^1 \\ \downarrow & & \downarrow \\ X & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m]. \end{array}$$

This diagram is equivalent to the following one

$$\begin{array}{ccc} Y & \xrightarrow{s} & \mathbb{A}_X^1 \\ \downarrow & & \downarrow \\ X & \longrightarrow & [\mathbb{A}^1/\mathbb{G}_m]_X, \end{array}$$

and the top arrow is \mathbb{G}_m -equivariant, namely a morphism of line bundles $s : L \rightarrow \mathcal{O}_X$. This finishes the proof.

In fact, in the three fibered categories, one can replace \mathbb{N} by \mathbb{N}^r , and rank 1 DF-log structure by rank r DF-log structure, and replace $[\mathbb{A}^1/\mathbb{G}_m]$ by $[\mathbb{A}^r/\mathbb{G}_m^r]$ (which is equivalent to $[\mathbb{A}^1/\mathbb{G}_m]^r$), and they are still equivalent.

More generally, let P be a fine monoid and S a scheme, and let $S[P]$ be the product $S \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}[P]$, which has a fine log structure coming from the chart $P \rightarrow \mathbb{Z}[P]$ (2.8). For an affine S -scheme $\text{Spec } R$, the set of R -points $S[P](R)$ is the set of monoid homomorphisms $\text{Hom}_{\text{mon}}(P, R)$, where R is regarded as a multiplicative monoid. Let P^{gp} be the group associated to P . For any affine S -scheme $\text{Spec } R$, the group $\text{Hom}_{\text{mon}}(P^{\text{gp}}, R) = \text{Hom}_{\text{gp}}(P^{\text{gp}}, R^*)$ acts on the set $\text{Hom}_{\text{mon}}(P, R)$ by pointwise multiplication. This induces an action of the S -group scheme $S[P^{\text{gp}}]$ on $S[P]$. When $S = \text{Spec } k$ for a field k and P is saturated and

torsion-free, the k -group variety $S[P^{\text{gp}}]$ is a torus, and $S[P]$ is a toric variety with respect to this torus action.

We have the following.

Lemma 6.3. ([46], 5.14, 5.15) *The following two categories fibered in groupoids over the category of S -schemes are equivalent:*

- (1) *the category of pairs $(X, \beta : P \rightarrow \overline{\mathcal{M}}_X)$ consisting of a fine log scheme X with a morphism $\underline{X} \rightarrow S$ and a morphism of sheaves of monoids β that fppf locally lifts to a chart: $\tilde{\beta} : P \rightarrow \mathcal{M}_X$;*
- (2) *the quotient stack $\mathcal{S}_P := [S[P]/S[P^{\text{gp}}]]$.*

If in addition P is fs, then one can replace “fppf” by “étale”.

6.4. In fact, the action of $S[P^{\text{gp}}]$ on $S[P]$ extends to an action on the log structure on $S[P]$, and so this log structure descends to a log structure $\mathcal{M}_{\mathcal{S}_P}$ on the stack \mathcal{S}_P (cf. Section 4 for the definition of log structures on stacks), and there is a natural morphism $\pi_P : P \rightarrow \overline{\mathcal{M}}_{\mathcal{S}_P}$ of sheaves of monoids that fppf locally lifts to a chart. This is the universal pair $(\mathcal{M}_{\mathcal{S}_P}, \pi_P)$ on \mathcal{S}_P that induces the equivalence in (6.3) above.

Moreover, for a morphism $h : Q \rightarrow P$ of fine monoids, the induced morphism

$$S[h] : S[P] \rightarrow S[Q]$$

is compatible with the actions of $S[P^{\text{gp}}]$, $S[Q^{\text{gp}}]$ and the homomorphism $S[h^{\text{gp}}] : S[P^{\text{gp}}] \rightarrow S[Q^{\text{gp}}]$, hence it descends to a morphism

$$S(h) : \mathcal{S}_P \rightarrow \mathcal{S}_Q$$

of S -stacks. The map $h : Q \rightarrow P$, regarded as a morphism of constant sheaves, induces a morphism $S(h)^* \mathcal{M}_{\mathcal{S}_Q} \rightarrow \mathcal{M}_{\mathcal{S}_P}$ of log structures, making $S(h)$ into a morphism of S -log stacks.

The stack of log structures

Now we can discuss the stack LOG_S parameterizing fine log structures.

Let S be a fine log scheme. Define LOG_S to be the category with

- objects: morphisms $X \rightarrow S$ of fine log schemes, and
- morphisms: strict morphisms $X \rightarrow Y$ over S .

With the functor $(X \rightarrow S) \mapsto (\underline{X} \rightarrow \underline{S})$ from $LOG_S \rightarrow \text{Sch}_{\underline{S}}$, this defines a fibered category over \underline{S} . One of the main results in [46] is the following.

Theorem 6.5. ([46], 1.1) *LOG_S is an algebraic stack locally of finite presentation over \underline{S} .*

Here for an algebraic stack we use a slightly different definition from [36, 4.1]. Namely the first axiom there that the diagonal is representable, separated and quasi-compact, is replaced by that the diagonal is representable and of finite presentation. In fact the stack LOG_S is not quasi-separated [46, 3.17].

Here are two basic properties of LOG_S .

Proposition 6.6. ([46], 3.19, 3.20) (1). *The natural map $i_S : \underline{S} \rightarrow LOG_S$ corresponding to the identity morphism $S \rightarrow S$ is an open immersion;*
 (2). *The 2-functor*

$$S \mapsto LOG_S : \{\text{fine log schemes}\} \rightarrow \{\text{algebraic stacks}\}$$

preserves fiber product. More precisely, if

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ S' & \longrightarrow & S \end{array}$$

is a Cartesian square of fine log schemes, then the induced diagram

$$\begin{array}{ccc} LOG_{X'} & \longrightarrow & LOG_X \\ \downarrow & & \downarrow \\ LOG_{S'} & \longrightarrow & LOG_S \end{array}$$

is a 2-Cartesian square of algebraic stacks.

What is LOG_S good for?

One can use this stack LOG_S to reinterpret many concepts in log geometry. Note that for a morphism $f : X \rightarrow S$ of fine log schemes, the induced morphism $LOG(f) : LOG_X \rightarrow LOG_S$ is faithful, hence representable.

Definition 6.7. Let P be a property of representable morphisms of algebraic stacks. Then we say that $f : X \rightarrow S$ *has property* $LOG(P)$ if $LOG(f) : LOG_X \rightarrow LOG_S$ has property P . We say that f *has property weak* $LOG(P)$ if the map $\underline{X} \rightarrow LOG_S$ corresponding to the given morphism $f : X \rightarrow S$ has property P .

6.8. Caution: The diagram

$$\begin{array}{ccc} \underline{X} & \xrightarrow{i_X} & LOG_X \\ \downarrow f & & \downarrow LOG(f) \\ \underline{S} & \xrightarrow{i_S} & LOG_S \end{array}$$

does not necessarily commute. It commutes if and only if f is strict. In [49, Section 2] a device is introduced in order to fix this issue, using stacks of diagrams of logarithmic structures.

Recall from (3.10) the notion of log smoothness and log étaleness.

Theorem 6.9. *For a morphism $f : X \rightarrow S$ of fine log schemes, f is LOG smooth (resp. LOG étale) if and only if f is log smooth (resp. log étale), if and only if f is weakly LOG smooth (resp. weakly LOG étale).*

This is part of [46, 4.6].

6.10. Another application is the following. Consider the deformation problem for a log smooth integral morphism $f_0 : X_0 \rightarrow B_0$ of fine log schemes and a strict square-zero thickening $B_0 \rightarrow B$ defined by an ideal $J \subset \mathcal{O}_B$. In (3.20) we gave the relation between this deformation problem and the cohomology groups of the log tangent bundle T_{X_0/B_0} . The stack LOG_S provides another way of thinking of this problem.

By (6.9), the log smooth morphism $f_0 : X_0 \rightarrow B_0$ induces a representable smooth morphism $\underline{X}_0 \rightarrow LOG_{B_0}$, denoted \mathcal{L}_{f_0} , and the deformation problem

$$\begin{array}{ccc} X_0 & \hookrightarrow & X \\ f_0 \downarrow & & \downarrow f \\ B_0 & \hookrightarrow & B \end{array}$$

is equivalent to the following

$$\begin{array}{ccc} \underline{X}_0 & \dashrightarrow & \underline{X} \\ \mathcal{L}_{f_0} \downarrow & & \downarrow \mathcal{L}_f \\ LOG_{B_0} & \longrightarrow & LOG_B. \end{array}$$

The solution to this deformation problem is the cohomology groups of the *ordinary tangent bundle* $T_{\underline{X}_0/LOG_{B_0}}$, therefore, theorem (3.20) holds with T_{X_0/B_0} replaced by $T_{\underline{X}_0/LOG_{B_0}}$. In fact, we have $\Omega_{X_0/B_0}^1 \cong \Omega_{\underline{X}_0/LOG_{B_0}}$ (cf. ([49], 3.8)).

See section 7 for the general deformation theory of log schemes.

Local structure of LOG_S .

For a fine log scheme S , the relation between the quotient stacks \mathcal{S}_P and LOG_S is that, the stack LOG_S can be covered by the relative versions of the \mathcal{S}_P 's.

Let $u : U \rightarrow S$ be a strict morphism of fine log schemes, such that the underlying morphism \underline{u} is étale. We will just say that u is an étale strict morphism, if there is no confusion. Let $\beta : Q \rightarrow \mathcal{M}_U$ be a chart, and let $h : Q \rightarrow P$ be a morphism of fine monoids. The chart β induces a strict morphism $U \rightarrow \underline{\mathcal{S}}[Q]$, which we also denote by β .

Let \mathcal{S}_P be the quotient stack $[\underline{\mathcal{S}}[P]/\underline{\mathcal{S}}[P^{\text{gp}}]]$ with the natural fine log structure $\mathcal{M}_{\mathcal{S}_P}$ in (6.4), and let $\underline{\mathcal{S}}_P$ be the underlying stack. Consider the 2-commutative

diagram

$$\begin{array}{ccc} \underline{\mathcal{S}}_P \times_{\underline{\mathcal{S}}_Q} \underline{U} & \xrightarrow{pr_2} \underline{U} & \xrightarrow{\underline{\beta}} \underline{\mathcal{S}}[Q] \\ pr_1 \downarrow & & \downarrow \pi \\ \underline{\mathcal{S}}_P & \xrightarrow{\underline{\mathcal{S}}(h)} & \underline{\mathcal{S}}_Q. \end{array}$$

Let Z be the \underline{U} -log stack with underlying stack $\underline{Z} = \underline{\mathcal{S}}_P \times_{\underline{\mathcal{S}}_Q} \underline{U}$ and the inverse image log structure $\mathcal{M}_Z = pr_1^* \mathcal{M}_{\underline{\mathcal{S}}_P}$. Applying pr_1^* to the morphism of log structures $\underline{\mathcal{S}}(h)^* \mathcal{M}_{\underline{\mathcal{S}}_Q} \rightarrow \mathcal{M}_{\underline{\mathcal{S}}_P}$ (cf. (6.4)), noting that $\pi \circ \beta : \underline{U} \rightarrow \underline{\mathcal{S}}_Q$ is strict, we obtain a morphism $pr_2^* \mathcal{M}_{\underline{U}} \rightarrow \mathcal{M}_Z$, making pr_2 into a morphism of log stacks $pr_2 : Z \rightarrow \underline{U}$. This gives a morphism

$$\underline{Z} \rightarrow LOG_{\underline{U}} \rightarrow LOG_{\underline{\mathcal{S}}_Q}.$$

Proposition 6.11. ([46], 5.25) *For any fine log scheme S , the natural morphism*

$$\coprod_{(U, \beta, h)} \underline{\mathcal{S}}_P \times_{\underline{\mathcal{S}}_Q} \underline{U} \rightarrow LOG_S$$

is a representable étale surjection, where the disjoint union is taken over the isomorphism classes of all triples (U, β, h) consisting of an étale strict morphism $U \rightarrow S$, a chart $Q \xrightarrow{\beta} \mathcal{M}_U$, and a morphism $h : Q \rightarrow P$ of fine monoids, for some fine monoids P and Q .

7. Log deformation theory in general

The main reference here is [49].

As is well known, the general deformation theory of schemes and morphisms of schemes is not as easy as in the smooth case. To understand deformation theory of general morphisms, one has to use the full power of the cotangent complex, see [19, 20]. In log geometry, one can generalize it to get a reasonable theory of logarithmic cotangent complex.

This log cotangent complex will be compatible with the usual cotangent complex when the morphism in question is strict and is also compatible with log smooth deformation theory for log smooth morphisms. Basically, this is an application of the deformation theory of representable morphisms to algebraic stacks ([50]) to the classifying morphisms from the underlying scheme of X to the stack LOG_Y ([46], see also section 6 of this chapter.)

Convention. We will focus on the category of fine log schemes. For a log scheme X , \underline{X} means the underlying scheme of X .

Remark 7.1. We will work with the category $D'(\underline{X}_{\text{ét}})$ and similar categories, and one can talk about distinguished triangles and Ext's in these categories. For relevant definitions, see [49] and [50].

Our presentation here follows [49]. An alternative approach to the cotangent complex due to Gabber is also explained in [49, Section 8].

The Log Cotangent Complex

In Section 6, an Artin stack LOG_Y is defined for a log scheme Y . It has the property that morphisms of log schemes $X \rightarrow Y$ are equivalent to morphisms $\underline{X} \rightarrow LOG_Y$. Thus one may interpret deformations of a morphism of log schemes $X \rightarrow Y$ as deformations of the associated representable morphism $\underline{X} \rightarrow LOG_Y$. In [50], the deformation theory of representable morphisms of stacks was studied in detail. As an application of this theory, one makes the following definition:

Definition 7.2. For a morphism of log schemes $f : X \rightarrow Y$, the *logarithmic cotangent complex of f* is the complex $L_f = L_{\underline{X}/LOG_Y}$, where the right hand side is the cotangent complex of the morphism $\underline{X} \rightarrow LOG_Y$ defined in Section 6.

Remark 7.3. One should think about L_f as an object of the category $D'_{qcoh}(\underline{X}_{ét})$. In the above definition, the right hand side is an object of the category $D'_{qcoh}(\underline{X}_{lis-ét})$. As the restriction functor $D'_{qcoh}(\underline{X}_{lis-ét}) \rightarrow D'_{qcoh}(\underline{X}_{ét})$ is an equivalence of categories, no information of the cotangent complex would be lost.

Basic Properties

For every morphism of fine log schemes $f : X \rightarrow Y$ the log cotangent complex is a projective system

$$L_f = (\cdots \rightarrow L_{\bar{f}}^{\geq -n-1} \rightarrow L_{\bar{f}}^{\geq -n} \rightarrow \cdots \rightarrow L_{\bar{f}}^{\geq 0})$$

where each $L_{\bar{f}}^{\geq -n}$ is an essentially constant ind-object in $D^{[-n,0]}(\mathcal{O}_X)$ (The derived category of \mathcal{O}_X -modules supported in $[-n, 0]$).

The log cotangent complex L_f has the following properties:

- (1) For any $n \geq 0$, the natural map $\tau_{\geq -n} L_{\bar{f}}^{\geq -n-1} \rightarrow L_{\bar{f}}^{\geq -n}$ is an isomorphism.
- (2) If f is strict, then the system $(\tau_{\geq -n} L_{\bar{f}'})$ represents L_f , where $L_{\bar{f}'}$ is the usual cotangent complex of the underlying morphism of schemes f' .
- (3) If $f : X \rightarrow Y$ is log smooth, then the sheaf of log differentials $\Omega_{X/Y}^1$ represents L_f .
- (4) If

$$\begin{array}{ccc} X' & \xrightarrow{a} & X \\ g \downarrow & & \downarrow f \\ Y' & \xrightarrow{b} & Y \end{array}$$

is a commutative diagram of fine log schemes, then there is a natural map

$$a^* L_f \rightarrow L_g$$

which is an isomorphism if the square above is cartesian and f is log flat. Furthermore, if the composite $X' \rightarrow Y' \rightarrow Y$ satisfies the condition (T) below, then the map

$$g^*L_b \oplus a^*L_f \rightarrow L_{bg}$$

is also an isomorphism.

(5) Given a composite

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

satisfying condition (T) below, there is a natural map

$$L_f \rightarrow f^*L_g[1]$$

making the resulting triangle

$$(7.4) \quad f^*L_g \rightarrow L_{gf} \rightarrow L_f \rightarrow f^*L_g[1]$$

distinguished.

Remark 7.5. In (4) and (5) above, f^* , a^* , g^* should be understood in the derived sense.

Remark 7.6. One might hope for a theory of log cotangent complex in which every triangle (7.4) is distinguished. This is unfortunately not the case - an example due to W. Bauer is given in [49, Section 7].

On the other hand, Gabber has shown (see [49, Section 8]) that if one loosens the requirement 3, then one can obtain a theory of log cotangent complexes for which one has a distinguished triangle (7.4) for all composites $X \xrightarrow{f} Y \xrightarrow{g} Z$.

The Condition (T) mentioned above is the following:

There exists a family of commutative diagrams

$$\begin{array}{ccccccc}
 X_i & \xrightarrow{\pi_{X_i}} & X \times_Y Y_i & \longrightarrow & X \times_Z Z_i & \longrightarrow & X \\
 & \searrow & \downarrow & & \downarrow & & \downarrow f \\
 & & Y_i & \xrightarrow{\pi_{Y_i}} & Y \times_Z Z_i & \longrightarrow & Y \\
 & & & \searrow & \downarrow & & \downarrow g \\
 & & & & Z_i & \xrightarrow{\pi_{Z_i}} & Z
 \end{array}$$

such that

- (1) The underlying schemes of X_i, Y_i, Z_i are all affine.
- (2) The π 's are all strict, and their underlying morphisms are flat and locally of finite presentation.
- (3) The underlying family of morphisms of schemes of $\{X_i \rightarrow X\}$ is jointly surjective.

(4) There exists charts

$$\beta_{X_i} : Q_{X_i} \rightarrow \mathcal{M}_{X_i}, \beta_{Y_i} : Q_{Y_i} \rightarrow \mathcal{M}_{Y_i}, \beta_{Z_i} : Q_{Z_i} \rightarrow \mathcal{M}_{Z_i}$$

and injective maps

$$Q_{Z_i} \rightarrow Q_{Y_i} \rightarrow Q_{X_i}$$

compatible with the morphisms f_i, g_i and

$$\mathrm{Tor}_{\mathcal{O}_{Z_i} \otimes_{\mathbb{Z}[Q_{Z_i}]} \mathbb{Z}[Q_{Y_i}]}^j(\mathcal{O}_{Z_i} \otimes_{\mathbb{Z}[Q_{Z_i}]} \mathbb{Z}[Q_{X_i}], \mathcal{O}_{Y_i}[G]) = 0 \text{ for all } j > 0.$$

Here $G := \mathrm{Coker}(Q_{Z_i}^{gp} \rightarrow Q_{Y_i}^{gp})$ and $\mathcal{O}_{Y_i}[G]$ is viewed as an $\mathcal{O}_{Z_i} \otimes_{\mathbb{Z}[Q_{Z_i}]} \mathbb{Z}[Q_{Y_i}]$ -algebra via the map

$$\mathcal{O}_{Z_i} \otimes_{\mathbb{Z}[Q_{Z_i}]} \mathbb{Z}[Q_{Y_i}] \rightarrow \mathcal{O}_{Y_i}[G], t \otimes e_q \mapsto g_i^*(t) \beta_{Q_{Y_i}}(q) \cdot \bar{q}$$

where \bar{q} denotes the image of q in G .

Deformation Theory of Log Schemes in General

In this section, we explain the relation between the log cotangent complex and deformation theory of log schemes. Let $f : X \rightarrow Y$ be a morphism of fine log schemes and let I be a quasi-coherent sheaf on \underline{X} . Define a Y -extension of X by I to be a commutative diagram of log schemes

$$\begin{array}{ccc} X & \xrightarrow{j} & X' \\ f \downarrow & \swarrow f' & \\ Y & & \end{array}$$

where j is a strict closed immersion defined by a square-zero ideal, together with an isomorphism $\epsilon_j : I \cong \mathrm{Ker}(\mathcal{O}_{X'} \rightarrow \mathcal{O}_X)$. The set of Y -extensions of X by I forms, in a natural way, a category $\underline{\mathrm{Exal}}_Y(X, I)$. Let $\mathrm{Exal}_Y(X, I)$ be the set of isomorphism classes of this category.

There is a tautological equivalence of categories (see [50, Problem 1] for the meaning of the right hand side):

$$\underline{\mathrm{Exal}}_Y(X, I) \cong \underline{\mathrm{Exal}}_{\mathrm{LOG}_Y}(\underline{X}, I).$$

Hence, by [50, Theorem 1.1] and our definition of L_f , we obtain the following result:

Theorem 7.7. ([49, Theorem 5.2]) *There is a natural bijection*

$$\mathrm{Exal}_Y(X, I) \cong \mathrm{Ext}^1(L_f, I).$$

It is precisely the theorem above that guarantees that general deformation theory is controlled by our logarithmic cotangent complex.

Definition 7.8. Let $j_0 : Y_0 \hookrightarrow Y$ be an strict closed immersion of fine log schemes defined by a square-zero ideal $I \subset \mathcal{O}_Y$, and let $f_0 : X_0 \rightarrow Y_0$ be a LOG flat morphism (Definition 6.7). A *log flat deformation of X_0 to Y* is a cartesian square

$$\begin{array}{ccc} X_0 & \xrightarrow{j} & X \\ f_0 \downarrow & & \downarrow f \\ Y_0 & \xrightarrow{j_0} & Y \end{array}$$

with f LOG flat.

To give a log flat deformation as above is equivalent to give a 2-commutative diagram

$$\begin{array}{ccc} \underline{X}_0 & \xrightarrow{j} & \underline{X} \\ \mathcal{L}_{f_0} \downarrow & & \downarrow \mathcal{L}_f \\ \underline{LOG}_{Y_0} & \xrightarrow{j_0} & \underline{LOG}_Y \end{array}$$

with \mathcal{L}_f flat. Thus from ([50], 1.4) we obtain the following:

Theorem 7.9. *Let J denote the ideal of \underline{LOG}_{Y_0} in \underline{LOG}_Y . Then*

- (1) *There exists a canonical class $o \in \text{Ext}^2(L_{f_0}, \mathcal{L}_{f_0}^* J)$ whose vanishing is equivalent to the existence of a log flat deformation of X_0 to Y .*
- (2) *If $o = 0$, then the set of isomorphism classes of log flat deformations of X_0 to Y is naturally a torsor under $\text{Ext}^1(L_{f_0}, \mathcal{L}_{f_0}^* J)$.*
- (3) *The automorphism group of any log flat deformation of X_0 to Y is canonically isomorphic to $\text{Ext}^0(L_{f_0}, \mathcal{L}_{f_0}^* J)$.*

This theorem gives an answer to the question of general deformation theory of log schemes.

Deformations of morphisms

As in the case of usual schemes, once one understands deformations of log schemes, one obtains a solution to the related problem of deformations of a log morphisms.

We are given a commutative diagram of solid arrows

$$(7.10) \quad \begin{array}{ccccc} X_0 & \xrightarrow{i} & X & & \\ & \searrow f_0 & & \searrow f & \\ & & Y_0 & \xrightarrow{j} & Y \\ & \searrow h_0 & \downarrow g_0 & & \downarrow g \\ & & Z_0 & \xrightarrow{k} & Z \end{array}$$

where i, j, k are strict closed immersion defined by square-zero ideal sheaves I, J, K living on X, Y, Z respectively. The question is to find a dotted arrow f fitting in the diagram. To nail down f , we need some more data.

The morphisms h and g induce morphisms $w : h_0^*K \rightarrow I$ and $v : g_0^*K \rightarrow J$. Assume given a morphism $u : f_0^*J \rightarrow I$ such that the composite

$$h_0^*K = f_0^*g_0^*K \rightarrow f_0^*J \rightarrow I$$

is equal to w .

What we want to find is an f fitting in the diagram (7.10) such that the morphism $f_0^*J \rightarrow I$ induced by f is equal to u . This can also be solved by the logarithmic cotangent complex.

Theorem 7.11. *In the situation above, assume in addition that u induces a map $\mathcal{L}_{f_0}^*J' \rightarrow I$, where J' is the ideal of LOG_{Y_0} in LOG_Y , then there is a canonical class $o \in \text{Ext}^1(f_0^*L_{Y_0/Z_0}, I)$ which vanishes iff there exists f fitting in the diagram (7.10) such that the morphism $f_0^*J \rightarrow I$ induced by f is equal to u . If $o = 0$, then the set of such maps f is a torsor under the group $\text{Ext}^0(f_0^*L_{Y_0/Z_0}, I)$.*

For a proof of this theorem, see [49], Theorem 5.9.

8. Rounding

The main reference for this section is [30]. We have benefitted from a lecture by A. Ogus, who reported on results in [43].

What is rounding?

The process of “rounding”, in its most basic form, produces a manifold *with corners* from a *smooth* analytic space with a normal crossings divisor. So the corners are not rounded but rather the opposite: they are created. On the other hand, these corners are rather round and shapely. Also, anybody who has seen the construction under any name and hears the name “rounding” immediately knows what this is about. Evidently then, even though “rounding” might be something of a misnomer, it is a very good name. The origin of the name seems to be in work of Kajiwara, Nakayama and Ogus [23, 43].

In various moduli problems, the rounding of the moduli space often has a more natural topological interpretation than the moduli space itself. A good example is the Deligne-Mumford moduli space $\overline{\mathcal{M}}_{g,n}$ of marked nodal curves, whose boundary is a normal crossings divisor. The “interior” $\mathcal{M}_{g,n}$ can be described topologically as a quotient of Teichmüller space by the appropriate mapping class group. There is a natural generalization of Teichmüller space involving 2-manifolds decorated with circles, due to Harvey [18]; the analogous quotient yields a topological description of the *rounding* of $\overline{\mathcal{M}}_{g,n}$, rather than the moduli space $\overline{\mathcal{M}}_{g,n}$ itself.

Another example is that of twisted curves, discussed in section 11. A twisted curve is an algebraic stack with a log structure, so it is a bit exotic. But its rounding is a good old topological space. A similar example occurs in current work of one of us (Gillam): the relative Hilbert stack of a marked Riemann surface can be defined algebraically using Jun Li style expansions, but it is not representable (except in some trivial cases). However its rounding is a topological space (even a manifold with corners) which is relatively easy to describe. A similar phenomenon occurs in many moduli problems involving expansions, discussed briefly below.

The topological preeminence of the rounding of moduli spaces might ultimately be traced back to the preference in topology for operations involving real codimension one subspaces (e.g. connected sum of manifolds) as opposed to the algebro-geometric preference for complex codimension one operations (e.g. pushout of two smooth varieties along a common divisor). From this point of view, one might think of log geometry as an attempt to speak algebraically about various “real codimension one” phenomena.

The oriented real blowup

The most general rounding operation is the *Kato–Nakayama logarithmic space* associated to a log analytic space [30]. In the basic example of a smooth analytic space with log structure from a normal crossings divisor, the Kato–Nakayama space can be described in terms of *oriented real blowup*, which is a relatively simple rounding operation that can be described as follows.

Suppose X is a topological space, $\pi : L \rightarrow X$ is a complex line bundle, and $s : X \rightarrow L$ is a section of π . Locally on X we can choose a trivialization $(\pi, \phi) : L \rightarrow X \times \mathbb{C}$ and consider the subspace

$$B_{L,s,\phi} X := \left\{ l \in L : |\phi(l)| \cdot (\phi s \pi)(l) = \phi(l) \cdot |(\phi s \pi)(l)| \right\}$$

of L . A continuous function $u : X \rightarrow \mathbb{C}^*$ yields a new trivialization $(\pi, u \cdot \phi)$, where $(u \cdot \phi)(l) := (u\pi)(l)\phi(l)$. The key observation is that $B_{L,s,\phi} X = B_{L,s,u \cdot \phi} X$, so the subspace $B_{L,s,\phi} X$ is independent of the choice of ϕ , hence one can define a subspace $B_{L,s} X \subseteq L$ by defining it locally on X using a trivialization, then gluing the locally defined subspaces. From the local picture using a trivialization, it is clear that the subspace $B_{L,s} X$ contains the zero section and $L|_{Z(s)}$ (where $Z(s) \subseteq X$ is the zero locus of s) and is invariant under the $\mathbb{R}_{>0}$ action on L inherited from the full \mathbb{C}^* scaling action. We let $B_{L,s}^* X$ be the complement of the zero section in $B_{L,s}$ and we call

$$\text{Blo}_{L,s} X := (B_{L,s}^* X) / \mathbb{R}_{>0}$$

the *oriented real blowup* of X along (L, s) .

The space $\text{Blo}_{L,s} X$ is a closed subspace of the oriented circle bundle $S^1 L := L^* / \mathbb{R}_{>0}$ associated to L and is, in particular, proper over X . The projection

$\tau : \text{Blo}_{L,s} X \rightarrow X$ is an isomorphism away from $Z(s)$ and $\tau^{-1}(Z(s))$ is oriented circle bundle $S^1 L|_{Z(s)}$. The spaces $\text{B}_{L,s} X$ and $\text{Blo}_{L,s} X$ are natural under pulling back line bundles and sections.

If X is an analytic space and $D \subseteq X$ is a Cartier divisor, then D determines a line bundle $\mathcal{O}_X(D)$ together with a section s whose zero locus is D . In this situation, we will write $\text{B}_D X$, $\text{Blo}_D X$, etc. and speak of the *oriented real blowup of X along D* . The space $\text{Blo}_D X$ inherits a differentiable structure from its inclusion in $S^1 \mathcal{O}_X(D)$.

The basic example to keep in mind is the oriented real blowup $\text{Blo}_0 \mathbb{C}$ of the complex plane \mathbb{C} at the origin. The origin is the zero locus of the identity map $\text{Id} : \mathbb{C} \rightarrow \mathbb{C}$, hence

$$\begin{aligned} \text{Blo}_0 \mathbb{C} &= \{(z, Z) \in \mathbb{C} \times \mathbb{C}^* : |z|Z = z|Z|\} / \mathbb{R}_{>0} \\ &= \{(z, Z) \in \mathbb{C} \times S^1 : |z|Z = z\} \\ &\cong \mathbb{R}_{\geq 0} \times S^1, \end{aligned}$$

where the last isomorphism from $\mathbb{R}_{\geq 0} \times S^1$ is given by $(\lambda, Z) \mapsto (\lambda Z, Z)$. Evidently $\text{Blo}_0 \mathbb{C}$ is a half-infinite annulus whose boundary S^1 is the exceptional locus of $\tau : \text{Blo}_0 \mathbb{C} \rightarrow \mathbb{C}$ (the fiber over the origin).

The Kato–Nakayama space

Let (X, \mathcal{M}_X) be a fine and saturated logarithmic analytic space.

Definition 8.1. [[30, 1.2]] We define its *canonical rounding*, or *Kato–Nakayama space*, denoted X^{log} , as the space whose points are pairs (x, F) where $x \in X$ and $F : \mathcal{M}_{X,x} \rightarrow S^1$ is a monoid homomorphism satisfying $F(u) = u(x)/|u(x)|$ for every $u \in \mathcal{O}_{X,x}^* \subseteq \mathcal{M}_{X,x}$.

This space has a natural topology. Let us describe the topology in the special case where \mathcal{M}_X is the canonical log structure associated to a Cartier divisor $D \subseteq X$, see Example 2.7. Locally on X we can find $f_1, \dots, f_n \in \mathcal{M}_X(X)$ which, together with the units, generate \mathcal{M}_X . The map

$$\begin{aligned} X^{\text{log}} &\rightarrow X \times (S^1)^n \\ (x, F) &\mapsto (x, F(f_{1,x}), \dots, F(f_{n,x})) \end{aligned}$$

is then easily seen to be a monomorphism onto a closed subset of $X \times (S^1)^n$, so we give X^{log} the subspace topology so that this is a closed embedding. Since one can check easily that this topology does not depend on the choice of generators f_1, \dots, f_n , the locally defined topologies glue to a topology on X^{log} making the projection $\tau : X^{\text{log}} \rightarrow X$ given by $\tau(x, F) := x$ a proper map.

8.2. Relating Kato–Nakayama spaces to oriented blowups

There is a morphism

$$\begin{aligned} \phi : X^{\log} &\rightarrow \mathrm{Blo}_D X \\ (x, F) &\mapsto (x, f \mapsto F(\bar{f})) \end{aligned}$$

of topological spaces over X which requires a little explanation. Here $f \in S^1 \mathcal{O}_X(-D)|_x$ is in the circle bundle associated to the fiber $\mathcal{O}_X(-D)|_x \cong \mathbb{C}$ and $\bar{f} \in \mathcal{O}_{X,x}(-D)$ is a lifting of f to the stalk (one shows that any such \bar{f} is actually in $\mathcal{M}_{X,x}$ and that $F(\bar{f})$ does not depend on this choice of lifting \bar{f}). If we use the identification

$$S^1 \mathcal{O}_X(D) = \mathrm{Hom}_{S^1}(S^1 \mathcal{O}_X(-D), S^1),$$

then we can think of ϕ as a map from X^{\log} to $S^1 \mathcal{O}_X(D)$; one then shows that this ϕ is continuous and that ϕ factors through $\mathrm{Blo}_D X \subseteq S^1 \mathcal{O}_X(D)$.

When X is a smooth analytic space and D is a smooth divisor, the map ϕ is easily seen to be an isomorphism since one can reduce to the case $(X, D) = (\mathbb{C}, 0)$ on formal grounds. Slightly more generally, if X is smooth, but D is only a normal crossings divisor, then locally we can write D as a union of smooth divisors D_1, \dots, D_i which look like the first i coordinate hyperplanes in \mathbb{C}^n ($n = \dim X$), and we can define a variant of the oriented real blowup

$$\mathrm{Blo}'_D X := (\mathrm{Blo}_{D_1} X) \times_X \cdots \times_X (\mathrm{Blo}_{D_i} X)$$

and a map $\phi : X^{\log} \rightarrow \mathrm{Blo}'_D X$. In this local picture, the log structure \mathcal{M}_X is the direct sum (in the category of log structures) of the log structures \mathcal{M}_X^j from D_1, \dots, D_i , the associated Kato–Nakayama space X^{\log} is the fibered product over X of the $(X, \mathcal{M}_X^j)^{\log}$, and ϕ is just the fibered product over X of the previously constructed isomorphisms $\phi_j : (X, \mathcal{M}_X^j)^{\log} \rightarrow \mathrm{Blo}_{D_j} X$. The locally defined variants can be glued to define a global variant $\mathrm{Blo}'_D X$ of the oriented real blowup and an isomorphism $\phi : X^{\log} \cong \mathrm{Blo}'_D X$ of topological spaces over X .

8.3. Topology, cohomology, and the Kato–Nakayama space

Locally, if $X = \mathbb{C}^n$ and D is the union of the first i coordinate hyperplanes, then D is the zero locus of $(z_1, \dots, z_n) \mapsto z_1 \cdots z_i \in \mathbb{C}$ and we have

$$\begin{aligned} \mathrm{Blo}_D X &= \{(z_1, \dots, z_n, Z) \in \mathbb{C}^n \times S^1 : |z_1 \cdots z_i| Z = z_1 \cdots z_i\} \\ \mathrm{Blo}'_D X &= \{(\bar{z}, \bar{Z}) \in \mathbb{C}^n \times (S^1)^i : |z_j| Z_j = z_j \text{ for } j = 1, \dots, i\}. \end{aligned}$$

In the general normal crossings divisor situation, the fiber of $\tau : X^{\log} \rightarrow X$ over a point $x \in X$ is naturally identified with

$$S^1 N_{D_1/X}|_x \times \cdots \times S^1 N_{D_i/X}|_x,$$

where D_1, \dots, D_i are the branches of D containing x . When $\dim X = n$, a point $y \in \tau^{-1}(x)$ has a neighborhood diffeomorphic to a neighborhood of the origin in $\mathbb{R}_{\geq 0}^i \times \mathbb{R}^{2n-i}$. (Note $i \leq n$, so the depth of the corners in a Kato–Nakayama space is somewhat constrained.) Recall that the *topology* near the origin only depends

on whether $i > 0$, but the differentiable structure depends on the actual value of i . In particular, the topological boundary of the manifold X^{\log} is given by $\tau^{-1}(D)$, and this manifold with boundary is homotopy equivalent to its interior, so $H^*(X^{\log}) = H^*(X \setminus D)$.

The topology of morphisms of Kato–Nakayama spaces is also very nice, as shown by the following beautiful general result, see [43, Theorem 0.3]:

Theorem 8.4. *Let $f : X \rightarrow Y$ be a log smooth and integral morphism of fine log analytic spaces. Then the associated map $X^{\log} \rightarrow Y^{\log}$ is a topological submersion.*

In fact, Nakayama and Ogus prove a more general result, replacing integrality by K. Kato’s notion of *exact morphisms*, see [27, Definition 4.6].

The fact that the topology is nice suggests that one expects nice cohomological implications. This is indeed the original motivation leading Kato and Nakayama to define X^{\log} , see [30, Theorem 0.2 (1)]:

Theorem 8.5. *Let (X, M) be a fine and saturated log scheme with X of finite type over \mathbb{C} . Let F be a constructible sheaf on the log-étale site $X_{\log\text{-ét}}$, and let F^{\log} be its pullback to the topological space X^{\log} . Then for all $q \in \mathbb{Z}$ we have*

$$H^q(X_{\log\text{-ét}}, F) = H^q(X^{\log}, F^{\log}).$$

Very strong results hold true for de Rham cohomology. In fact the Kato–Nakayama space is a model for the log de Rham cohomology of X in the sense that

$$H^*(X^{\log}, \mathbb{C}) = \mathbb{H}^*(X, \wedge^{\bullet} \Omega_{(X, M)})$$

under mild assumptions on X . We discuss this in Section 9 below.

Kato–Nakayama spaces of expanded pairs

Given a pair (X, D) consisting of a smooth variety X over \mathbb{C} with a smooth divisor $D \subseteq X$, the notion of an *expanded pair* $t : \mathcal{X} \rightarrow B$ over a base B arises in various relative curve counting theories. The fiber of t over a point $b \in B$ always looks like

$$X[n]_0 = X \prod_D \Delta_1 \prod_D \cdots \prod_D \Delta_n$$

(for an appropriate n), where $\Delta_i = \mathbb{P}(N_{D/X} \oplus \mathcal{O}_D)$ is a \mathbb{P}^1 bundle over X . Both \mathcal{X} and B have natural log structures making t a log smooth map of log schemes. The fiber of $t^{\log} : \mathcal{X}^{\log} \rightarrow B^{\log}$ over a point $c \in \tau_B^{-1}(b)$ looks like

$$X^{\log} \prod_{c_1: S^1 N_{D/X} \cong S^1 N_{D/\Delta_1}} \Delta_1^{\log} \cdots \prod_{c_n: S^1 N_{D/\Delta_{n-1}} \cong S^1 N_{D/\Delta_n}} \Delta_n^{\log},$$

where the choice of $c \in \tau^{-1}(b) \cong (S^1)^n$ determines the choice of orientation reversing S^1 bundle isomorphisms c_1, \dots, c_n . Here each Δ_j has the log structure

from the two copies of D , and Δ_j^{\log} is a cylinder bundle over D (better: an I -bundle over D^{\log}).

The action of $(\mathbb{C}^*)^n$ on $X[n]_0$ given by scaling the fibers of the \mathbb{P}^1 bundles Δ_i is an action by isomorphisms of log schemes, so it lifts to an action on Kato–Nakayama spaces. This lifted action is nontrivial on B^{\log} as the $(S^1)^n$ factor of $(\mathbb{C}^*)^n$ acts simply transitively on $\tau_B^{-1}(b) \cong (S^1)^n$. In the usual moduli problems involving expansions, the isotropy group of a point b involves elements of $(\mathbb{C}^*)^n$ such that the induced action on $X[n]_0$ respects a map from a curve to $X[n]_0$, a subscheme of $X[n]_0$, etc., and this isotropy is usually required to be finite to have a good moduli problem. Since $G \cap \mathbb{R}_{>0} = \{\text{Id}\}$ for any finite subgroup G of \mathbb{C}^* , the Kato–Nakayama space of the moduli problem is often representable even if the moduli problem itself is not. This is always the case for moduli problems involving, say, quotients of sheaves on $X[n]_0$ pulled back from X , since these quotients themselves have no automorphisms and the only isotropy comes from the subgroup of $(\mathbb{C}^*)^n$ preserving the quotient.

9. Log de Rham and Hodge structures

The main references of this section are [31, 30, 29]. This section owes much to a lecture by Phillip Griffiths [13].

Moduli spaces of polarized Hodge structures.

We assume the reader to be familiar with some basic concepts of Hodge theory. First of all, we briefly summarize the classical theory of the moduli spaces of polarized Hodge structures.

9.1. The moduli space $M_h = \Gamma \backslash D_h$. Let n be an integer, and let h be a sequence of non-negative integers $(h^{n,0}, h^{n-1,0}, \dots, h^{0,n})$ satisfying $h^{p,q} = h^{q,p}$, called the *Hodge numbers*. Let $H_{\mathbb{Z}}$ be a free abelian group of rank $\sum h^{p,n-p}$, with a non-degenerate bilinear form $Q : H_{\mathbb{Z}} \otimes H_{\mathbb{Z}} \rightarrow \mathbb{Z}$, which is symmetric (resp. anti-symmetric) if n is even (resp. odd). Let $G_{\mathbb{Z}}$ be the group functor $\text{Aut}(H_{\mathbb{Z}}, Q)$ on commutative rings, sending a ring R to the group of automorphisms on the free R -module $H_R := H_{\mathbb{Z}} \otimes R$ preserving the bilinear form Q . It is an affine group scheme of finite type over \mathbb{Z} (which is clear if we write down the matrix representing the bilinear form Q with respect to some basis of $H_{\mathbb{Z}}$). Let Γ be an arithmetic subgroup of $G_{\mathbb{Z}}(\mathbb{Z})$.

The set of Hodge structures of weight n on $H_{\mathbb{R}}$ with prescribed Hodge numbers h , such that Q induces a *polarization* on $H_{\mathbb{R}}$ (i.e. it induces a morphism $H_{\mathbb{R}} \otimes H_{\mathbb{R}} \rightarrow \mathbb{R}(-n)$ of Hodge structures, and the bilinear form $Q_C(u, v) := Q(u, Cv)$, where C is the Weil operator, is symmetric and positive definite), is parameterized by the homogeneous space $D_h = G_{\mathbb{R}}/K$, where K is the stabilizer

group of a fixed polarized Hodge structure F_0 on $H_{\mathbb{R}}$. See for instance ([9]) for these concepts.

This homogeneous space $D = D_h = G_{\mathbb{R}}/K$ has a complex structure defined as follows. It is clear that $Q : H_{\mathbb{R}} \otimes H_{\mathbb{R}} \rightarrow \mathbb{R}(-n)$ is a morphism of Hodge structures if and only if $Q(F^p, F^{n-p+1}) = 0$ for all p . Let $f^p = \sum_{r \geq p} h^{r, n-r}$, and let D^{\vee} , the *compact dual of D* , be the subspace of the product of the Grassmannians $\prod_p \text{Gr}(f^p, H_{\mathbb{R}})$ consisting of all flags F^{\bullet} :

$$\dots \subset F^{p+1} \subset F^p \subset \dots$$

such that $Q(F^p, F^{n-p+1}) = 0$. Then $D^{\vee} \simeq G_{\mathbb{C}}/P$, where P is a parabolic subgroup preserving a fixed flag. This gives D^{\vee} a complex structure. We see that $D \subset D^{\vee}$ is the locus of flags satisfying

- (i) $F^p \cap \overline{F}^{n-p+1} = 0$ (so that $F^p \oplus \overline{F}^{n-p+1} \cong H_{\mathbb{C}}$) for all p , and
- (ii) $Q(\overline{u}, Cu) > 0$ for all $u \neq 0$ in $H_{\mathbb{C}}$.

They are both open conditions, so $D \subset D^{\vee}$ is an open complex submanifold. The group Γ acts on D_h properly discontinuously, and the quotient $M_h = \Gamma \backslash D_h$ is the moduli space of Γ -*equivalence classes of Q -polarized Hodge structures on $H_{\mathbb{C}}$ with Hodge type h* . See ([31], 0.3.6, 0.3.7).

Variations of Hodge structures.

Definition 9.2. ([9], 3.11, 3.12) Let S be a complex manifold. A *variation of Hodge structures \mathcal{H} of weight n on S* is given by

- a local system $\mathcal{H}_{\mathbb{Z}}$ of free abelian groups of finite rank on S ;
- a finite decreasing filtration $F^{\bullet} \mathcal{H}_{\mathbb{C}}$ of the vector bundle $\mathcal{H}_{\mathbb{C}} := \mathcal{H}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_S$

by holomorphic sub-bundles,

such that the following conditions are satisfied:

- 1) (Griffiths transversality) the natural flat connection $\nabla = d \otimes \text{id}_{\mathcal{H}_{\mathbb{C}}} : \mathcal{H}_{\mathbb{C}} \rightarrow \Omega_S^1 \otimes \mathcal{H}_{\mathbb{C}}$ takes $F^p \mathcal{H}_{\mathbb{C}}$ into $\Omega_S^1 \otimes F^{p-1} \mathcal{H}_{\mathbb{C}}$, for every p ;
- 2) for each point $s \in S$, the fiber $F^{\bullet}(s)$ over s is a Hodge structure of weight n .

A *polarization* of the variation of Hodge structures \mathcal{H} is a locally constant bilinear form

$$\mathcal{Q} : \mathcal{H}_{\mathbb{Z}} \otimes \mathcal{H}_{\mathbb{Z}} \rightarrow \mathbb{Z}$$

such that on each fiber over $s \in S$, it induces a polarization of the fiber Hodge structure.

Suppose we have a polarized family of Hodge structures $(\mathcal{H}, \mathcal{Q} : \mathcal{H}_{\mathbb{Z}} \otimes \mathcal{H}_{\mathbb{Z}} \rightarrow \mathbb{Z})$ of weight n on S (not necessarily a variation of Hodge structures), and a global section of the sheaf

$$\Gamma \backslash \underline{\text{Isom}}((\mathcal{H}_{\mathbb{Z}}, \mathcal{Q}), (H_{\mathbb{Z}}, Q)),$$

where $H_{\mathbb{Z}}$ is regarded as a constant sheaf on S . If the monodromy group of this family of Hodge structures on S is contained in Γ , i.e. the image of the homomorphism

$$\pi_1(S) \rightarrow G_{\mathbb{Z}}(\mathbb{Z})$$

is contained in Γ , then there is a well-defined function

$$\varphi : S \rightarrow M_h$$

inducing this family of Hodge structures. This map is locally liftable to D_h .

If $f : X \rightarrow S$ is a projective smooth morphism between quasi-projective complex algebraic manifolds, with a relative hyperplane section $\eta \in H^0(S, R^2 f_* \mathbb{Z})$, then the family of the primitive parts $P^n(X_s, \mathbb{Z})$ of the cohomology groups $H^n(X_s, \mathbb{Z})$ modulo torsion form a polarized variation of Hodge structures of weight n on S , and it induces a map

$$\varphi : S \rightarrow M_h$$

if Γ contains the monodromy group. The map φ is called the *period map*. To be precise, the family of $H^n(X_s, \mathbb{C})$'s are the stalks of $R^n f_*(f^{-1} \mathcal{O}_S)$, and the Hodge filtration on $R^n f_*(f^{-1} \mathcal{O}_S)$ is given by the degenerate spectral sequence

$$E_1^{pq} = R^q f_* \Omega_{X/S}^p \implies R^{p+q} f_*(f^{-1} \mathcal{O}_S),$$

which is induced from the natural quasi-isomorphism (the relative holomorphic Poincaré lemma)

$$f^{-1} \mathcal{O}_S \longrightarrow \Omega_{X/S}^\bullet.$$

Since η is a global section, the primitive part form a family of sub-Hodge structures on S . Griffiths proved that the period map is holomorphic, and the family of Hodge structures is a variation of Hodge structures. See ([9], 3) for more detail.

For instance, one can take S to be the moduli space A_g (resp. M_g) of principally polarized abelian varieties of dimension g (resp. projective smooth curves of genus g), and take X to be the universal family of such objects over S , and $n = 1$. In this case, the local system $\mathcal{H}_{\mathbb{Z}} = R^1 f_* \mathbb{Z}$ is torsion-free and is equal to its primitive part, and the filtration $F^\bullet \mathcal{H}_{\mathcal{O}}$ is given by $F^0 \mathcal{H}_{\mathcal{O}} = \mathcal{H}_{\mathcal{O}}$, $F^1 \mathcal{H}_{\mathcal{O}} = R^0 f_* \Omega_{X/S}$ and $F^2 \mathcal{H}_{\mathcal{O}} = 0$. This variation of Hodge structures (and its tensor powers) are of great arithmetic interest.

Logarithmic Hodge structures.

Consider the following situation. Let $f : X \rightarrow S$ be a family of projective manifolds, where S is the complement of a normal crossing divisor D in a compact manifold \bar{S} , and suppose one can extend the family f to a family $\bar{f} : \bar{X} \rightarrow \bar{S}$ which is log smooth (here \bar{S} has the log structure induced by the divisor D (2.7)). Then one can ask if it is possible to enlarge the moduli space M_h to some \bar{M}_h so that the period map extends to $\bar{\varphi} : \bar{S} \rightarrow \bar{M}_h$.

To study the degenerations of Hodge structures, Kato and Usui introduced the notion of logarithmic Hodge structures [31].

Log de Rham complex. Let $(X, \alpha : M_X \rightarrow \mathcal{O}_X)$ be an fs log analytic space over \mathbb{C} , and let X^{\log} be the Kato-Nakayama space (Definition 8.1), with $\tau = \tau_X : X^{\log} \rightarrow X$ the natural proper map. Over the open set $X^* \subset X$ where the log structure is trivial, the map τ is a homeomorphism, and the section $j^{\log} : X^* \hookrightarrow X^{\log}$ is a homotopy equivalence. For $x \in X$, the fiber $\tau^{-1}(x)$ is a compact torus $(S^1)^m$, where m is the rank of $\overline{M}_{X,x}^{\text{gp}}$. For instance, let Δ be the open unit disk $\{|z| < 1\}$ equipped with the log structure induced by the center $\{z = 0\}$ (cf. 2.7). Then Δ^{\log} is homeomorphic to $[0, 1) \times S^1$. See sections 8.2 and 8.3 for more examples.

One can define a sheaf of rings $\mathcal{O}_{X^{\log}}$ on X^{\log} . Roughly speaking, this is the subsheaf of rings of $j_*^{\log} \mathcal{O}_{X^*}$ on X^{\log} generated over $\tau^{-1} \mathcal{O}_X$ by “ $\log(q)$ ”, for all $q \in M_X^{\text{gp}}$. See ([30], Section 3) for the precise definition.

For example, if $x \in X$ and $y \in \tau^{-1}(x)$, and the free abelian group $\overline{M}_{X,x}^{\text{gp}}$ has rank m and is generated by $f_1, \dots, f_m \in M_{X,x}^{\text{gp}}$, then the stalk $\mathcal{O}_{X^{\log},y}$ is isomorphic to the polynomial ring $\mathcal{O}_{X,x}[\log(f_1), \dots, \log(f_m)]$. This shows that in general, $(X^{\log}, \mathcal{O}_{X^{\log}})$ is not a locally ringed space.

For a morphism $f : X \rightarrow Y$ of fs log analytic spaces, one can define the *sheaf of relative log differentials* $\Omega_{X/Y}^1$ in the same way as Definition 3.5, namely it is the sheaf representing the functor of Y -log derivations (Definition 3.3), where we use the sheaf \mathcal{O}_X of holomorphic functions as the structure sheaf. The explicit description in Proposition 3.4 still applies. If (∂, D) is the universal Y -log derivation of X to $\Omega_{X/Y}^1$, the morphism $D : M_X \rightarrow \Omega_{X/Y}^1$ is also written as $d \log$, and it can be extended by linearity to M_X^{gp} . In the explicit description

$$\Omega_{X/Y}^1 = (\Omega_{X/Y} \oplus \mathcal{O}_X \otimes_{\mathbb{Z}} M_X^{\text{gp}}) / \mathcal{K},$$

$d \log(a)$ is the image of $0 \oplus (1 \otimes a)$, for a local section a of M_X^{gp} .

The sheaf $\Omega_{X/Y}^1$ is an analytic coherent \mathcal{O}_X -module. For an integer $r \geq 1$, let $\Omega_{X/Y}^r$ be the r -th exterior power of $\Omega_{X/Y}^1$. The derivation $\partial : \mathcal{O}_X \rightarrow \Omega_{X/Y}^1$ can be prolonged to a complex $\Omega_{X/Y}^\bullet$:

$$\mathcal{O}_X \xrightarrow{\partial} \Omega_{X/Y}^1 \xrightarrow{d} \Omega_{X/Y}^2 \xrightarrow{d} \dots \longrightarrow \Omega_{X/Y}^r \longrightarrow \dots$$

by imposing that $d(d \log(a)) = 0$ for $a \in M_X^{\text{gp}}$. This is a complex of $f^{-1} \mathcal{O}_Y$ -modules, called the *relative log de Rham complex on X with respect to f* .

For any sheaf F of \mathcal{O}_X -modules, define

$$\tau^* F := \tau^{-1} F \otimes_{\tau^{-1} \mathcal{O}_X} \mathcal{O}_{X^{\log}}$$

as a sheaf on X^{\log} . For an integer $r \geq 1$, define

$$\Omega_{X^{\log}/Y^{\log}}^r := \tau^* \Omega_{X/Y}^r.$$

The structure sheaf $\mathcal{O}_{X^{\log}}$ comes with a natural derivation $d : \mathcal{O}_{X^{\log}} \rightarrow \Omega_{X^{\log}/Y^{\log}}^1$ (see ([30], 3.5)), which can be prolonged to a complex $\Omega_{X^{\log}/Y^{\log}}^\bullet$ of $(f^{\log})^{-1}\mathcal{O}_{Y^{\log}}$ -modules

$$\mathcal{O}_{X^{\log}} \xrightarrow{d} \Omega_{X^{\log}/Y^{\log}}^1 \xrightarrow{d} \Omega_{X^{\log}/Y^{\log}}^2 \xrightarrow{d} \cdots \longrightarrow \Omega_{X^{\log}/Y^{\log}}^r \longrightarrow \cdots,$$

which can be called the *relative log de Rham complex on X^{\log} with respect to f* .

When Y is a point with trivial log structure, we denote $\Omega_{X/Y}^\bullet$ (resp. $\Omega_{X^{\log}/Y^{\log}}^\bullet$) by Ω_X^\bullet (resp. $\Omega_{X^{\log}}^\bullet$), and call it the *absolute log de Rham complex on X* (resp. X^{\log}).

Under mild conditions, F. Kato proved the relative log Poincaré lemma and the logarithmic analogue of the de Rham theorem. We state in the following a weaker version. See [25] for the more general version. We say an fs log analytic space is *log smooth* if it is so over a point with trivial log structure (pt, \mathbb{C}^*) .

Theorem 9.3. ([25], 3.4.2, 3.2.5) *Let $f : X \rightarrow Y$ be a log smooth morphism of fs log analytic spaces, with Y log smooth. Assume that the induced morphism $f^{-1}\overline{M}_Y \rightarrow \overline{M}_X$ is injective, that the stalk $\overline{M}_{X/Y,x}$ of the relative characteristic (4.1) is torsion-free for every $x \in X$, and that f is exact ([27], 4.6). Then there is a natural quasi-isomorphism*

$$(f^{\log})^{-1}\mathcal{O}_{Y^{\log}} \longrightarrow \Omega_{X^{\log}/Y^{\log}}^\bullet.$$

Corollary 9.4. ([25], 4.1.5) *Let $f : X \rightarrow Y$ be as in Theorem 9.3 above, and assume in addition that \underline{f} is proper. Then there is a natural quasi-isomorphism*

$$\tau_Y^* Rf_* \Omega_{X/Y}^\bullet \longrightarrow Rf_*^{\log} \underline{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_{Y^{\log}}.$$

These results apply in particular to semi-stable degenerations. Also, the absolute log Poincaré lemma was proved earlier by Kato and Nakayama ([30], Theorem 3.8).

Log variations of polarized Hodge structures. For $y \in X^{\log}$ and $x = \tau(y) \in X$, let $\text{sp}(y)$ be the set of all ring homomorphisms $s : \mathcal{O}_{X^{\log},y} \rightarrow \mathbb{C}$ that extend the evaluation map $\text{ev}_x : \mathcal{O}_{X,x} \rightarrow \mathbb{C}$. Since $\mathcal{O}_{X^{\log},y}$ is isomorphic to the polynomial ring over $\mathcal{O}_{X,x}$ generated by log of a basis for $\overline{M}_{X,x}$, if we fix an $s_0 \in \text{sp}(y)$, then we have a bijection:

$$s \mapsto (f \mapsto s(\log(f)) - s_0(\log(f))) : \text{sp}(y) \xrightarrow{\sim} \text{Hom}_{\text{group}}(\overline{M}_{X,x}^{\text{gp}}, \mathbb{C}),$$

where \mathbb{C} is viewed as an additive group.

Definition 9.5. Let X be an fs log analytic space. A *log variation of polarized Hodge structures of weight n* (abbreviated as *LVPHS*) on X is given by

- a local system of free abelian groups of finite rank $\mathcal{H}_{\mathbb{Z}}$ on X^{\log} ,
- a bilinear form $\mathcal{Q} : \mathcal{H}_{\mathbb{Z}} \otimes \mathcal{H}_{\mathbb{Z}} \rightarrow \underline{\mathbb{Z}}$,

• a finite decreasing filtration $F^\bullet \mathcal{H}_\theta$ of $\mathcal{H}_\theta := \mathcal{H}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_{X^{\log}}$ by $\mathcal{O}_{X^{\log}}$ -submodules,

such that the following conditions are satisfied:

1) there exist a locally free \mathcal{O}_X -module \mathcal{E} and a finite decreasing filtration $F^\bullet \mathcal{E}$ by \mathcal{O}_X -submodules, such that $\mathrm{Gr}_p(\mathcal{E})$ is locally free and

$$F^p \mathcal{H}_\theta \cong \tau^* F^p \mathcal{E}$$

for each p ;

2) for $y \in X^{\log}$ and $x = \tau(y) \in X$, let $s \in \mathrm{sp}(y)$ and let $f_1, \dots, f_r \in M_{X,x} - \mathcal{O}_{X,x}^*$ generate the monoid $\overline{M}_{X,x}$. If the real numbers $|\exp(s(\log(f_i)))|$ are sufficient small for all i , then $(\mathcal{H}_{\mathbb{Z},y}, \mathcal{Q}, F^\bullet(s))$ is a polarized Hodge structure of weight n ;

3) (Griffiths transversality) the connection $\nabla = d \otimes \mathrm{id}_{\mathcal{H}_{\mathbb{Z}}} : \mathcal{H}_\theta \rightarrow \Omega_{X^{\log}}^1 \otimes_{\mathcal{O}_{X^{\log}}} \mathcal{H}_\theta$ takes $F^p \mathcal{H}_\theta$ into $\Omega_{X^{\log}}^1 \otimes F^{p-1} \mathcal{H}_\theta$.

Here $F^\bullet(s)$, the *specialization of F^\bullet at s* , is the decreasing filtration of $\mathcal{H}_{\mathbb{C},y} := \mathbb{C} \otimes_{\mathbb{Z}} \mathcal{H}_{\mathbb{Z},y}$ defined by $F^p(s) = \mathbb{C} \otimes_{s, \mathcal{O}_{X^{\log},y}} (F^p \mathcal{H}_\theta)_y$. For a fixed point $y \in X^{\log}$, the family $(\mathcal{H}_{\mathbb{Z},y}, \mathcal{Q}, F^\bullet(s))_{s \in \mathrm{sp}(y)}$ is called a *polarized log Hodge structure of weight n* on the log point $(x, M_{X,x})$; this is the same as a log variation of polarized Hodge structures of weight n on the log point $(x, M_{X,x})$.

9.6. Variant. The definition we gave here follows ([31], 0.2.19), except that our polarization \mathcal{Q} is integral. This definition differs slightly from the one as in ([41], 5.3) and ([29], 2.3), and is weaker. The main difference is that, in *loc. cit.* the locally free \mathcal{O}_X -module \mathcal{E} with its filtration $F^\bullet \mathcal{E}$ is part of the data of the definition, and the flat connection is for \mathcal{E} on X (namely $\nabla : \mathcal{E} \rightarrow \Omega_X^1 \otimes \mathcal{E}$) and is required to satisfy the Griffiths transversality.

9.7. LVPHS from geometry. Log variations of polarized Hodge structures arise from geometry in the following way. Let $f : X \rightarrow Y$ be a projective vertical log smooth morphism between log smooth fs log analytic spaces, and we fix a line bundle on \underline{X} which is relatively very ample over \underline{Y} . Here “vertical” means $f^{-1}(Y^*) = X^*$. By a theorem of Kajiwara and Nakayama ([23], 0.3), for every integer n , the sheaf $R^n f_*^{\log} \underline{\mathbb{Z}}$ is a local system on Y^{\log} . We take $\mathcal{H}_{\mathbb{Z}}$ to be $R^n f_*^{\log} \underline{\mathbb{Z}}$ modulo torsion, take \mathcal{Q} to be the pairing induced by the fixed line bundle (which is obtained in the same way as ([29], 8.2) where we replace all local systems of \mathbb{Q} -vector spaces by the integral lattices in them), take \mathcal{E} to be $R^n f_*(\Omega_{X/Y}^\bullet)$, with filtration $F^p \mathcal{E} = R^n f_*(\Omega_{X/Y}^{\geq p}) \subset \mathcal{E}$, and take $F^p \mathcal{H}_\theta$ to be $\tau^* F^p \mathcal{E}$. Then by a theorem of Kato, Matsubara and Nakayama ([29], Theorem 8.1), this is a log variation of polarized Hodge structures on Y , even in the stronger sense ([29], 2.3). See ([29], Theorem 8.1) for a more general version with coefficients. The special case when Y is the unit disk $\{|z| < 1\}$ in the complex plane with log structure

induced by the divisor $\{z = 0\}$ and f is family of projective manifolds with semi-stable reduction over $\{z = 0\}$ was proved earlier by Matsubara ([41], Theorem C), except for the polarization part.

9.8. Relation to Deligne’s canonical extensions. Suppose that \underline{X} and \underline{Y} are smooth, that the log structure on Y is induced by a normal crossing divisor $D \subset Y$, and that $f : X \rightarrow Y$ is a morphism of semi-stable reduction. Let $f' : X^* \rightarrow Y^*$ be the restriction of f . Then the flat connection $(R^n f'_* \Omega_{X^*/Y^*}^\bullet, \nabla')$ on Y^* has unipotent local monodromy around each component of D , and the flat connection $(R^n f_* \Omega_{X/Y}^\bullet, \nabla)$ on Y is its canonical extension in the sense of Deligne [8].

Kato-Usui spaces.

We fix $n, h, H_{\mathbb{Z}}, Q, G_{\mathbb{Z}}, D$ and D^\vee as in (9.1). For a ring R , let $\mathfrak{g}_R = \text{Lie}(G_R)$. A subset $\sigma \subset \mathfrak{g}_{\mathbb{R}}$ is called a *nilpotent cone* if it is a cone

$$\sigma = \sum_{i=1}^n \mathbb{R}_{\geq 0} N_i$$

generated by mutually commutative nilpotent operators $N_i \in \mathfrak{g}_{\mathbb{R}} \subset \text{End}(H_{\mathbb{R}})$. It is called a *rational nilpotent cone* if it can be generated by nilpotent operators in $\mathfrak{g}_{\mathbb{Q}}$. Let Γ be a *neat subgroup* of $G_{\mathbb{Z}}(\mathbb{Z})$, i.e. for every element $\gamma \in \Gamma$, its eigenvalues on $H_{\mathbb{C}}$ generate a torsion-free subgroup of \mathbb{C}^* .

Nilpotent orbits.

Definition 9.9. Let $\sigma = \sum_i \mathbb{R}_{\geq 0} N_i$ be a nilpotent cone. A subset $Z \subset D^\vee$ is called a σ -*nilpotent orbit*, if there exists an $F_0 \in D^\vee$ such that

- $Z = \exp(\sum_i \mathbb{C} N_i) F_0$,
- $N F_0^p \subset F_0^{p-1}$ for all $p \in \mathbb{Z}$ and $N \in \sigma$,
- $\exp(\sum_i z_i N_i) F_0 \in D$ if $\text{Im}(z_i) \gg 0$ for all i .

We also call the pair (σ, Z) a *nilpotent orbit*.

Let Σ be a *fan* in $\mathfrak{g}_{\mathbb{Q}}$, i.e. Σ is a non-empty set of rational nilpotent cones in $\mathfrak{g}_{\mathbb{R}}$ such that

- if $\sigma \in \Sigma$, then all faces of σ are in Σ ,
- for $\sigma, \sigma' \in \Sigma$, the intersection $\sigma \cap \sigma'$ is a face of both σ and σ' ,
- for every $\sigma \in \Sigma$, we have $\sigma \cap (-\sigma) = 0$.

One can then define the set $D_{h, \Sigma}$ (or just D_Σ , if there is no confusion) of *nilpotent orbits in the directions in Σ* to be the set of nilpotent orbits (σ, Z) where $\sigma \in \Sigma$. There is a natural injection

$$F \mapsto (0, \{F\}) : D \hookrightarrow D_\Sigma.$$

The moduli space M_Σ . Let Σ be a fan in $\mathfrak{g}_\mathbb{Q}$ and let $\Gamma \subset G_\mathbb{Z}(\mathbb{Z})$ be a subgroup. Then we say that Γ is compatible with Σ if for every $\gamma \in \Gamma$ and $\sigma \in \Sigma$, we have $Ad(\gamma)(\sigma) \in \Sigma$. In this case, there is an action of Γ on D_Σ given by

$$(\sigma, Z) \xrightarrow{\gamma} (Ad(\gamma)(\sigma), \gamma Z).$$

We say that Γ is strongly compatible with Σ if every cone $\sigma \in \Sigma$ is generated by elements in $\log \Gamma$. Kato and Usui showed that when Γ is strongly compatible with Σ and the arithmetic subgroup Γ is neat, the quotient set $\Gamma \backslash D_\Sigma$ can be given the structure of a log locally ringed space over \mathbb{C} , in fact a *log manifold* (see ([31], 3.5.7)). Roughly speaking, a log manifold is a log locally ringed space over \mathbb{C} , which is locally isomorphic to the “zero locus” of some log differential forms on a log smooth analytic space.

Informally speaking, Kato and Usui proved the following. First, there is a one-to-one correspondence between D_Σ and the set of polarized log Hodge structures of the given type. Second, if $\overline{X} \rightarrow \overline{S}$ is a log smooth family extending the projective smooth family $X \rightarrow S$, where $S \subset \overline{S}$ is the complement of a normal crossing divisor, then the period map extends to $\overline{S} \rightarrow M_\Sigma$. We briefly explain the first part in the following.

We shall show how to get a nilpotent orbit from a polarized log Hodge structure on a log point ([31], 0.4.24). Let x be an fs log point with log structure M_x . Then \overline{M}_x is a sharp fs monoid and $\overline{M}_x^{\text{gp}}$ if a free abelian group of finite rank, say r . Fix $y \in x^{\text{log}}$. We have $x^{\text{log}} = \text{Hom}(\overline{M}_x^{\text{gp}}, S^1) \simeq (S^1)^r$ and hence $\pi_1(x^{\text{log}}) = \text{Hom}(\overline{M}_x^{\text{gp}}, \mathbb{Z}) \simeq \mathbb{Z}^r$. Let $\pi_1^+(x^{\text{log}}) \subset \pi_1(x^{\text{log}})$ be the subset consisting of those homomorphisms $a : \overline{M}_x^{\text{gp}} \rightarrow \mathbb{Z}$ that take \overline{M}_x into \mathbb{N} ; this subset is an fs monoid.

Let $(H_\mathbb{Z}, Q, F^\bullet H_\sigma)$ be a polarized log Hodge structure on x . Let $(h_i)_{i=1}^n$ be a family of generators for $\pi_1^+(x^{\text{log}})$ and fix an $s_0 \in \text{sp}(y)$. Let z_1, \dots, z_r be complex numbers, and let $s \in \text{sp}(y)$ be such that

$$s \left(\frac{\log(f)}{2\pi i} \right) - s_0 \left(\frac{\log(f)}{2\pi i} \right) = \sum_{i=1}^r z_i h_i(f), \quad \text{for } f \in \overline{M}_x^{\text{gp}}.$$

Let $N_i : H_{\mathbb{Q}, y} \rightarrow H_{\mathbb{Q}, y}$ be the logarithm of h_i . Then we have

$$F(s) = \exp \left(\sum_{i=1}^n z_i N_i \right) F(s_0),$$

which shows that $(F(s))_{s \in \text{sp}(y)}$ is an orbit of filtrations under $\exp(\sigma \otimes \mathbb{C})$ for $\sigma = \sum_i \mathbb{R}_{\geq 0} N_i$. Moreover, the condition 2) in Definition (9.5) implies that $F(s) \in D$ if $\text{Im}(z_i) \gg 0$ for all i , and the condition 3) in Definition (9.5) implies that $NF(s_0)^p \subset F(s_0)^{p-1}$ for all $p \in \mathbb{Z}$ and $N \in \sigma$. In other words, the family $(F(s))_s$ is a σ -nilpotent orbit.

10. The main component of moduli spaces

Moduli: compactness and main components

In Section 4, we gave an overview of F. Kato’s work [26] in which he uses log geometry to compactify the moduli space $\mathcal{M}_{g,n}$ of curves. Specifically, he shows that the moduli space of log smooth curves agrees with the Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$. The key philosophic idea in that section was that since moduli spaces of log smooth objects already includes degenerate objects, it is reasonable to expect that such a moduli space is a compactification of the moduli of objects with trivial log structure.

While the Deligne–Mumford space of stable curves $\overline{\mathcal{M}}_{g,n}$ turns out to be irreducible, it is an unfortunate fact of life that if \mathfrak{X} is a moduli space of higher dimensional objects, then moduli-theoretic “compactifications” $\tilde{\mathfrak{X}}$ of \mathfrak{X} tend to have many irreducible components. If \mathfrak{X} is irreducible, then it sits entirely within one of these many components of $\tilde{\mathfrak{X}}$ and so it is natural then to ask if this “main component” can itself be given a moduli interpretation.

In Section 4.14 we stated a second philosophic principle: log geometry controls degenerations; that is, moduli of log smooth objects does not incorporate “too many” degenerate objects. This provides a type of converse to the aforementioned philosophy that moduli of log smooth objects should be compact. As explained in Section 4.14, the log structure gives us a fighting chance to show that our moduli space is irreducible (although it is of course too naïve to expect that moduli of log smooth objects is always irreducible). Combining the two principles, one may hope that if \mathfrak{X} is an irreducible moduli space and $\tilde{\mathfrak{X}}$ a moduli-theoretic compactification, then by appropriately incorporating log structures into the objects parameterized by $\tilde{\mathfrak{X}}$, one will isolate the main component.

This technique of using log geometry to isolate the main component of a moduli space has been carried out by M. Olsson in several different settings. In [53], Olsson gives a moduli interpretation to the normalization of the main component of the toric Hilbert scheme. In [52], he isolates the normalization of the main component of V. Alexeev’s compactification of the moduli space of principally polarized abelian varieties given in [3]; he further constructs a moduli-theoretic irreducible compactification of the moduli space of abelian varieties with higher degree polarization. In [48], he gives an irreducible modular compactification of the moduli space of polarized $K3$ surfaces.

Example: the toric Hilbert scheme

Our goal in this section is to explain the technique of isolating the main component of a moduli space by following Olsson’s work [53]. We begin with the definition of the toric Hilbert scheme. Let k be a field and let P and Q be finitely-generated integral monoids with Q sharp and P^{gp} and Q^{gp} torsion-free.

Fix a surjective morphism $\pi : P \rightarrow Q$. This yields a closed immersion from $A_Q := \text{Spec } k[Q]$ to $A_P := \text{Spec } k[Q]$, which is T_Q -equivariant, where T_Q (resp. T_P) denotes the torus associated to Q^{gp} (resp. P^{gp}). Consider the functor \mathcal{H} whose S -valued points are diagrams

$$\begin{array}{ccc} Z & \xrightarrow{i} & A_{P,S} \\ & \searrow g & \downarrow \\ & & S \end{array}$$

where i is a T_Q -invariant closed immersion and for every $q \in Q^{gp}$, the q -eigenspace of $g_*\mathcal{O}_Z$ is a finitely-presented projective \mathcal{O}_S -module of rank 1 if $q \in Q$ and rank 0 otherwise. By [16, Thm 1.1], this functor is representable by a quasi-projective scheme, which we call the *toric Hilbert scheme*.

Given a closed subscheme Z of $A_{P,S}$ as above, we can move Z by the action of T_P on $A_{P,S}$. This yields an action of T_P on \mathcal{H} . Since Z is T_Q -invariant, this action factors through $T_K = T_P/T_Q$, where K denotes the kernel of π^{gp} . We therefore obtain a map

$$T_K \longrightarrow \mathcal{H}$$

by letting $u \in T_K$ act on the distinguished point of $\mathcal{H}(k)$ given by the closed immersion $A_Q \rightarrow A_P$. By [6, 3.6(2)], this map is an open immersion. Therefore the normalization \mathcal{S} of the scheme-theoretic closure of its image is a normal toric variety, and hence carries a natural fs log structure $\mathcal{M}_{\mathcal{S}}$ which makes it log smooth over $\text{Spec } k$ (endowed with the trivial log structure). The goal of [53] is to give a moduli-theoretic interpretation of $(\mathcal{S}, \mathcal{M}_{\mathcal{S}})$.

Consider the functor \mathcal{H}^{log} on the category of fs log schemes over k whose (S, \mathcal{M}_S) -valued points are given by diagrams

$$\begin{array}{ccc} (Z, \mathcal{M}_Z) & \xrightarrow{i} & (A_P, \mathcal{M}_{A_P}) \times (S, \mathcal{M}_S) \\ & \searrow g & \downarrow \\ & & (S, \mathcal{M}_S) \end{array}$$

where the underlying maps on schemes defines a point of $\mathcal{H}(S)$, where g is log smooth and integral, and where the map

$$P \longrightarrow \mathcal{M}_{(Z, \mathcal{M}_Z)/(S, \mathcal{M}_S)} := \text{coker}(g^* \mathcal{M}_S \rightarrow \mathcal{M}_Z)$$

induced by i factors through Q . The main theorem of [53] is then

Theorem 10.1 ([53, Thm 1.6]). *The functor \mathcal{H}^{log} is representable by $(\mathcal{S}, \mathcal{M}_{\mathcal{S}})$.*

We explain briefly how Olsson obtains a natural morphism

$$F : \mathcal{H}_{\mathcal{S}} \longrightarrow \mathcal{H}^{log}$$

which he then shows is an equivalence; here $\mathcal{H}_{\mathcal{S}}$ denotes the functor of points of $(\mathcal{S}, \mathcal{M}_{\mathcal{S}})$. Olsson obtains F by showing that the pullback to \mathcal{S} of the universal family over \mathcal{H} yields a point of $\mathcal{H}^{\text{log}}(\mathcal{S}, \mathcal{M}_{\mathcal{S}})$. Explicitly, if $i : \mathcal{Z} \rightarrow A_P \times \mathcal{S}$ is the pullback of the universal family, he constructs a log structure on \mathcal{Z} as follows. Since \mathcal{S} is a toric variety with torus T_K , it can be covered by open affines of the form $\text{Spec } k[L]$ with L a submonoid of K whose associated group is K . Olsson proves ([53, 2.4]) that over such an open affine, the closed immersion i is given by

$$\mathcal{Z} \times_{\mathcal{S}} \text{Spec } k[L] = \text{Spec } k[E_L] \longrightarrow \text{Spec } k[P \oplus L] = A_P \times \text{Spec } k[L],$$

where E_L is the image of $P \oplus L$ in P^{gp} under the map $(p, \ell) \mapsto p + \ell$.

Example 10.2. Let $\pi : \mathbb{N}^2 \rightarrow \mathbb{N}$ send e_1 to 2 and e_2 to 1. Then the kernel K of π^{gp} is all integer multiples of $2e_2 - e_1$. Let L be the submonoid of K consisting of the non-negative multiples of $e_1 - 2e_2$ and let M be the submonoid consisting of the non-positive multiples. Then E_L is generated by e_1 , e_2 , and $2e_2 - e_1$; hence,

$$\text{Spec } k[E_L] \simeq k[x, y, z]/(xy - z^2).$$

We see that E_M is freely generated by e_2 and $e_1 - 2e_2$, so $\text{Spec } k[E_M] \simeq \mathbb{A}_k^2$.

We see then that $\mathcal{Z} \times_{\mathcal{S}} \text{Spec } k[L]$ carries a natural log structure. These log structures glue to give a log structure on \mathcal{Z} (by [53, Lemma 2.8]) which makes i a closed immersion of log schemes. This therefore yields F above.

Example: moduli of K3 surfaces

In this section we discuss Olsson's work [48] on the moduli of K3 surfaces. Recall that a surface $\underline{X}/\underline{S}$ is called *K3* if for all geometric points s of \underline{S} , the canonical divisor $K_{\underline{X}_s}$ is trivial and $H^1(\underline{X}_s, \mathcal{O}_{\underline{X}_s}) = 0$. In his thesis, R. Friedman constructs partial compactifications of the coarse space of the moduli of polarized K3 surfaces over \mathbb{C} . The key notion in his construction is that of a combinatorial K3 surface, which we now recall (see [11, p.2]).

Definition 10.3. If k is an algebraically closed field, then a *combinatorial K3 surface* over $\underline{S} = \text{Spec } k$ is a k -scheme \underline{X} with normal crossings (see Section 5) which is d -semistable, has trivial dualizing sheaf, and satisfies one of the following: \underline{X} is a smooth K3 surface, \underline{X} is a chain of elliptic ruled surfaces with rational surfaces on either end, or \underline{X} is a union of rational surfaces where the double curves on each component form a cycle of rational curves with the dual graph of \underline{X} a triangulation of S^2 .

In light of the d -semistability condition and our discussion in Section 5, we should expect to be able to “explain” Friedman's moduli from the point of view of log geometry. With this in mind, we give the following definition of a log K3 surface (see [48, Def 5.1]).

Definition 10.4. A morphism $f : X \rightarrow S$ of log algebraic spaces is a *log K3 surface* if f is log smooth and integral, \underline{f} is proper, the cokernel of $f^* \mathcal{M}_S \rightarrow \mathcal{M}_X$ in the category of integral sheaves of monoids is a sheaf of groups, and for every geometric point s of \underline{S} , we have $\Omega_{X_s/s}^2 = \mathcal{O}_{X_s}$, $H^1(\mathcal{O}_{X_s}) = 0$, and \underline{X}_s is a normal crossing variety. The log K3 surface is called *stable* if it has no infinitesimal automorphisms.

As explained in [48, Rmk 5.3], if X/S is a log K3 surface, then for every geometric point s of \underline{S} , the fiber \underline{X}_s is a combinatorial K3 surface.

Now that we have a definition of log K3 surfaces, we discuss the logarithmic counterpart to the polarization. In order to ease the exposition, in what follows we will always assume that our log K3 surfaces satisfy the technical *special* condition defined in [48, Def 2.7]. Olsson introduces a notion of logarithmic Picard functor which generalizes the usual Picard functor in the case of trivial log structure (see Definition 4.5 and Corollary 5.6 of [48]):

Definition 10.5. If X/S is a log K3 surface, then the *log Picard functor* $\underline{Pic}(X/S)$ is the sheafification of the presheaf associating to any \underline{S} -scheme \underline{T} the isomorphism classes of $\mathcal{M}_{X_T}^{gp}$ -torsors on $\underline{X}_{T, \acute{e}t}$.

A polarization on a log K3 surface X/S is then defined to be ([48, Def 5.7]) a morphism $\lambda : \underline{S} \rightarrow \underline{Pic}(X/S)$ such that on each geometric fiber \underline{X}_s , there is a line bundle \mathcal{L} which lifts λ_s to $H^1(\underline{X}_s, \mathcal{O}_{\underline{X}_s}^*)$ and satisfies the following. There is some $N > 0$ such that \mathcal{L}^N is generated by global sections and the map defined by \mathcal{L}^N only contracts finitely many curves to points.

With these definitions in place, Olsson fixes a positive integer k and considers the stack $\mathbb{M}_{2k}/\mathbb{Q}$ whose fiber over \underline{T} is the groupoid of triples $(\mathcal{M}_T, X/(\underline{T}, \mathcal{M}_T), \lambda)$ where \mathcal{M}_T is a log structure on \underline{T} and $(X/(\underline{T}, \mathcal{M}_T), \lambda)$ is a stable polarized log K3 surface such that $\lambda_t^2 = 2k$ for every geometric point t of \underline{T} . Note that \mathbb{M}_{2k} carries a natural log structure given by base log structure \mathcal{M}_T in each fiber. One of the main results of [48] is then:

Theorem 10.6 ([48, Thm 6.2]). *The stack \mathbb{M}_{2k} is smooth, log smooth, Deligne-Mumford, and contains an open substack \mathbb{M}_{2k}^{sm} parameterizing polarized K3 surfaces in the classical sense. The compliment of \mathbb{M}_{2k}^{sm} in \mathbb{M}_{2k} is a smooth divisor and the induced log structure agrees with the natural one on \mathbb{M}_{2k} .*

11. Twisted curves and log twisted curves

Twisted curves are a central object in the theory of *twisted stable maps* [2, 5, 1]: in order to have a complete moduli space of stable maps $C \rightarrow X$ of type Γ , where X is a proper tame stack with projective coarse moduli space and $\Gamma = (g, n, \beta)$ are the relevant discrete data, one must allow the curve C itself to be a certain type of stack, called *twisted curve*.

The original treatments of twisted curves relied on ad-hoc methods. The more recent approach of [1] relies on a method introduced in [51], which uses a construction with logarithmic structures.

Twisted curves

For simplicity we will stick with the case of Deligne–Mumford stacks.

First consider the geometric objects: fix an algebraically closed field k .

Definition 11.1. A *twisted curve* over k is a tame, purely 1-dimensional Deligne–Mumford stack \mathcal{C}/k , with at most nodes as singularities, satisfying the following conditions:

- (1) Let $\pi : \mathcal{C} \rightarrow C$ be the morphism to the coarse moduli space. Then $\mathcal{C}^{sm} = \pi^{-1}C^{sm}$, and $\pi : \mathcal{C} \rightarrow C$ is an isomorphism over a dense open subset of C .
- (2) Consider a node $\bar{x} \rightarrow C$, where the strictly henselian local ring $\mathcal{O}_{C,\bar{x}}$ is the strict henselization of $k[x, y]/(xy)$. Then

$$\mathcal{C} \times_C \text{Spec } \mathcal{O}_{C,\bar{x}} \simeq \left[\text{Spec } \mathcal{O}_{C,\bar{x}}[z, w]/(zw, z^m - x, w^m - y) \Big/ \mu_m \right],$$

where $\zeta \in \mu_m$ acts by $(z, w) \mapsto (\zeta z, \zeta^{-1}w)$.

An action such as (2) above is called *balanced* - it is crucial to our discussion of log structures below. Note that \mathcal{C} may have a stack structure at isolated smooth points as well - such points will behave like $[\mathbb{A}^1/\mu_a]$, where μ_a acts by multiplication.

Over a general base S twisted curves are detected by their geometric fibers: a *twisted curve* $\mathcal{C} \rightarrow S$ is a flat, tame Deligne–Mumford stack locally of finite presentation, all of whose geometric fibers are twisted curves as in the definition above.

The *genus* of \mathcal{C} is simply the genus of C . One typically needs to consider n -pointed twisted curves, where the markings are described in families as follows:

Definition 11.2. An *n -pointed twisted curve* \mathcal{C}/S marked by disjoint closed substacks $\{\Sigma_i\}_{i=1}^n$ in \mathcal{C} is assumed to satisfy the following:

- (1) the Σ_i are contained in the smooth locus \mathcal{C}^{sm} ,
- (2) each Σ_i is a tame étale gerbe over S , and
- (3) $\mathcal{C}_{gen} := \mathcal{C}^{sm} \setminus \cup_i \Sigma_i \rightarrow C$ is an open embedding.

Remark 11.3. When $S = \text{Spec } k$ where $k = \bar{k}$, then $\Sigma_i = B\mu_{a_i}$, and moreover a_i is locally constant in families.

Remark 11.4. When $(\mathcal{C}/S, \{\Sigma_i\})$ is an n -pointed twisted curve, then the coarse moduli space of Σ_i is isomorphic to S . This means that the composite morphism $\Sigma_i \rightarrow \mathcal{C} \rightarrow C$ factors through a section $p_i : S \rightarrow C$. It follows that $(C, \{p_i\})$ is an n -pointed curve in the usual sense. This gives a functor

$$(\mathcal{C}/S, \{\Sigma_i\}) \mapsto (C, \{p_i\})$$

One can ask oneself: what does one need in order to recover a twisted n -pointed curve $(\mathcal{C}/S, \{\Sigma_i\})$ from a usual n -pointed curve $(C, \{p_i\})$? In other words, can we enrich the functor above to something like

$$(\mathcal{C}/S, \{\Sigma_i\}) \mapsto (C, \{p_i\}) + ?$$

which is nice and explicit and actually an equivalence of categories?

The stack structure at the marking definitely needs the data of the integers a_i , but in fact this is all that is necessary for the markings: near p_i , the curve \mathcal{C} is canonically isomorphic to the root stack $C(\sqrt[a_i]{p_i})$. If x is a local generator of the ideal of p_i , then Zariski locally we have

$$\mathcal{C} \simeq \left[\text{Spec } \mathcal{O}_C[z]/(z^{a_i} - x) \right] / \mu_{a_i}.$$

The story is a bit more interesting at a node. It has to be - a twisted curve \mathcal{C} with a node of index $m > 1$ has “ghost” automorphisms in μ_m which are not detectible on the coarse curve C : using the local coordinates given in Definition 11.1 (2), the μ_m action

$$(z, w) \mapsto (\zeta z, w)$$

on $\text{Spec } \mathcal{O}_{C, \bar{x}}[z, w]/(zw, z^m - x, w^m - y)$ commutes with the action defining the quotient stack. It therefore descends to a nontrivial action on \mathcal{C} which becomes trivial on the coarse moduli space C .

Log twisted curves

Let X be a Deligne–Mumford stack. Recall from Definition 2.16 that a fine log structure M on X is said to be *locally free* if for every geometric point $\bar{x} \rightarrow X$ we have that the characteristic sheaf $\overline{M}_{\bar{x}}$ is isomorphic to \mathbb{N}^r for some r .

In this situation we say that a morphism of sheaves of monoids $M \rightarrow M'$ is *simple* if for every geometric point $\bar{x} \rightarrow X$ we can identify the map as the diagonal map

$$\begin{array}{ccc} \overline{M}_{\bar{x}} & \longrightarrow & \overline{M}'_{\bar{x}} \\ \downarrow \simeq & & \downarrow \simeq \\ \mathbb{N}^r & \xrightarrow{(m_1, \dots, m_r)} & \mathbb{N}^r \end{array}$$

where all m_i are prime to the characteristic of the field.

Definition 11.5. An n -pointed log twisted curve over S is the data

$$(C/S, \{\sigma_i, a_i\}, \ell : M_S \rightarrow M'_S)$$

where

- $(C, \{\sigma_i\})/S$ is an n -pointed nodal curve.
- M_S is the canonical log structure coming from the family $(C, M_C) \rightarrow (S, M_S)$ (see section 4).

- $a_i : S \rightarrow \mathbb{Z}_{>0}$ are locally constant, with $a_i(s)$ invertible in the residue field $k(s)$.
- $\ell : M_S \rightarrow M'_S$ is a simple morphism.

And we have the following:

Theorem 11.6 ([51, Theorem 1.8]). *The fibered category of n -pointed twisted curves is naturally equivalent to the stack of n -pointed log twisted curves.*

The picture is as follows: we have already noted that we can replace a marking p_i by a stacky marking just using the data a_i . Now the j -th node which looks like $\text{Spec } \mathcal{O}_S[x, y]/(xy-t)$ needs to be replaced by $[\text{Spec } \mathcal{O}_S[z, w]/(zw - t^{1/m_j}) / \mu_{m_j}]$, and the data is encoding by deviding the j -th generator of \mathbb{N}^r by m_j .

Remark 11.7. We can decompose the stack according to a_i :

$$\coprod_{\underline{a}} \mathcal{M}_{g,n,\underline{a}}^{tw} = \mathcal{M}_{g,n}^{tw} = \text{stack of } n\text{-pointed twisted curves}$$

$$\downarrow$$

$$\mathcal{S}_{g,n} = \text{stack of } n\text{-pointed nodal curves}$$

and it can be deduced form the theorem that $\mathcal{M}_{g,n,\underline{a}}^{tw}$ is obtained from $\mathcal{S}_{g,n}$ using a root construction applied to the boundary divisor of $\mathcal{S}_{g,n}$, see [51, Remark 1.10].

In fact we have to apply all possible roots, accounting for all possible twisting of nodes, and glue together, so $\mathcal{M}_{g,n,\underline{a}}^{tw}$ is highly non-separated.

Below we sketch the main ideas in proving this. We stress that the assumption that our twisted curves are *balanced* is crucial - the case of unbalanced curves has not been treated.

From twisted curves to log twisted curves

Fix a twisted curve $f : \mathcal{C} \rightarrow S$.

We can follow F. Kato [26], giving log structures on nodal curves: consider all possible triples

$$(M_S, M_{\mathcal{C}}, f^b : f^{-1}M_S \rightarrow M_{\mathcal{C}})$$

such that

- (1) $(\mathcal{C}, M_{\mathcal{C}}) \rightarrow (S, M_S)$ is log smooth;
- (2) $M_{\mathcal{C}}, M_S$ are locally free; and
- (3) for all geometric points $\bar{x} \rightarrow \mathcal{C}$ mapping to nodes we have

$$\begin{array}{ccc} \overline{M}_{\mathcal{C},\bar{x}} & \xrightarrow{\sim} & \mathbb{N}^{r-1} \oplus \mathbb{N}^2 \\ \uparrow & & \uparrow \text{id} \oplus \Delta \\ \overline{M}_{S,f(\bar{x})} & \xrightarrow{\sim} & \mathbb{N}^{r-1} \oplus \mathbb{N} \end{array}$$

If $S = \text{Spec } k$ we get a natural map

$$\mathbb{N}^{\text{number of nodes}} \rightarrow \overline{M}_S.$$

We say that (M_C, M_S, f^b) is *special* if for every geometric point $\bar{s} \rightarrow S$ this map is an isomorphism.

A result [51, Theorem 3.6] analogous to F. Kato's [26, Theorem 2.1] says that there is a unique special triple (M_C, M'_S, f^b) associated to $f : \mathcal{C} \rightarrow S$. Analyzing the coarse moduli space we obtain a unique diagram

$$\begin{array}{ccc} (\mathcal{C}, M_C) & \longrightarrow & (C, M_C) \\ \downarrow & & \downarrow \\ (S, M'_S) & \xrightarrow{(id, \ell)} & (S, M_S) \end{array}$$

where $\ell : M_S \rightarrow M'_S$ is simple. In particular we obtain a log twisted curve

$$(C/S, \{\sigma_i, a_i\}_{i=1}^n, \ell : M_S \rightarrow M'_S)$$

From log twisted curves to twisted curves

Now we fix a log twisted curve $(C/S, \{\sigma_i, a_i\}_{i=1}^n, \ell : M_S \rightarrow M'_S)$. In particular we have the log smooth curve

$$(C, M_C) \rightarrow (S, M_S)$$

which is the coarse moduli space of a putative twisted curve. We want to describe \mathcal{C}/C as the stack parametrizing natural objects over $T \rightarrow C$. Here it is! If we denote the relevant maps as follows

$$\begin{array}{ccc} T & \xrightarrow{s} & C \\ & \searrow h & \downarrow \\ & & S, \end{array}$$

then \mathcal{C} is the groupoid of diagrams

$$\begin{array}{ccc} h^* M_S & \xrightarrow{\ell} & H^* M'_S \\ \downarrow & & \downarrow \\ s^* M_C & \xrightarrow{k} & M'_C, \end{array}$$

where

- (1) k is simple and for every geometric point $\bar{t} \rightarrow T$ the map $\overline{M}'_{S, \bar{t}} \rightarrow \overline{M}'_{C, \bar{t}}$ is either an isomorphism (at a general point), or of the form

$$\mathbb{N}^r \hookrightarrow \xrightarrow{(id, 0)} \mathbb{N}^r \oplus \mathbb{N}$$

(at a marked point), or

$$\mathbb{N}^{r-1} \oplus \mathbb{N} \xrightarrow{(id, \Delta)} \mathbb{N}^{r-1} \oplus \mathbb{N}^2$$

(at a node).

- (2) for all i and geometric point $\bar{t} \rightarrow T$ with $s(\bar{t}) \subset \sigma_i(S) \subset C$ the group

$$\text{Coker}(\overline{M}_{C, \bar{t}}^{gp} \rightarrow \overline{M}_{C, \bar{t}}^{gp})$$

is cyclic of order a_i .

12. Log stable maps

From curves to maps and expansions

Curves In section 4 we discussed how prestable curves can be encoded as log smooth curves, and how in particular the stack of Deligne–Mumford stable curves can be interpreted as a logarithmic stack, representing log smooth stable curves over the category of fine and saturated log schemes. Stability in this situation just means that $\Omega_{(C, \mathcal{M}_C)}^1$ is an ample line bundle. This is the same as saying that $\omega_C(D)$ is ample, where D is the divisor of markings.

Maps Kontsevich [35] introduced the moduli of *stable maps* of prestable curves into a projective target variety X . This is a proper Deligne–Mumford stack $\overline{\mathcal{M}}_\Gamma(X)$ having projective coarse moduli space, where $\Gamma = (g, n, \beta)$ is the relevant numerical data: genus, number of markings and homology class of the image curve. It parametrizes maps $f : C \rightarrow X$, where this time stability means that ω_C is f -ample.

Kontsevich’s moduli space has the property that it carries a *perfect relative obstruction theory* giving rise to a *virtual fundamental class* $[\overline{\mathcal{M}}_\Gamma(X)]^{vir}$, see [40, 4]. This is a key ingredient in defining Gromov–Witten invariants, with their applications in enumerative geometry and theoretical physics. The simplest GW invariants are

$$\langle \gamma_1 \cdots \gamma_n \rangle := \int_{[\overline{\mathcal{M}}_\Gamma(X)]^{vir}} e_1^* \gamma_1 \cdots e_n^* \gamma_n,$$

where $\gamma_i \in H^*(X, \mathbb{Q})$ and $e_i : \overline{\mathcal{M}}_\Gamma(X) \rightarrow X$ are the natural *evaluation maps* at the n markings.

So far no logarithmic structures are necessary.

Degenerations Among the methods of computing GW invariants, the *degeneration formula* is among the most powerful ones. It was introduced by A.-M. Li and Y. Ruan in symplectic geometry [37], see also Ionel–Parker [21, 22]. For our purposes, Jun Li’s treatment in algebraic geometry [38, 39] is most relevant.

We are interested in the invariants of the smooth projective variety X . Since these are deformation invariant, it is natural to consider a degeneration of X , with smooth total space, into a union $X_0 = Y_1 \sqcup_D Y_2$ of two smooth varieties Y_i meeting

transversally along a smooth divisor D . One wants to show that invariants of X coincide with suitably defined invariants of X_0 , and these in turn can be computed in terms of appropriately defined invariants of Y_i relative to D .

This is where logarithmic structures begin to show up, but there is still some way to go.

Perfect obstruction and Li's approach The difficulty with the degeneration is precisely the fact that the variety X_0 is singular, and therefore the natural obstruction theory on the moduli space of stable maps is not perfect in general. A similar situation occurs when considering the pair (Y_i, D) , but we will not get into this discussion.

The problem occurs when a component of the source curve C maps entirely into D .

Jun Li's approach uses *expanded degenerations*. There are similar ideas in [37], but the symplectic approach builds in deformations of Cauchy-Riemann equations and has, at least on the surface, a significantly different flavor.

The idea is, that just as in stable pointed curves, if a marking travels towards a node one sprouts a new component of the curve, Jun Li says that when a component of C travels into D we can let X_0 sprout a new component.

Expansions Here is how the new component looks like. Denote by $N_{D \subset Y_i}$ the normal bundle of D in Y_i . Since the total space is smooth, we have $N_{D \subset Y_1} \cong N_{D \subset Y_2}^\vee$. Let $\mathbb{P} = \mathbb{P}roj_D(1_D \oplus N_{D \subset Y_1})$. We have $\mathbb{P} \cong \mathbb{P}roj_D(N_{D \subset Y_2} \oplus 1_D)$ so we can denote by D^+ and D^- the smooth divisors in \mathbb{P} which correspond to the normal bundle $N_{D \subset Y_1}$ and $N_{D \subset Y_2}$ respectively. Note that the divisor D^+ and D^- are canonically isomorphic to D .

We now glue things up. Let \mathbb{P}_i for $i \in \mathbb{N}$ be copies of \mathbb{P} . We can glue Y_1 and \mathbb{P}_1 along D and D^- respectively, \mathbb{P}_i and \mathbb{P}_{i+1} along D^+ and D^- respectively, and \mathbb{P}_n and Y_2 along D^+ and D respectively. We denote the resulting gluing by

$$(12.1) \quad X_0[n] = Y_1 \bigsqcup_{D_1} \mathbb{P}_1 \bigsqcup_{D_2} \cdots \bigsqcup_{D_n} \mathbb{P}_n \bigsqcup_{D_{n+1}} Y_2,$$

where D_1, \dots, D_{n+1} are the disjoint singular loci of $X_0[n]$.

Such a beast is known as an *expanded degeneration*, or by the more folksy name *an n -accordion*.

This led Jun Li to define *degeneration stable maps with target X_0* as *nondegenerate maps $C \rightarrow X_0[n]$* : a map is nondegenerate if no component of C maps into any of the D_i .

Predeformability But here Jun Li meets another phenomenon, already present in the space of admissible covers of Harris–Mumford [17]: nondegenerate maps to $C \rightarrow X_0[n]$ have many redundant components which have nothing to do with maps to the generic fiber X . Near a singular point of $X_0[n]$ which looks like $\{xy = 0\}$,

a curve map locally deforms to a smoothing of X_0 if and only if the curve looks like $\{uv = 0\}$, and the map given by $x = u^m, y = v^m$. Such nice maps are called *predeformable*. But predeformable maps are clearly not open in the space of maps - they are actually closed among nondegenerate maps. This means that the restricted obstruction theory on them is “wrong” - definitely not perfect

Of course the virtual fundamental class of Gromov–Witten theory has no problem dealing with the total moduli space with its extra components, but these extra components do get in the way of decomposing invariants of X_0 in terms of (Y_i, D) . So we really do want to stick by predeformable maps.

Logarithmic methods: from Jun Li to Bumsig Kim

Predeformable deformations At this point Jun Li’s reasoning arrives at a point where a new obstruction theory on predeformable maps is needed. Having read this chapter, the reader will immediately recognize that

- (1) nodal curves are log-smooth,
- (2) n -accordions are d -semistable and admit a canonical log-smooth structure, and
- (3) predeformable maps can be viewed as log maps from the log-smooth curve to the d -semistable target.

Jun Li also recognized this fact, as did Shin Mochizuki before him [42] when he in his turn revisited the space of admissible covers of Harris–Mumford. What he lacked at the time was a formalism for logarithmic deformation theory of singular spaces, such as the moduli space itself: a perfect obstruction theory is a two-term complex mapping to the cotangent complex of the moduli space, but the moduli space is, as usual highly singular, even taking its logarithmic structure into account.

So Jun Li resorted to an ad-hoc construction of his perfect obstruction theory. This is the most difficult part of his work.

In section 7 we saw how log deformation theory works in the necessary generality. This is where Bumsig Kim’s paper [33] comes in: he provides a correct formalism for nondegenerate logarithmic stable maps into expanded degenerations, and shows that it carries a perfect obstruction theory. The degeneration formula in this formalism has been worked out by one of us (Q. Chen), and should appear as part of a larger project indicated below.

One aspect that deserves mention is Kim’s notion of *minimal log structures* on maps. Recall that the log stack $\overline{\mathcal{M}}_{g,n}$ can be constructed as a stack over the category of fine saturated log schemes, whose objects over $S = (\underline{S}, M)$ are log smooth stable pointed curves over S . But in order to exhibit the underlying stack, one needs to use the *canonical* log structure on \underline{S} , which is initial among all possible ones carrying the log smooth curve.

Kim describes his stack similarly - given a predeformable map $\underline{C} \rightarrow \underline{X}$ of underlying schemes over \underline{S} , it amounts to describing what he calls *minimal* log structure S on \underline{S} carrying a log map $C \rightarrow X$. Kim does this by explicitly describing the combinatorial structure of such log structures.

Unexpanded log maps: from Siebert into the future

A very different approach was proposed in a 2001 lecture by Bernd Siebert, but lay dormant for almost a decade.

The point is this. If one embraces logarithmic structures, and logarithmic maps from log smooth curves to some logarithmic scheme, then expansions are no longer necessary. Defined correctly, the space of log maps automatically has a perfect log-obstruction theory, which in view of sections 6 and 7 can be viewed as an obstruction theory relative to the stack LOG . This automatically results in a virtual fundamental class.

With this way of thinking, one can consider much more general logarithmic stable maps, gaining access to invariants of much more general degenerations of varieties. This has been a desired goal for a number of years.

So what is the correct definition? Consider a fine and saturated log scheme X . Following the work of F. Kato [26] as discussed in section 4, one comes up with a definition of a category $\overline{\mathcal{M}}_\Gamma(X)$ fibered over the category $LSch_{f_s}$ of fine and saturated log schemes: an object over a fine saturated log scheme S is a log smooth curve $C \rightarrow S$ and a log map $f : C \rightarrow X$. We further require it to be *stable*: the line bundle of logarithmic differentials $\Omega_{C/S}^1$ is required to be f -ample. This is tantamount to requiring the map of underlying schemes to be a stable map.

So the main claim is: $\overline{\mathcal{M}}_\Gamma(X)$ is represented by a logarithmic Deligne–Mumford stack with projective coarse moduli space. In fact this stack is proper and quasi-finite over the usual stack of stable maps $\overline{\mathcal{M}}_\Gamma(\underline{X})$ of the underlying scheme. As in the discussion of Kim’s work, the underlying stack can be viewed as a moduli of log maps with *minimal* log structure.

This is the subject of current work - of Gross and Siebert on the one hand and of two of us (mainly Chen, and to a lesser extent Abramovich) on the other, so it would not be appropriate to get into details until definite results appear.

Let us instead put this in a larger context. Consider logarithmic schemes $Z \rightarrow B$ and X , and assume we are given a morphism of underlying schemes $\underline{f} : \underline{Z} \rightarrow \underline{X}$. We can define a category $\text{Lift}_{\underline{f}}$ fibered over $LSch_{f_s}$ whose objects over a fine saturated log B -scheme $S \rightarrow B$ are lifts $f_S : Z_S \rightarrow X$ of the morphism of underlying schemes $\underline{f}_S : \underline{Z}_S \rightarrow \underline{X}$.

One can ask the following general questions:

- Question 12.2.* (1) Under what conditions is $\text{Lift}_{\underline{f}}$ a log stack locally of finite type over B ?
- (2) What natural numerical data cut out a substack of finite type?

(3) Under what conditions is the result proper?

We want to stress our belief that this question is natural, important and quite tractable. For instance, the case where $B = \text{Spec } \mathbb{C}$ with trivial structure and $Z = X = \text{Spec}(\mathbb{N} \rightarrow \mathbb{C})$ the result is a countable union of components. It is similar in nature to an inertia stack. The more general case where $B = X$, $Z = X \times \text{Spec}(\mathbb{N} \rightarrow \mathbb{C})$ and \underline{f} is the diagonal, is the relevant analogue of the inertia stack of X . It is important for Gromov–Witten theory - up to \mathbb{C}^* action the result is the natural target for evaluation maps associated to log smooth curves. Its components account for the contact orders of relative stable maps. Further examples of a similar nature govern gluing of nodes of log-smooth curves.

References

- [1] Dan Abramovich, Martin Olsson, and Angelo Vistoli. Twisted stable maps to tame artin stacks, 2008. <http://arxiv.org/abs/0801.3040>. 50, 51
- [2] Dan Abramovich and Angelo Vistoli. Compactifying the space of stable maps. *J. Amer. Math. Soc.*, 15(1):27–75 (electronic), 2002. 50
- [3] Valery Alexeev. Complete moduli in the presence of semiabelian group action. *Ann. of Math. (2)*, 155(3):611–708, 2002. 47
- [4] K. Behrend and B. Fantechi. The intrinsic normal cone. *Invent. Math.*, 128(1):45–88, 1997. 55
- [5] Weimin Chen and Yongbin Ruan. Orbifold Gromov-Witten theory. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 25–85. Amer. Math. Soc., Providence, RI, 2002. 50
- [6] Olga V. Chuvashova. The main component of the toric Hilbert scheme. *Tohoku Math. J. (2)*, 60(3):365–382, 2008. 48
- [7] P. Deligne and D. Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, (36):75–109, 1969. 18
- [8] Pierre Deligne. *Équations différentielles à points singuliers réguliers*. Lecture Notes in Mathematics, Vol. 163. Springer-Verlag, Berlin, 1970. 45
- [9] Pierre Deligne. Travaux de Griffiths. In *Séminaire Bourbaki. Vol. 1969/70: Exposés 364–381*, Lecture Notes in Mathematics, Vol. 180, pages 213–237: Exposé 376. Springer-Verlag, Berlin, 1971. 40, 41
- [10] Robert Friedman. Global smoothings of varieties with normal crossings. *Ann. of Math. (2)*, 118(1):75–114, 1983. 3, 19, 21, 22
- [11] Robert Friedman and Francesco Scattone. Type III degenerations of $K3$ surfaces. *Invent. Math.*, 83(1):1–39, 1985. 49

- [12] Ofer Gabber and Lorenzo Ramero. Foundations of p -adic hodge theory, 2009. <http://arxiv.org/abs/0409584>. 4
- [13] Phillip Griffiths. Logarithmic hodge structures: Report on the work of kato-usui. Notes from a talk given at MSRI on March 18, 2009. <http://math.brown.edu/~abrmovic/Griffiths-log.pdf>. 39
- [14] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I. *Inst. Hautes Études Sci. Publ. Math.*, (20):259, 1964. 8
- [15] A. Grothendieck and M. Raynaud. *Revêtements étales et groupe fondamental*. Springer-Verlag, Berlin, 1971. Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de M. Raynaud, Lecture Notes in Mathematics, Vol. 224. 14
- [16] Mark Haiman and Bernd Sturmfels. Multigraded Hilbert schemes. *J. Algebraic Geom.*, 13(4):725–769, 2004. 48
- [17] Joe Harris and David Mumford. On the Kodaira dimension of the moduli space of curves. *Invent. Math.*, 67(1):23–88, 1982. With an appendix by William Fulton. 56
- [18] W. J. Harvey. Boundary structure of the modular group. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 245–251. Princeton Univ. Press, Princeton, N.J., 1981. 34
- [19] Luc Illusie. *Complexe cotangent et déformations. I*. Lecture Notes in Mathematics, Vol. 239. Springer-Verlag, Berlin, 1971. 29
- [20] Luc Illusie. *Complexe cotangent et déformations. II*. Lecture Notes in Mathematics, Vol. 283. Springer-Verlag, Berlin, 1972. 29
- [21] Eleny-Nicoleta Ionel and Thomas H. Parker. Relative Gromov-Witten invariants. *Ann. of Math. (2)*, 157(1):45–96, 2003. 55
- [22] Eleny-Nicoleta Ionel and Thomas H. Parker. The symplectic sum formula for Gromov-Witten invariants. *Ann. of Math. (2)*, 159(3):935–1025, 2004. 55
- [23] Takeshi Kajiwara and Chikara Nakayama. Higher direct images of local systems in log Betti cohomology. *J. Math. Sci. Univ. Tokyo*, 15(2):291–323, 2008. 34, 44
- [24] Fumiharu Kato. Log smooth deformation theory. *Tohoku Math. J. (2)*, 48(3):317–354, 1996. 19, 21, 22
- [25] Fumiharu Kato. The relative log Poincaré lemma and relative log de Rham theory. *Duke Math. J.*, 93(1):179–206, 1998. 43
- [26] Fumiharu Kato. Log smooth deformation and moduli of log smooth curves. *Internat. J. Math.*, 11(2):215–232, 2000. 14, 15, 16, 17, 19, 47, 53, 54, 58
- [27] Kazuya Kato. Logarithmic structures of Fontaine-Illusie. In *Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988)*, pages 191–224. Johns Hopkins Univ. Press, Baltimore, MD, 1989. 1, 4, 8, 11, 13, 14, 15, 19,

- 24, 38, 43
- [28] Kazuya Kato. Toric singularities. *Amer. J. Math.*, 116(5):1073–1099, 1994. 12
 - [29] Kazuya Kato, Toshiharu Matsubara, and Chikara Nakayama. Log C^∞ -functions and degenerations of Hodge structures. In *Algebraic geometry 2000, Azumino (Hotaka)*, volume 36 of *Adv. Stud. Pure Math.*, pages 269–320. Math. Soc. Japan, Tokyo, 2002. 39, 44
 - [30] Kazuya Kato and Chikara Nakayama. Log Betti cohomology, log étale cohomology, and log de Rham cohomology of log schemes over C . *Kodai Math. J.*, 22(2):161–186, 1999. 34, 35, 36, 38, 39, 42, 43
 - [31] Kazuya Kato and Sampei Usui. *Classifying spaces of degenerating polarized Hodge structures*, volume 169 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009. 39, 40, 42, 44, 46
 - [32] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat. *Toroidal embeddings. I*. Lecture Notes in Mathematics, Vol. 339. Springer-Verlag, Berlin, 1973. 12
 - [33] Bumsig Kim. Logarithmic stable maps. <http://arxiv.org/abs/0807.3611>, 2008. 57
 - [34] János Kollár. Algebraic geometry. In Timothy Gowers, June Barrow-Green, and Imre Leader, editors, *The Princeton companion to mathematics*, pages 363–372. Princeton University Press, Princeton, NJ, 2008. 3
 - [35] Maxim Kontsevich. Enumeration of rational curves via torus actions. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 335–368. Birkhäuser Boston, Boston, MA, 1995. 55
 - [36] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000. 26
 - [37] An-Min Li and Yongbin Ruan. Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds. *Invent. Math.*, 145(1):151–218, 2001. 55, 56
 - [38] Jun Li. Stable morphisms to singular schemes and relative stable morphisms. *J. Differential Geom.*, 57(3):509–578, 2001. 55
 - [39] Jun Li. A degeneration formula of GW-invariants. *J. Differential Geom.*, 60(2):199–293, 2002. 55
 - [40] Jun Li and Gang Tian. Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. *J. Amer. Math. Soc.*, 11(1):119–174, 1998. 55
 - [41] Toshiharu Matsubara. On log Hodge structures of higher direct images. *Kodai Math. J.*, 21(2):81–101, 1998. 44, 45
 - [42] Shinichi Mochizuki. The geometry of the compactification of the Hurwitz scheme. *Publ. Res. Inst. Math. Sci.*, 31(3):355–441, 1995. 57

- [43] Chikara Nakayama and Arthur Ogus. Relative rounding, 2009. <http://math.berkeley.edu/~ogus/preprints/nakayamaogus.pdf>. 34, 38
- [44] Arthur Ogus. F -crystals, Griffiths transversality, and the Hodge decomposition. *Astérisque*, (221):ii+183, 1994. 9
- [45] Arthur Ogus. Lectures on logarithmic algebraic geometry. TeXed notes, 2001. http://math.berkeley.edu/~ogus/preprints/log_book/logbook.pdf. 4, 6, 7, 9, 10, 12
- [46] Martin C. Olsson. Logarithmic geometry and algebraic stacks. *Ann. Sci. École Norm. Sup. (4)*, 36(5):747–791, 2003. 24, 26, 27, 28, 29
- [47] Martin C. Olsson. Universal log structures on semi-stable varieties. *Tohoku Math. J. (2)*, 55(3):397–438, 2003. 19, 21, 22, 23
- [48] Martin C. Olsson. Semistable degenerations and period spaces for polarized $K3$ surfaces. *Duke Math. J.*, 125(1):121–203, 2004. 47, 49, 50
- [49] Martin C. Olsson. The logarithmic cotangent complex. *Math. Ann.*, 333(4):859–931, 2005. 14, 27, 28, 29, 30, 31, 32, 34
- [50] Martin C. Olsson. Deformation theory of representable morphisms of algebraic stacks. *Math. Z.*, 253(1):25–62, 2006. 29, 30, 32, 33
- [51] Martin C. Olsson. (Log) twisted curves. *Compos. Math.*, 143(2):476–494, 2007. 51, 53, 54
- [52] Martin C. Olsson. *Compactifying moduli spaces for abelian varieties*, volume 1958 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2008. 47
- [53] Martin C. Olsson. Logarithmic interpretation of the main component in toric Hilbert schemes. In *Curves and abelian varieties*, volume 465 of *Contemp. Math.*, pages 231–252. Amer. Math. Soc., Providence, RI, 2008. 47, 48, 49
- [54] Ulf Persson and Henry Pinkham. Some examples of nonsmoothable varieties with normal crossings. *Duke Math. J.*, 50(2):477–486, 1983. 21

BROWN UNIVERSITY, BOX 1917, PROVIDENCE, RI 02912, U.S.A.

E-mail address, Abarmovich: abrmovic@math.brown.edu

E-mail address, Chen: q.chen@math.brown.edu

E-mail address, Gillam: wdgillam@math.brown.edu

UNIVERSITY OF CALIFORNIA, BERKELEY

E-mail address, Huang: huang@math.berkeley.edu

E-mail address, Olsson: molsson@math.berkeley.edu

E-mail address, Satriano: satriano@math.berkeley.edu

E-mail address, Sun: shenghao@math.berkeley.edu