

# FAILURE OF THE HASSE PRINCIPLE ON GENERAL K3 SURFACES

BRENDAN HASSETT AND ANTHONY VÁRILLY-ALVARADO

ABSTRACT. We show that transcendental elements of the Brauer group of an algebraic surface can obstruct the Hasse principle. We construct a general K3 surface  $X$  of degree 2 over  $\mathbb{Q}$ , together with a two-torsion Brauer class  $\alpha$  that is unramified at every finite prime, but ramifies at real points of  $X$ . Motivated by Hodge theory, the pair  $(X, \alpha)$  is constructed from a double cover of  $\mathbb{P}^2 \times \mathbb{P}^2$  ramified over a hypersurface of bi-degree  $(2, 2)$ .

## 1. INTRODUCTION

Let  $X$  be a smooth projective geometrically integral variety over a number field  $k$ . If  $X$  has a  $k_v$ -point for every place  $v$  of  $k$  (equivalently, if its set  $X(\mathbf{A}_k)$  of adelic points is nonempty), yet it does not have a  $k$ -point, then we say that  $X$  does not satisfy the **Hasse principle**. Manin [Man71] showed that any subset  $\mathcal{S}$  of the Brauer group  $\mathrm{Br}(X) := H_{\text{ét}}^2(X, \mathbb{G}_m)$  may be used to construct an intermediate set

$$X(k) \subseteq X(\mathbf{A}_k)^{\mathcal{S}} \subseteq X(\mathbf{A}_k)$$

that often explains failures of the Hasse principle, in the sense that  $X(\mathbf{A}_k)^{\mathcal{S}}$  may be empty, even if  $X(\mathbf{A}_k)$  is not. In this case, we say there is a **Brauer-Manin obstruction** to the Hasse principle for  $X$ . See §4 for the definition of  $X(\mathbf{A}_k)^{\mathcal{S}}$ .

There is a filtration of the Brauer group  $\mathrm{Br}_0(X) \subseteq \mathrm{Br}_1(X) \subseteq \mathrm{Br}(X)$ , where

$$\begin{aligned} \mathrm{Br}_0(X) &:= \mathrm{im}(\mathrm{Br}(k) \rightarrow \mathrm{Br}(X)), \\ \mathrm{Br}_1(X) &:= \ker(\mathrm{Br}(X) \rightarrow \mathrm{Br}(\bar{X})), \end{aligned}$$

and  $\bar{X} = X \times_k \bar{k}$  for a fixed algebraic closure  $\bar{k}$  of  $k$ . Elements in  $\mathrm{Br}_0(X)$  are said to be **constant**; class field theory shows that if  $\mathcal{S} \subseteq \mathrm{Br}_0(X)$ , then  $X(\mathbf{A}_k)^{\mathcal{S}} = X(\mathbf{A}_k)$ , so these elements cannot obstruct the Hasse principle. Elements in  $\mathrm{Br}_1(X)$  are called **algebraic**; the remaining elements of the Brauer group are **transcendental**.

There is a large body of literature, spanning the last four decades, on algebraic Brauer classes and algebraic Brauer-Manin obstructions to the Hasse principle and the related notion of weak approximation (i.e., where sets  $\mathcal{S} \subseteq \mathrm{Br}_1(X)$  suffice to explain failures of these phenomena); see, for example [Man74, BSD75, CTC80, CTSSD87, CTKS87, SD93, SD99, KT04, Bri06, BBFL07, Cun07, Cor07, KT08, Log08, VA08, LvL09, EJ10a, EJ10b, Cor10, EJ11a]. The systematic study of these obstructions benefits in no small part from an isomorphism

$$(1) \quad \mathrm{Br}_1(X)/\mathrm{Br}_0(X) \xrightarrow{\sim} H^1(k, \mathrm{Pic}(\bar{X})),$$

coming from the Hochschild-Serre spectral sequence.

---

2000 *Mathematics Subject Classification*. Primary 11G35; Secondary 14G05, 14F22.

This research was supported by National Science Foundation Grants 0554491, 0901645, and 1103659.

Obstructions arising from transcendental elements, on the other hand, remain mysterious, because it is difficult to get a concrete handle on transcendental elements of the Brauer group; there is no known analogue of (1) for the group  $\mathrm{Br}(X)/\mathrm{Br}_1(X)$ .

If  $X$  is a curve, or a surface of negative Kodaira dimension, then  $\mathrm{Br}(\overline{X}) = 0$ , so the Brauer group is entirely algebraic. On the other hand, in 1996 Harari constructed a 3-fold with a transcendental Brauer-Manin obstruction to the Hasse principle [Har96]. This begs the question: what about algebraic surfaces? Can transcendental Brauer classes obstruct the Hasse principle on an algebraic surface? A natural place to study this question is the class of K3 surfaces; they are arguably some of the simplest surfaces of nonnegative Kodaira dimension in the Castelnuovo-Enriques-Manin classification. The group  $\mathrm{Br}(X)/\mathrm{Br}_1(X)$  is finite for a K3 surface [SZ08], but it can be nontrivial.

With arithmetic applications in mind, several authors over the last decade have constructed explicit transcendental elements on K3 surfaces [Wit04, SSD05, HS05, Ier10, Pre10, ISZ11, SZ12]. Wittenberg, Ieronymou and Preu have used these elements to exhibit obstructions to **weak approximation** (i.e., density of  $X(k)$  in  $\prod_v X(k_v)$  for the product of the  $v$ -adic topologies). In all cases the K3 surfaces considered have elliptic fibrations that play a vital role in the construction of transcendental classes.

Inspired by Hodge-theoretic work of van Geemen and Voisin [vG05, Voi86], we recently constructed a K3 surface with geometric Picard number 1 (and hence no elliptic fibrations), together with a transcendental Brauer class  $\alpha$  obstructing weak approximation; see [HVAV11] (joint with Varilly). The pair  $(X, \alpha)$  was obtained from a cubic fourfold containing a plane. At the time, we were unable to extend our work to obtain counterexamples to the Hasse principle, in part because we were unable to control the invariants of  $\alpha$  at real points of  $X$ —ironically, this is precisely the reason we obtain a counterexample to weak approximation! See Remarks 1.3 as well.

Taking advantage of some recent developments (see Remarks 1.3), our goal in this paper is to rectify the above situation and show, once and for all, that transcendental Brauer classes on algebraic surfaces can obstruct the Hasse principle.

**Theorem 1.1.** *Let  $X$  be a K3 surface of degree 2 over a number field  $k$ , with function field  $\mathbf{k}(X)$ , given as a sextic in the weighted projective space  $\mathbb{P}(1, 1, 1, 3) = \mathrm{Proj} k[x_0, x_1, x_2, w]$  of the form*

$$(2) \quad w^2 = -\frac{1}{2} \cdot \det \begin{pmatrix} 2A & B & C \\ B & 2D & E \\ C & E & 2F \end{pmatrix},$$

where  $A, \dots, F \in k[x_0, x_1, x_2]$  are homogeneous quadratic polynomials. Then the class  $\mathcal{A}$  of the quaternion algebra  $(B^2 - 4AD, A)$  in  $\mathrm{Br}(\mathbf{k}(X))$  extends to an element of  $\mathrm{Br}(X)$ .

When  $k = \mathbb{Q}$ , there exist particular polynomials  $A, \dots, F \in \mathbb{Z}[x_0, x_1, x_2]$  such that  $X$  has geometric Picard rank 1 and  $\mathcal{A}$  gives rise to a transcendental Brauer-Manin obstruction to the Hasse principle on  $X$ .

*Remark 1.2.* In [vG05, §9], Van Geemen showed that every Brauer class  $\alpha$  of order 2 on a polarized complex K3 surface  $(X, f)$  of degree 2 with  $\mathrm{Pic}(X) = \mathbb{Z}f$  gives rise to (and must arise from) one of three types of varieties:

- a smooth complete intersection of three quadrics in  $\mathbb{P}^5$  (itself a K3 surface), or

- a cubic fourfold containing a plane, or
- a double cover of  $\mathbb{P}^2 \times \mathbb{P}^2$  ramified along a hypersurface of bidegree  $(2, 2)$ .

More precisely, the class  $\alpha$  determines a sublattice  $T_\alpha \subseteq T_X$  of the transcendental lattice of  $X$  which is a polarized Hodge structure, a twist of which is Hodge isometric to a primitive sublattice of the middle cohomology of one the three types of varieties above. See §2 for more details.

The Azumaya algebra  $\mathcal{A}$  of Theorem 1.1 represents a class arising from a double cover of  $\mathbb{P}^2 \times \mathbb{P}^2$  ramified along a hypersurface of bidegree  $(2, 2)$ .

*Remarks 1.3.* We record a few remarks on the computational subtleties behind the second part of Theorem 1.1:

- (1) For computational purposes, we go in a direction “opposite” to van Geemen: starting from one of the three types of varieties described in Remark 1.2, defined over a number field  $k$ , we recover a K3 surface  $X$  over  $k$  of degree 2, together with a 2-torsion Azumaya algebra  $\mathcal{A}$ . Unfortunately, there is no guarantee that  $X$  has geometric Picard number  $\rho = 1$ ; in fact, it need not. We use a recent theorem of Elsenhans and Jahnel [EJ11c] to certify that our example has  $\rho = 1$ .
- (2) Curiously, one of the most delicate steps in the proof of Theorem 1.1 is determining the primes of bad reduction of  $X$ . We have to factor an integer with 318 decimal digits, whose smallest prime factor turns out to have 66 digits!
- (3) We use some of our work on varieties parametrizing maximal isotropic subspaces of families of quadrics admitting at worst isolated singularities to show that  $\mathcal{A}$  can ramify only at the real place, 2-adic places and primes of bad reduction for  $X$  [HVAV11, §3]. These are thus the only places where the local invariants of  $\mathcal{A}$  can be nontrivial.
- (4) We rely on recent work of Colliot-Thélène and Skorobogatov [CTS13] to control the local invariants for the algebra  $\mathcal{A}$  at odd primes of bad reduction.

*Remark 1.4.* The Azumaya algebra of Theorem 1.1 looks remarkably similar to the algebra we used in [HVAV11] to exhibit counter-examples to weak approximation. This is not a coincidence: compare Theorem 3.2 with [HVAV11, Theorem 5.1].

**Outline of the paper.** In §2 we explain the content of Remark 1.2 in detail, following van Geemen [vG05]. The section is not logically necessary for the paper, but we include it for completeness because it explains how to construct, in principle, Azumaya algebras representing every two-torsion Brauer class on a general K3 surface of degree 2.

In §3, we explain how to explicitly construct, from a double cover of  $\mathbb{P}^2 \times \mathbb{P}^2$  ramified along a hypersurface of bidegree  $(2, 2)$ , a pair  $(X, \alpha)$  where  $X$  is a K3 surface of degree 2 and  $\alpha \in \text{Br}(X)[2]$  is an Azumaya algebra. We work mostly over a discrete valuation ring (see Theorem 3.2). This flexibility later affords us control, when working over number fields, of local invariants at places where  $\alpha$  ramifies; see Lemma 4.4. In §4, we give a collection of sufficient conditions to control the evaluation maps of  $\alpha$  over number fields, specializing ultimately to the case  $k = \mathbb{Q}$ . Notably, Proposition 4.1 (due to Colliot-Thélène and Skorobogatov) together with Lemma 4.2 show that the evaluation maps of  $\alpha$  are constant at non-2-adic finite places of bad reduction of  $X$  whenever the singular locus consists of  $r < 8$  ordinary double points.

We use this preparatory work to give an example in §5 of a surface witnessing the second part of Theorem 1.1. In §6 we give details of how we found the example of §5, using a computer.

**Acknowledgments.** We thank Olivier Wittenberg for helpful comments and correspondence on transcendental Brauer classes on general K3 surfaces of degree 2 [Wit]. We also thank the referee for their careful reading of the manuscript and suggesting valuable improvements. Our computations were carried out using Macaulay 2 [GS], Magma [BCP97] and SAGE [S<sup>+</sup>09]

## 2. LATTICES AND HODGE THEORY

In this section, all varieties are defined over  $\mathbb{C}$ . Our goal here is to outline van Geemen's geometric constructions representing two-torsion Brauer classes on a K3 surface of degree 2 and Picard rank 1. Strictly speaking, this section is not logically necessary in the proof of Theorem 1.1, and we use only one of the three constructions described. We include it, however, so that readers not acquainted with these ideas get a clear sense of the geometric motivation behind Theorem 1.1.

Let  $X$  be a complex K3 surface. Regarding its middle cohomology as a lattice with respect to the intersection form, we can write [LP81, §1]

$$(3) \quad H^2(X, \mathbb{Z}) \simeq U^3 \oplus E_8(-1)^2 =: \Lambda$$

where

$$U = \langle e, f \rangle, \quad \text{with intersections} \quad \begin{array}{c|cc} & e & f \\ \hline e & 0 & 1 \\ f & 1 & 0 \end{array}$$

and  $E_8$  is the positive definite lattice arising from the corresponding root system, i.e., the unique positive definite even unimodular lattice of rank eight. Let  $e$  and  $f$  denote the generators of the first summand  $U$  in (3), and  $h \in H^2(X, \mathbb{Z})$  a primitive vector with  $h \cdot h = 2d > 0$ . The isomorphism (3) can be chosen so that

$$h \mapsto e + df.$$

Writing  $v = e - df$ , we have

$$h^\perp \simeq \mathbb{Z}v \oplus \Lambda', \quad \text{where } \Lambda' := U^2 \oplus E_8(-1)^2.$$

Let  $(X, h)$  be a polarized K3 surface of degree  $2d$ , i.e.,  $h \cdot h = 2d$ ; assume that  $\text{Pic}(X)$  is generated by  $h$ . Since  $H^3(X, \mathbb{Z}) = 0$ , the long exact sequence in cohomology associated to the exponential sequence gives rise to the short exact sequence

$$0 \rightarrow H^2(X, \mathbb{Z})/\langle h \rangle \rightarrow H^2(X, \mathcal{O}_X) \rightarrow \text{Br } X \rightarrow 0.$$

Applying the snake lemma to the diagram obtained by multiplication by 2 on this exact sequence, we see that two-torsion elements of the Brauer group of  $X$  may be interpreted as elements

$$\alpha \in H^2(X, \mathbb{Z}/2\mathbb{Z})/\langle h \rangle.$$

Under this identification, we can express

$$\alpha = n\bar{f} + \bar{\lambda}_\alpha, \quad n = 0, 1,$$

where  $\bar{f}$  is the image of  $f$  and  $\bar{\lambda}_\alpha$  is the image of some  $\lambda_\alpha \in \Lambda'$ . Using the non-degenerate cup product on  $H^2(X, \mathbb{Z})$ , let  $\alpha^\perp$  be the kernel of the map  $h^\perp \rightarrow \mathbb{Z}/2\mathbb{Z}$ ,  $x \mapsto x \cdot (nf + \lambda_\alpha)$ . We have

$$\alpha^\perp \subset h^\perp \subset H^2(X, \mathbb{Z}),$$

where the first subgroup has index two when  $\alpha \neq 0$ .

Assume from now on that  $\alpha \neq 0$ . If  $n = 1$  and  $\lambda_\alpha = 0$ , then  $\alpha^\perp = \mathbb{Z}(2v) \oplus \Lambda'$ , a lattice with discriminant group  $\mathbb{Z}/8d\mathbb{Z}$ , generated by  $v/2d$ . If  $\lambda_\alpha \neq 0$ , choose  $\mu \in \Lambda'$  satisfying

$$(4) \quad \mu \cdot \lambda_\alpha \equiv 1 \pmod{2}.$$

In this case

$$\alpha^\perp = \mathbb{Z}(v + \mu) + \{\lambda' \in \Lambda' : \lambda' \cdot \lambda_\alpha \equiv 0 \pmod{2}\},$$

a lattice with discriminant group generated by  $(-v + 2d\lambda_\alpha)/4d$ , which is therefore  $\mathbb{Z}/8d\mathbb{Z}$ .

If  $n = 0$  then  $\lambda_\alpha \neq 0$  and we can also choose  $\mu \in \Lambda'$  satisfying (4). We have

$$\alpha^\perp = \mathbb{Z}v \oplus \{\lambda' \in \Lambda' : \lambda' \cdot \lambda_\alpha \equiv 0 \pmod{2}\},$$

a lattice with discriminant group  $(\mathbb{Z}/2d\mathbb{Z}) \oplus (\mathbb{Z}/2\mathbb{Z})^2$ , where the last two summands are generated by  $\lambda_\alpha/2$  and  $\mu$ .

General results on quadratic forms (see, for example, [Nik79]) make it possible to classify even indefinite quadratic forms with prescribed rank and discriminant group  $H$ , provided the rank of the form is significantly larger than the number of generators of  $H$ . In particular, van Geemen [vG05, Proposition 9.2] classifies isomorphism classes of lattices  $\alpha^\perp$  arising from this construction:

- if  $n = 0$  there is a unique such lattice, up to isomorphism;
- if  $n = 1$  and  $d$  is even then there is a unique such lattice up to isomorphism;
- if  $n = 1$  and  $d$  is odd then there are two such lattices up to isomorphism, depending on the parity of  $\lambda_\alpha \cdot \lambda_\alpha/2$ .

He goes further when  $d = 1$ , offering geometric constructions of varieties having primitive Hodge structure isomorphic to  $\alpha^\perp$ . We elaborate on his description:

**Case  $n = 0$ :** Let  $W \subset \mathbb{P}^2 \times \mathbb{P}^2$  denote a smooth hypersurface of bidegree  $(2, 2)$  and  $Y \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$  the double cover branched along  $W$ . Let  $h_1$  and  $h_2$  denote the divisors on  $Y$  obtained by pulling back the hyperplane classes from the factors. We have intersections:

	$h_1^2$	$h_1 h_2$	$h_2^2$
$h_1^2$	0	0	2
$h_1 h_2$	0	2	0
$h_2^2$	2	0	0

The non-zero Hodge numbers of  $Y$  are:

$$h^{00} = h^{44} = 1, h^{11} = h^{33} = 2, h^{13} = h^{31} = 1, h^{22} = 22.$$

Consider the weight-two Hodge structure

$$\langle h_1^2, h_1 h_2, h_2^2 \rangle^\perp \subset H^4(Y)(1)$$

having underlying lattice  $M$ , with respect to the intersection form. The lattice  $M$  is even indefinite, and it has the same rank and discriminant group as  $\alpha^\perp$ . Thus

$$M \simeq \alpha^\perp.$$

**Case  $n = 1, \lambda_\alpha \cdot \lambda_\alpha \equiv 0 \pmod{4}$ :** In this case, there exists a *primitive* embedding

$$\alpha^\perp \hookrightarrow U^3 \oplus E_8(-1)^2,$$

unique up to automorphisms of the source and target. We can interpret the image as the primitive cohomology of a polarized K3 surface  $(S, f)$  with  $f \cdot f = 8$ .

**Case  $n = 1, \lambda_\alpha \cdot \lambda_\alpha \equiv 2 \pmod{4}$ :** Let  $Y$  be a cubic fourfold containing a plane  $P$ , with hyperplane class  $h$ . We have the intersections:

$$\begin{array}{c|cc} & h^2 & P \\ \hline h^2 & 3 & 1 \\ P & 1 & 3 \end{array}$$

The non-zero Hodge numbers of  $Y$  are

$$h^{00} = h^{44} = 1, h^{11} = h^{33} = 1, h^{13} = h^{31} = 1, h^{22} = 21.$$

The weight-two Hodge structure

$$\langle h^2, P \rangle^\perp \subset H^4(Y)(1)$$

has underlying lattice isomorphic to  $\alpha^\perp$ .

The last two geometric constructions yield explicit unramified Azumaya algebras over the degree two K3 surface. The connection between cubic fourfolds containing planes and quaternion algebras over the K3 surface can be found in [HVAV11]; the other construction goes back to Mukai [Muk84]: A degree eight K3 surface  $S$  is generally a complete intersection of three quadrics in  $\mathbb{P}^5$ , and the discriminant curve of the corresponding net is a smooth plane sextic. Let  $X$  be a degree-two K3 surface obtained as the double cover of  $\mathbb{P}^2$  branched along this sextic. The variety  $\mathcal{F}$  parametrizing maximal isotropic subspaces of the quadrics cutting out  $S$  admits a morphism (cf. [HVAV11, §3])  $\mathcal{F} \rightarrow X$ , which is smooth with geometric fibers isomorphic to  $\mathbb{P}^3$ .

In this paper, we focus on the first case, and use the resulting Azumaya algebra for arithmetic purposes.

### 3. UNRAMIFIED CONIC BUNDLES

Let  $k$  be an algebraically closed field of characteristic  $\neq 2$ , and let  $W$  be an irreducible **type  $(2, 2)$  divisor** on  $\mathbb{P}^2 \times \mathbb{P}^2$ , that is, hypersurface of bidegree  $(2, 2)$ . The two projections  $\pi_1: W \rightarrow \mathbb{P}^2$  and  $\pi_2: W \rightarrow \mathbb{P}^2$  define conic bundle structures on  $W$ . Let  $Y \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$  be the double cover branched along  $W$ . Composing this map with the projections onto the factors we obtain two quadric surface bundles  $q_i: Y \rightarrow \mathbb{P}^2$ . Note that the  $\pi_i$  and  $q_i$  need not be flat morphisms.

Let  $x_0, x_1, x_2$  and  $y_0, y_1, y_2$  denote homogeneous coordinates on the  $\mathbb{P}^2$ 's. The equation for  $W$  may be expressed as a quadratic form in the  $y_j$ 's with coefficients quadratic in the  $x_j$ 's (or vice versa). The determinant of the associated symmetric  $3 \times 3$  matrix of quadratic forms gives the locus over which  $\pi_1$  (or  $\pi_2$ ) ramifies. This determinant might be identically zero. Otherwise, we obtain plane sextic curves  $C_1$  and  $C_2$  over which  $\pi_1$  and  $\pi_2$  (as well as  $q_1$  and  $q_2$ ) are ramified. Let  $\phi_i: X_i \rightarrow \mathbb{P}^2$  be a double cover of  $\mathbb{P}^2$  branched over  $C_i$ ; if  $C_i$  is a smooth curve then  $X_i$  is a K3 surface of degree 2.

If  $W$  has at worst isolated singularities then the geometric generic fiber of  $\pi_i$  (and hence of  $q_i$ ) is smooth. Indeed, if the geometric generic fiber of  $\pi_i$  were singular then there would exist a generically étale  $V \rightarrow \mathbb{P}^2$  such that  $V \times_{\mathbb{P}^2} W$  is singular over the generic point of  $V$ . Here we are using the fact that the characteristic  $\neq 2$ . Thus it follows that  $C_1$  and  $C_2$  are curves.

**Lemma 3.1.** *Retain the notation above and assume that  $C_1$  and  $C_2$  are curves. If  $Y$  (equivalently,  $W$ ) is not smooth then neither  $C_1$  nor  $C_2$  is smooth. On the other hand, if  $W$  is smooth then each  $\pi_i$  is flat and  $C_i$  is singular if  $\pi_i$  has a geometric fiber of rank 1,  $i = 1, 2$ .*

*Proof.* An easy application of the Jacobian criterion shows that  $Y$  is smooth if and only if  $W$  is smooth. We use the latter scheme to prove the remaining claims of the lemma.

Let  $w \in W$  be a singular point,  $c_i \in C_i$  its images under projection,  $(u_0, u_1)$  local coordinates of the first  $\mathbb{P}^2$  centered at  $c_1$ , and  $(v_0, v_1)$  local coordinates of the second  $\mathbb{P}^2$  centered at  $c_2$ . The defining equation of  $W$  takes the form

$$a(u_0, u_1)v_0^2 + b(u_0, u_1)v_0v_1 + 2d(u_0, u_1)v_1^2 + c(u_0, u_1)v_0 + e(u_0, u_1)v_1 = 0,$$

where the coefficients are quadratic in  $u_0$  and  $u_1$ , and  $c(0, 0) = e(0, 0) = 0$ . Note that  $W$  would be singular if the coefficients  $a(u_0, u_1), \dots, e(u_0, u_1)$  were all zero. The defining equation for  $C_1$  is therefore

$$\det \begin{pmatrix} 2a & b & c \\ b & 2d & e \\ c & e & 0 \end{pmatrix} = 0.$$

Expanding this out, we get

$$bce - ae^2 - c^2d = 0,$$

where each term vanishes to order  $\geq 2$  at  $c_1 = (0, 0)$ .

The last statement of the lemma is a consequence of [Bea77, Prop. 1.2].  $\square$

**Theorem 3.2.** *Let  $\mathcal{O}$  denote a discrete valuation ring with residue field  $\mathbb{F}$  of characteristic  $\neq 2$ . Let  $\mathcal{W}$  be a type  $(2, 2)$  divisor in  $\mathbb{P}^2 \times \mathbb{P}^2$  flat over  $\text{Spec } \mathcal{O}$ , and  $\mathcal{Y} \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$  a double cover simply branched along  $\mathcal{W}$ . For  $i = 1, 2$ , let  $q_i: \mathcal{Y} \rightarrow \mathbb{P}^2$  denote the quadric surface bundle obtained by projecting onto the  $i$ -th factor, and let  $C_i \subset \mathbb{P}^2$  be its discriminant divisor. Assume that for some  $j \in \{1, 2\}$ ,  $C_j$  is flat over  $\mathcal{O}$ , and that  $(C_j)_{\mathbb{F}}$  is smooth.*

*Let  $r_j: \mathcal{F}_j \rightarrow \mathbb{P}^2$  be the relative variety of lines of  $q_j$ . Then the Stein factorization*

$$r_i: \mathcal{F}_j \rightarrow \mathcal{X}_j \xrightarrow{\phi_j} \mathbb{P}^2$$

*consists of a smooth  $\mathbb{P}^1$ -bundle followed by a degree-two cover of  $\mathbb{P}^2$ , which is a K3 surface.*

*Proof.* By Lemma 3.1, smoothness of  $(C_j)_{\mathbb{F}}$  implies smoothness of  $\mathcal{W}_{\mathbb{F}}$ , and hence of  $\mathcal{Y}_{\mathbb{F}}$ . The same lemma shows that the fibers of  $(q_j)_{\mathbb{F}}$  have at worst isolated singularities. On the other hand, the morphism  $q_j: \mathcal{Y} \rightarrow \mathbb{P}^2$  is flat, and thus  $\mathcal{Y}$  is a regular scheme. Geometric fibers of  $q_j$  over the generic point of  $\text{Spec } \mathcal{O}$  with non-isolated singularities specialize to geometric fibers over the closed point with non-isolated singularities. Hence the fibers of  $q_j$  have isolated singularities. The theorem now follows directly from [HVAV11, Proposition 3.3]: The Stein factorization of the variety of maximal isotropic subspaces of a family of even-dimensional quadric hypersurfaces with (at worst) isolated singularities is isomorphic to the discriminant double cover of the base.

Since the morphism  $\mathcal{C}_j \rightarrow \mathbb{P}^2$  is flat, smoothness of  $(\mathcal{C}_j)_{\mathbb{F}}$  implies that  $\mathcal{C}_j$  is regular. Hence  $\mathcal{X}_j$  is a K3 surface over  $\text{Spec } \mathcal{O}$ .  $\square$

The smooth  $\mathbb{P}^1$ -bundle  $r_j: \mathcal{F}_j \rightarrow \mathcal{X}_j$  may be interpreted as a two-torsion element of  $\text{Br}(\mathcal{X}_j)$ . Without loss of generality, assume in the hypotheses of Theorem 3.2 that  $(\mathcal{C}_1)_{\mathbb{F}}$  is smooth; we omit the subscript  $j = 1$  from here on in. We give an explicit quaternion algebra over  $\mathbf{k}(\mathcal{X})$  representing the Brauer class of  $\mathcal{F} \rightarrow \mathcal{X}$ . Express

$$\mathbb{P}^2 \times \mathbb{P}^2 = \text{Proj } \mathcal{O}[x_0, x_1, x_2] \times_{\mathcal{O}} \text{Proj } \mathcal{O}[y_0, y_1, y_2]$$

so the equation for  $\mathcal{W}$  takes the form

$$(5) \quad 0 = A(x_0, x_1, x_2)y_0^2 + B(x_0, x_1, x_2)y_0y_1 + C(x_0, x_1, x_2)y_0y_2 \\ + D(x_0, x_1, x_2)y_1^2 + E(x_0, x_1, x_2)y_1y_2 + F(x_0, x_1, x_2)y_2^2,$$

for some homogeneous quadratic polynomials  $A, \dots, F \in \mathcal{O}[x_0, x_1, x_2]$ . The coefficients are unique modulo multiplication by a common unit in  $\mathcal{O}$ .

Consider the bigraded ring  $\mathcal{O}[x_0, x_1, x_2, y_0, y_1, y_2, v]$  where

$$\deg(x_i) = (1, 0), \quad \deg(y_i) = (0, 1), \quad \deg(v) = (1, 1),$$

and let

$$R := \bigoplus_{n \in \mathbb{Z}} \mathcal{O}[x_0, x_1, x_2, y_0, y_1, y_2, v]_{(n, n)}$$

denote the graded subring generated by elements of bidegree  $(n, n)$  for some  $n$ . Then an equation for  $\mathcal{Y} \subset \text{Proj } R$  is

$$(6) \quad v^2 = A(x_0, x_1, x_2)y_0^2 + B(x_0, x_1, x_2)y_0y_1 + C(x_0, x_1, x_2)y_0y_2 \\ + D(x_0, x_1, x_2)y_1^2 + E(x_0, x_1, x_2)y_1y_2 + F(x_0, x_1, x_2)y_2^2.$$

The quadric surface bundle  $q: \mathcal{Y} \rightarrow \mathbb{P}^2$  is ramified over the curve

$$(7) \quad \mathcal{C}: \quad \det \begin{pmatrix} 2A & B & C & 0 \\ B & 2D & E & 0 \\ C & E & 2F & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} = 0.$$

Thus, after rescaling, the K3 surface  $\mathcal{X}$  is described by the hypersurface

$$(8) \quad w^2 = -\frac{1}{2} \cdot \det(M)$$

in  $\mathbb{P}(1, 1, 1, 3)$ , where  $M \in \text{Mat}_3(\mathcal{O}[x_0, x_1, x_2])$  is the leading  $3 \times 3$  principal minor of the matrix in (7). The factor  $-\frac{1}{2}$  is dictated by the interpretation of the structure sheaf of  $\mathcal{X}$  as the discriminant algebra of our quadratic form in four variables (cf. [HVAV11, §3.1]).

The discussion in [HVAV11, §3.3] shows that the generic fiber of the map  $\mathcal{F} \rightarrow \mathcal{X}$  is the Severi-Brauer conic in  $\text{Proj } \mathbf{k}(\mathcal{X})[Y_0, Y_1, Y_2]$  given by

$$(9) \quad AY_0^2 + BY_0Y_1 + CY_0Y_2 + DY_1^2 + EY_1Y_2 + FY_2^2 = 0.$$

Essentially, given a smooth quadric surface whose discriminant double cover is split, each component of the variety of lines on the surface is isomorphic to a smooth hyperplane section of the surface. Let

$$M_A := 4DF - E^2, \quad M_D := 4AF - C^2, \quad \text{and} \quad M_F := 4AD - B^2.$$



Completing squares in (9), and renormalizing, we obtain

$$Y_0^2 = -\frac{M_F}{4A^2}Y_1^2 - \frac{\det(M)}{2A \cdot M_F}Y_2^2.$$

Hence, by [GS06, Corollary 5.4.8], the conic (9) corresponds to the Hilbert symbol

$$(10) \quad \left( -\frac{M_F}{4A^2}, -\frac{\det(M)}{2A \cdot M_F} \right).$$

Write  $\mathcal{A}$  for the class of this symbol in  $\text{Br}(\mathbf{k}(\mathcal{X}))$ ;  $\mathcal{A}$  is unaffected by multiplication by squares in either entry of a representative symbol. Since  $-\frac{1}{2}\det(M)$  is a square in  $\mathbf{k}(\mathcal{X})^\times$ , we see that

$$(-M_F, A \cdot M_F) = (-M_F, A)$$

is another representative of  $\mathcal{A}$  (the equality uses the multiplicativity of the Hilbert symbol and the relation  $(-M_F, M_F) = 1$  [Ser73, III, Proposition 2]). Here we have the usual abuse of notation: the entries are not rational functions, though they are homogeneous polynomials of even degree.

Depending on how we complete squares and renormalize (9), we may obtain several representatives of  $\mathcal{A}$ :

$$(11) \quad \begin{array}{lll} (-M_F, A), & (-M_D, A), & (-M_F, D), \\ (-M_A, D), & (-M_D, F), & (-M_A, F). \end{array}$$

**Proposition 3.3.** *Let  $X$  be a K3 surface of degree 2 over a number field  $k$ , given as a sextic in the weighted projective space  $\mathbb{P}(1, 1, 1, 3) = \text{Proj } k[x_0, x_1, x_2, w]$  of the form*

$$w^2 = -\frac{1}{2} \cdot \begin{pmatrix} 2A & B & C \\ B & 2D & E \\ C & E & 2F \end{pmatrix},$$

where  $A, \dots, F \in k[x_0, x_1, x_2]$  are homogeneous quadratic polynomials. Then the class  $\mathcal{A}$  of the quaternion algebra  $(B^2 - 4AD, A)$  in  $\text{Br}(\mathbf{k}(X))$  extends to an element of  $\text{Br}(X)$ .

*Proof.* Let  $\mathcal{O}$  be the valuation ring at some finite place of  $k$  where  $X$  has good reduction. The proposition follows directly from Theorem 3.2 and the subsequent discussion, keeping track of what is happening over the generic point of  $\text{Spec } \mathcal{O}$ . Indeed, define  $\mathcal{W}$  and  $\mathcal{Y}$ , respectively, by (5) and (6). The resulting curve  $(\mathcal{C}_1)_k$  is the branch curve of the double cover  $X \rightarrow \mathbb{P}^2$ , which is smooth because  $X$  is a K3 surface, by hypothesis.  $\square$

*Remark 3.4.* The assortment of quaternion algebras (11) representing the class  $\mathcal{A}$  of Proposition 3.3 is useful for the computation of the invariant map on the image of the evaluation map  $\text{ev}_{\mathcal{A}}: X(\mathbf{A}_k) \rightarrow \bigoplus_v \text{Br}(k_v), (P_v) \mapsto (\mathcal{A}(P_v))$ . The industrious reader can check that at every local point of  $X$ , either the first, fourth or fifth representative of  $\mathcal{A}$  in our list is well-defined; we shall not use this observation directly.

## 4. LOCAL INVARIANTS

Let  $X$  be a smooth projective geometrically integral variety over a number field  $k$ . For  $\mathcal{S} \subseteq \text{Br}(X)$ , let

$$X(\mathbf{A}_k)^{\mathcal{S}} := \left\{ (P_v) \in X(\mathbf{A}_k) : \sum_v \text{inv}_v \mathcal{A}(P_v) = 0 \text{ for all } \mathcal{A} \in \mathcal{S} \right\}.$$

The inclusion  $X(k) \subseteq X(\mathbf{A}_k)^{\mathcal{S}}$  follows from class field theory. See [Sko01, §5.2] for details. The local invariants  $\text{inv}_v \mathcal{A}(P_v)$  can be nonzero only at a finite number of places: the archimedean places of  $k$ , the places of bad reduction of  $X$ , and places where the class  $\mathcal{A}$  is ramified.

We begin this section by explaining how recent work of Colliot-Thélène and Skorobogatov [CTS13] shows that local invariants are *constant* at certain finite places  $v$  of bad reduction for  $X$  where the singular locus satisfies a technical hypothesis. Specializing to the case where  $X$  is a K3 surface over a number field  $k$  as in Proposition 3.3, this technical hypothesis is satisfied provided the singular locus at  $v$  consists of  $r < 8$  ordinary double points (Lemma 4.2).

We then show that the class  $\mathcal{A}$  of Proposition 3.3 can ramify only over infinite places, 2-adic places, and places of bad reduction for  $X$ . Finally, in the special case  $k = \mathbb{Q}$ , we give sufficient conditions for local invariants of  $\mathcal{A}$  to be trivial at 2-adic points and nontrivial at real points.

**4.1. Places of bad reduction with mild singularities.** In this section we use the following notation:  $k$  is a finite extension of  $\mathbb{Q}_p$  with a fixed algebraic closure  $\bar{k}$ ,  $\mathcal{O}$  denotes the ring of integers of  $k$ , and  $\mathbb{F}$  denotes its residue field. We let  $X$  be a smooth, proper, geometrically integral variety over  $k$  and write  $\pi: \mathcal{X} \rightarrow \text{Spec } \mathcal{O}$  for a flat proper morphism with  $X = \mathcal{X} \times_{\mathcal{O}} k$ .

The following proposition is a straightforward refinement of [CTS13, Proposition 2.4], using ideas in the remark on the case of bad reduction in [CTS13, §2]. We include the details here for the reader's convenience.

**Proposition 4.1.** *Let  $\ell \neq p$  be a prime. Assume that  $\mathcal{X}$  is regular with geometrically integral fibers over  $\text{Spec } \mathcal{O}$ , and that the smooth locus  $\mathcal{X}_{\mathbb{F}}^{\text{sm}}$  of the closed fiber is geometrically irreducible and has no connected unramified cyclic geometric coverings of degree  $\ell$ . If  $X(k) \neq \emptyset$ , then, for  $\mathcal{A} \in \text{Br}(X) \setminus \{\ell\}$ , the image of the evaluation map  $\text{ev}_{\mathcal{A}}: X(k) \rightarrow \text{Br}(k)$  consists of one element.*

*Proof.* Let  $\mathcal{Z}$  be the largest open subscheme of  $\mathcal{X}$  that is smooth over  $\text{Spec } \mathcal{O}$ ; note that  $\mathcal{Z} \times_{\mathcal{O}} k = X$ . Write  $\mathcal{Z}_{\mathbb{F}}$  for its closed fiber, and note that  $\mathcal{Z}_{\mathbb{F}} = \mathcal{X}_{\mathbb{F}}^{\text{sm}}$ . Let  $\mathcal{Z}_{\mathbb{F}}^{(1)}$  denote the set of closed integral subvarieties of  $\mathcal{Z}_{\mathbb{F}}$  of codimension 1. In [Kat86, Prop. 1.7], Kato shows there is a complex

$$\text{Br}(X)[\ell^n] \xrightarrow{\text{res}} H^1(\mathbf{k}(\mathcal{Z}_{\mathbb{F}}), \mathbb{Z}/\ell^n \mathbb{Z}) \rightarrow \bigoplus_{Y \in \mathcal{Z}_{\mathbb{F}}^{(1)}} H^0(\mathbf{k}(Y), \mathbb{Z}/\ell^n \mathbb{Z}(-1)).$$

(In Kato's notation, take  $q = -1$ ,  $i = -2$ ,  $n \mapsto \ell^n$ , and  $X \mapsto \mathcal{Z}$ .) We claim that for  $\mathcal{A} \in \text{Br}(X)[\ell^n]$ , the residue  $\text{res}(\mathcal{A}) \in H^1(\mathbf{k}(\mathcal{Z}_{\mathbb{F}}), \mathbb{Z}/\ell^n \mathbb{Z})$  lies in the subgroup  $H_{\text{ét}}^1(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z})$ . Indeed, the group  $H^1(\mathbf{k}(\mathcal{Z}_{\mathbb{F}}), \mathbb{Z}/\ell^n \mathbb{Z})$  classifies connected cyclic covers of  $\mathcal{Z}_{\mathbb{F}}$ . By Kato's complex, the cover  $\mathcal{W} \rightarrow \mathcal{Z}_{\mathbb{F}}$  corresponding to  $\text{res}(\mathcal{A})$  is unramified in codimension one, and

hence, by the Zariski-Nagata purity theorem [SGA03, Exposé X, Théorème 3.1],  $\mathcal{W} \rightarrow \mathcal{Z}_{\mathbb{F}}$  is unramified.

The long exact sequence of low degree terms associated to the spectral sequence

$$E_2^{p,q} := H^p(\mathbb{F}, H_{\text{ét}}^q(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z})) \implies H_{\text{ét}}^{p+q}(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z})$$

starts as follows:

$$(12) \quad 0 \rightarrow H^1(\mathbb{F}, \mathbb{Z}/\ell^n \mathbb{Z}) \rightarrow H_{\text{ét}}^1(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z}) \rightarrow H_{\text{ét}}^1(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z})$$

Since, by hypothesis,  $\mathcal{Z}_{\mathbb{F}}$  has no connected unramified cyclic geometric coverings of degree  $\ell$ , we have  $H_{\text{ét}}^1(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z}) = 0$ .

We claim there is an element  $\alpha \in \text{Br}(k)\{\ell\}$  such that  $\mathcal{A} - \alpha$  has trivial residues along any codimension one subvariety of  $\mathcal{Z}$ . Such a residue depends only on the local ring at the generic point of the subvariety, so it suffices to consider residues along codimension one subvarieties of generic fiber  $X$  of  $\mathcal{Z} \rightarrow \text{Spec } \mathcal{O}$ , as well as the residue along the generic point of the special fiber  $\mathcal{Z}_{\mathbb{F}}$ . Since  $\mathcal{A}$  belongs to  $\text{Br}(X)$ , it can have a nonzero residue only at the generic point of the special fiber  $\mathcal{Z}_{\mathbb{F}}$ . As above, this residue lies in  $H_{\text{ét}}^1(\mathcal{Z}_{\mathbb{F}}, \mathbb{Z}/\ell^n \mathbb{Z})$ . On the other hand, local class field theory shows that the invariant map  $\text{Br}(k)[\ell^n] \rightarrow H^1(\mathbb{F}, \mathbb{Z}/\ell^n \mathbb{Z})$  is an isomorphism. We pick  $\alpha \in \text{Br}(k)\{\ell\}$  so that its invariant in  $H^1(\mathbb{F}, \mathbb{Z}/\ell^n \mathbb{Z})$  coincides with the preimage of  $\text{res}(\mathcal{A})$  in  $H^1(\mathbb{F}, \mathbb{Z}/\ell^n \mathbb{Z})$  for the map in (12).

By Gabber's purity theorem [Fuj02], it follows that  $\mathcal{A} - \alpha \in \text{Br}(\mathcal{Z})\{\ell\} \subseteq \text{Br}(X)\{\ell\}$ . A valuation argument shows that  $X(k) = \mathcal{X}(\mathcal{O}) = \mathcal{Z}(\mathcal{O})$ ; see [Sko96, proof of Lemma 1.1(b)]. Since  $\text{Br}(\mathcal{O}) = 0$ , we conclude that the images of the evaluation maps  $\text{ev}_{\mathcal{A}}$  and  $\text{ev}_{\alpha}$  in  $\text{Br}(k)$  coincide; the latter consists of one element.  $\square$

**Lemma 4.2.** *Suppose that  $p \neq 2$ . Let  $X$  be a K3 surface defined over  $k$ , and let  $\pi: \mathcal{X} \rightarrow \text{Spec } \mathcal{O}$  be a flat proper morphism from a regular scheme with  $X = \mathcal{X} \times_{\mathcal{O}} k$ . Assume that the singular locus of the closed fiber  $\mathcal{X}_0 := \mathcal{X}_{\mathbb{F}}$  has  $r < 8$  points, each of which is an ordinary double point. Then the smooth locus  $U \subset \mathcal{X}_0$  has no connected unramified cyclic covers of prime degree  $\ell \neq p$ .*

*Proof.* Consider an algebraically closed field  $F$  of characteristic different from  $\ell$ . Let  $Y$  be a separated integral scheme over  $F$  with  $\Gamma(Y, \mathcal{O}_Y^*) = F^*$ ; this is the case if  $Y$  is proper, or a dense open subset of a proper scheme with complement of codimension  $\geq 2$ . Then degree  $\ell$  cyclic étale covers of  $Y$  are classified by  $H_{\text{ét}}^1(Y, \mu_{\ell})$  [Mil80, ch.III]. The Kummer exact sequence [Mil80, p.125] implies that  $H_{\text{ét}}^1(Y, \mu_{\ell}) = \text{Pic}(Y)[\ell]$ , the  $\ell$ -torsion subgroup.

Combining the canonical homomorphism from the Picard group to the Weil class group and the restriction homomorphism on class groups yields

$$\text{Pic}(\mathcal{X}_0) \subset \text{Cl}(\mathcal{X}_0) \simeq \text{Cl}(U) \simeq \text{Pic}(U).$$

The quotient  $\text{Pic}(U)/\text{Pic}(\mathcal{X}_0)$  is two-torsion. Indeed, ordinary double points are étale locally isomorphic to quadric cones, whose local class group equals  $\mathbb{Z}/2\mathbb{Z}$  (generated by the ruling). Thus for each closed point  $x \in \mathcal{X}_0$ , the quotient  $\text{Pic}(\text{Spec } \mathcal{O}_{\mathcal{X}_0, x} \setminus \{x\})/\text{Pic}(\text{Spec } \mathcal{O}_{\mathcal{X}_0, x})$  is annihilated by two [Lip69, §14]. If  $\ell \neq 2$  then this computation shows that  $\text{Pic}(U)[\ell] = \text{Pic}(\mathcal{X}_0)[\ell]$ , whence degree  $\ell$  cyclic étale covers of  $U$  extend to  $\mathcal{X}_0$ .

We claim that  $\text{Pic}(\mathcal{X}_0)[\ell] = 0$  for each prime  $\ell \neq p$ . To prove this, replace  $k$  by the ramified quadratic extension  $k'$  with ring of integers  $\mathcal{O}'$ , so that  $\mathcal{X}' = \mathcal{X} \times_{\mathcal{O}} \mathcal{O}'$  is singular over the

double points of  $\mathcal{X}_0$ . Concretely, given  $\mathfrak{p}$  a uniformizer of  $\mathcal{O}$ ,  $\mathfrak{p}' = \sqrt{\mathfrak{p}}$  the corresponding uniformizer of  $\mathcal{O}'$ , and  $x \in \mathcal{X}_0$  an ordinary double point, then the étale local equation of  $\mathcal{X}$

$$\mathfrak{p} = uv + w^2$$

pulls back to

$$(\mathfrak{p}')^2 = uv + w^2.$$

Let  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}'$  denote the blow-up of the resulting singularities, with central fiber the union of the proper transform of  $\mathcal{X}_0$  and the exceptional divisors

$$\tilde{\mathcal{X}}_0 = S \cup E_1 \cup \cdots \cup E_r, \quad E_j \simeq \mathbb{P}^1 \times \mathbb{P}^1.$$

(At the cost of passing to an algebraic space, we could blow down  $E_1, \dots, E_r$  along one of the rulings of  $\mathbb{P}^1 \times \mathbb{P}^1$ .) Note that  $S$  is the K3 surface obtained by resolving the ordinary double points of  $\mathcal{X}_0$ . The pull-back homomorphism

$$\text{Pic}(\mathcal{X}_0) \rightarrow \text{Pic}(S)$$

is injective since we may regard  $U$  as an open subset of both  $\mathcal{X}_0$  and  $S$ . However,  $\text{Pic}(S)$  has no  $\ell$ -torsion: the specialization homomorphism [Ful98, §20.3]

$$\text{Pic}(X_{\bar{k}}) \rightarrow \text{Pic}(S)$$

is injective and any torsion of its cokernel is annihilated by  $p$  [MP09, Prop. 3.6].

We now focus on the case  $\ell = 2$ . Let  $F_1, \dots, F_r$  denote the exceptional divisors of  $S \rightarrow \mathcal{X}_0$ , which satisfy

$$F_1^2 = \cdots = F_r^2 = -2, \quad F_i F_j = 0, i \neq j,$$

because  $\mathcal{X}_0$  has ordinary double points.

There may exist étale double covers of  $U$  that fail to extend to étale covers of  $\mathcal{X}_0$ . Given an étale double cover  $V \rightarrow U$ , let  $\varpi : T \rightarrow S$  denote the normalization of  $S$  in the function field of  $V$ . Since  $T$  is normal,  $\varpi$  is a flat morphism [Eis95, Ex. 18.17], étale away from  $F_1 \cup \cdots \cup F_r$ . Moreover, by purity of the branch locus,  $\varpi$  is branched over some subset

$$\{F_{j_1}, \dots, F_{j_s}\} \subset \{F_1, \dots, F_r\}.$$

Since the characteristic is odd,  $\varpi$  is simply branched over these curves. Consequently,  $\sum_{i=1}^s F_{j_i} = 2D$  for some  $D \in \text{Pic}(S)$ , hence  $4D^2 = -2s$ . Since  $D^2$  is even, we conclude that  $s \equiv 0 \pmod{4}$ , i.e.,  $s = 0$  or  $4$ . The case  $s = 0$  is impossible, since this would mean that  $S$  admits an étale cyclic cover with degree prime to the characteristic. The case  $s = 4$  is also impossible: By Riemann-Roch, we have

$$\chi(\mathcal{O}_S(-D)) = 1$$

but

$$h^2(\mathcal{O}_S(-D)) = h^0(\mathcal{O}_S(D)) = 0,$$

as any effective divisor supported in the  $F_j$  (like  $2D$ ) is rigid. On the other hand, since a divisor and its negative cannot both be effective, we find

$$h^0(\mathcal{O}_S(-2D)) = 0 \text{ which implies } h^0(\mathcal{O}_S(-D)) = 0.$$

Therefore  $h^1(\mathcal{O}_S(-D)) = -1$ , which is a contradiction.  $\square$

*Remark 4.3.* When  $r = 8$ , it is possible that a smooth resolution of  $\mathcal{X}_0$  is a K3 surface with a Nikulin involution, in which case the smooth locus  $U \subset \mathcal{X}_0$  has a connected unramified cyclic double cover [vGS07].

#### 4.2. Places where $\mathcal{A}$ can ramify.

**Lemma 4.4.** *Let  $X$  be a K3 surface over a number field  $k$  as in Proposition 3.3. Let  $v$  be a finite place of good reduction for  $X$ , and assume that  $v$  is not 2-adic. Then  $\mathcal{A}$  does not ramify at  $v$ . Consequently,  $\text{inv}_v \mathcal{A}(P) = 0$  for all  $P \in X(k_v)$ .*

*Proof.* We may assume without loss of generality that the coefficients of  $A, \dots, F$  are integral. Let  $\mathcal{O}_v$  be the ring of integers of  $k_v$ , and  $\mathbb{F}_v$  its residue field. Since  $X$  is smooth and proper over  $k$  and has good reduction at  $v$ , there is a smooth proper morphism  $\mathcal{X} \rightarrow \text{Spec } \mathcal{O}_v$  with  $X_{k_v} = \mathcal{X} \times_{\mathcal{O}_v} k_v$ . We will show that the class  $\mathcal{A} \otimes k_v$  can be spread out to a class in  $\text{Br}(\mathcal{X})$ . Since, by the valuative criterion of properness, we have  $\mathcal{X}(\mathcal{O}_v) = X(k_v)$ , it will follow that  $\mathcal{A}(P) \in \text{Br}(\mathcal{O}_v) = 0$  for every point  $P \in X(k_v)$ , establishing all the claims of the proposition.

Define  $\mathcal{W}$  and  $\mathcal{Y}$  over  $\mathcal{O}_v$ , respectively, by (5) and (6). The quadric surface bundle  $(q_1)_{\mathbb{F}}: \mathcal{Y}_{\mathbb{F}} \rightarrow \mathbb{P}_{\mathbb{F}}^2$  ramifies over the discriminant curve of  $\mathcal{X}_{\mathbb{F}_v} \rightarrow \mathbb{P}_{\mathbb{F}_v}^2$ , which is smooth, because  $X$  has good reduction at  $v$ . By Theorem 3.2, there exists a smooth  $\mathbb{P}^1$  bundle  $\mathcal{F} \rightarrow \mathcal{X}$ , whose corresponding two-torsion class in  $\text{Br}(\mathcal{X})$  is represented by the quaternion algebra  $(B^2 - 4AB, A)$ , by the discussion following Theorem 3.2. Thus  $\mathcal{A} \in \text{Br}(\mathcal{X})$ , as claimed.  $\square$

**4.3. Real and 2-adic invariants.** In this section we use the notation of Proposition 3.3, specializing to the case  $k = \mathbb{Q}$ . The following lemma gives a sufficient condition to guarantee that the local invariants of  $\mathcal{A}$  at real points of  $X$  are always non-trivial.

**Lemma 4.5.** *Suppose that the quadratic forms  $A, B, C, D, E$  and  $F$  satisfy*

- (1)  $A, D$  and  $F$  are negative definite,
- (2)  $B, C$  and  $E$  are positive definite.

*Then, for any real point of  $X$ , we have*

$$M_A > 0, \quad M_D > 0 \quad \text{and} \quad M_F > 0.$$

*Proof.* First, observe that we can write  $\frac{1}{2} \det(M)$  as

$$(13) \quad A \cdot M_A - (C^2 D + B^2 F - BCE).$$

Let  $P$  be a real point of  $X$ , so that  $\frac{1}{2} \det(M) \leq 0$  holds at  $P$ . Our hypotheses on  $A, \dots, F$  imply that

$$(C^2 D + B^2 F - BCE)(P) < 0.$$

Suppose first that  $M_A \leq 0$ . Then at  $P$  we have

$$\frac{1}{2} \det(M) = \underbrace{A}_{<0} \cdot \underbrace{M_A}_{\leq 0} - \underbrace{(C^2 D + B^2 F - BCE)}_{<0} > 0,$$

a contradiction. Hence  $M_A > 0$  at  $P$ . A similar argument shows the remaining two cases.  $\square$

**Corollary 4.6.** *Suppose the hypotheses of Lemma 4.5 hold. Then the local invariant of  $\mathcal{A}$  at every real point of  $X$  is nontrivial.*

*Proof.* It suffices to show that, for any real point  $P$  of  $X$ , there is a quaternion algebra representing  $\mathcal{A}$  whose entries are both negative at  $P$ . Using the six representatives (11) of  $\mathcal{A}$ , together with Lemma 4.5, the result follows.  $\square$

Next, we write down a sufficient condition to guarantee that the local invariant map on  $\mathcal{A}$  is constant and trivial on 2-adic points. Write  $v_2: \mathbb{Q}_2 \rightarrow \mathbb{Z} \cup \{\infty\}$  for the standard 2-adic valuation. Recall that  $a \in \mathbb{Q}_2^\times$  is a square if and only if  $v_2(a)$  is even and if  $a/2^{v_2(a)} \equiv 1 \pmod{8}$ .

Let  $P = [x_0 : x_1 : x_2 : w]$  denote a 2-adic point of  $X$ . We may assume without loss of generality that  $x_0, x_1$  or  $x_2$  are elements of  $\mathbb{Z}_2$ , at least one of which is a 2-adic unit. Suppose first that  $x_0$  is a 2-adic unit, so that  $v_2(x_0) = 0$ . We use the representative  $(-M_F, A)$  of  $\mathcal{A}$  to evaluate invariants at  $P$ . Write

$$A = A_1x_0^2 + A_2x_0x_1 + A_3x_0x_2 + A_4x_1^2 + A_5x_1x_2 + A_6x_2^2,$$

and suppose that the coefficients of  $A$  satisfy

$$A_1 \equiv 1 \pmod{8}, \quad \text{and} \quad v_2(A_i) \geq 3 \text{ for } i = 2, \dots, 6.$$

Then, at  $P$ , we have  $A \equiv 1 \pmod{8}$  (since  $v_2(x_0) = 0$ ) so  $A$  is a 2-adic square. It follows that  $\text{inv}_2 \mathcal{A}(P) = 0$ , provided that  $M_F(P) \neq 0$ . To ensure this, we impose restrictions on the coefficients of the quadratic form

$$B = B_1x_0^2 + B_2x_0x_1 + B_3x_0x_2 + B_4x_1^2 + B_5x_1x_2 + B_6x_2^2.$$

Suppose that

$$v_2(B_1) = 0, \quad \text{and} \quad v_2(B_i) \geq 1 \text{ for } i = 2, \dots, 6.$$

Then, since  $v_2(x_0) = 0$ , it follows that

$$v_2(M_F(P)) = v_2(B(P)) = 0$$

and hence  $M_F \neq 0$  at  $P$ .

To ensure that 2-adic invariants of  $\mathcal{A}$  are trivial at points where  $v_2(x_1) = 0$ , we use the representative  $(-M_A, D)$  of  $\mathcal{A}$  and constrain the coefficients of  $D$  and  $E$ , respectively, in a manner analogous to how we constrained the coefficients of  $A$  and  $B$ . We proceed similarly for 2-adic points with  $v_2(x_2) = 0$ , this time using the representative  $(-M_D, F)$  and we constrain the coefficients of  $C$  and  $F$ . We summarize our discussion in the following lemma.

**Lemma 4.7.** *Write*

$$\begin{aligned} A &= A_1x_0^2 + A_2x_0x_1 + A_3x_0x_2 + A_4x_1^2 + A_5x_1x_2 + A_6x_2^2, \\ B &= B_1x_0^2 + B_2x_0x_1 + B_3x_0x_2 + B_4x_1^2 + B_5x_1x_2 + B_6x_2^2, \\ C &= C_1x_0^2 + C_2x_0x_1 + C_3x_0x_2 + C_4x_1^2 + C_5x_1x_2 + C_6x_2^2, \\ D &= D_1x_0^2 + D_2x_0x_1 + D_3x_0x_2 + D_4x_1^2 + D_5x_1x_2 + D_6x_2^2, \\ E &= E_1x_0^2 + E_2x_0x_1 + E_3x_0x_2 + E_4x_1^2 + E_5x_1x_2 + E_6x_2^2, \\ F &= F_1x_0^2 + F_2x_0x_1 + F_3x_0x_2 + F_4x_1^2 + F_5x_1x_2 + F_6x_2^2. \end{aligned}$$

*Suppose that the coefficients of these quadratic forms satisfy:*

- (1)  $A_1 \equiv 1 \pmod{8}$ , and  $v_2(A_i) \geq 3$  for  $i \neq 1$ .
- (2)  $v_2(B_1) = 0$ , and  $v_2(B_i) \geq 1$  for  $i \neq 1$ .
- (3)  $v_2(C_6) = 0$ , and  $v_2(C_i) \geq 1$  for  $i \neq 6$ .
- (4)  $D_4 \equiv 1 \pmod{8}$ , and  $v_2(D_i) \geq 3$  for  $i \neq 4$ .

- (5)  $v_2(E_4) = 0$ , and  $v_2(E_i) \geq 1$  for  $i \neq 4$ .  
(6)  $F_6 \equiv 1 \pmod{8}$ , and  $v_2(F_i) \geq 3$  for  $i \neq 6$ .

Then, for every 2-adic point  $P$  of  $X$ , we have  $\text{inv}_2 \mathcal{A}(P) = 0$ . □

## 5. AN EXAMPLE

Consider the quadrics

$$\begin{aligned}
(14) \quad A &:= -7x_0^2 - 16x_0x_1 + 16x_0x_2 - 24x_1^2 + 8x_1x_2 - 16x_2^2 \\
B &:= 3x_0^2 + 2x_0x_2 + 2x_1^2 - 4x_1x_2 + 4x_2^2 \\
C &:= 10x_0^2 + 4x_0x_1 + 4x_0x_2 + 4x_1^2 - 2x_1x_2 + x_2^2 \\
D &:= -16x_0^2 + 8x_0x_1 - 23x_1^2 + 8x_1x_2 - 40x_2^2 \\
E &:= 4x_0^2 - 4x_0x_2 + 11x_1^2 - 4x_1x_2 + 6x_2^2 \\
F &:= -40x_0^2 + 32x_0x_1 - 40x_1^2 - 8x_1x_2 - 23x_2^2.
\end{aligned}$$

Let  $W \subset \text{Proj } \mathbb{Q}[x_0, x_1, x_2] \times \text{Proj } \mathbb{Q}[y_0, y_1, y_2]$  be the type  $(2, 2)$  divisor given by the vanishing of the bihomogeneous polynomial

$$\begin{aligned}
(15) \quad &A(x_0, x_1, x_2)y_0^2 + B(x_0, x_1, x_2)y_0y_1 + C(x_0, x_1, x_2)y_0y_2 \\
&+ D(x_0, x_1, x_2)y_1^2 + E(x_0, x_1, x_2)y_1y_2 + F(x_0, x_1, x_2)y_2^2.
\end{aligned}$$

As in §3, the projections  $\pi_i: W \rightarrow \mathbb{P}^2$  give conic bundle structures on  $W$  ramified over plane sextics  $C_i$ ,  $i = 1, 2$ . An equation for  $C_1$  is then given by  $f := -\frac{1}{2} \det(M) = 0$ , with  $M$  as in (8). An equation for  $C_2$  can be found analogously. The Jacobian criterion shows that both  $C_1$  and  $C_2$  are smooth; thus, for  $i = 1, 2$ , the double cover  $X_i \rightarrow \mathbb{P}^2$  ramified along  $C_i$  is a K3 surface of degree 2.

**5.1. Primes of bad reduction.** The primes of bad reduction of  $X_1$  and  $C_1$  coincide. The latter divide the generator  $m$  of the ideal obtained by saturating

$$\left\langle f, \frac{\partial f}{\partial x_0}, \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2} \right\rangle \subset \mathbb{Z}[x_0, x_1, x_2]$$

by the irrelevant ideal and eliminating  $x_0$ ,  $x_1$  and  $x_2$ . We obtain

$$\begin{aligned}
m = &1115508232640214856843363784231663793779083264535962688555888430968 \\
&8933364438401787008291918987282105867611490800785997644322303281186 \\
&8922614222749465991103128446037422257623280138072129654879995620391 \\
&0907629715637695773281604080143775185215794393627484442538367517916 \\
&8651952191024387026109016400178074232186309443422761817391984342483 \\
&34511814400.
\end{aligned}$$

Standard factorization methods quickly reveal a few small prime power factors of  $m$ :

$$m = 2^8 \cdot 5^2 \cdot 7 \cdot 89 \cdot 173 \cdot 257^2 \cdot 263 \cdot 650779^2 \cdot m'.$$

The remaining factor  $m'$  has 318 decimal digits. Factoring  $m'$  with present day mathematical and computational technology is a difficult problem. However, the presence of the second

K3 surface  $X_2$  supplies a backdoor solution: by Lemma 3.1, a prime of bad reduction for  $W$  is a prime of bad reduction for *both*  $X_1$  and  $X_2$ .

Another Groebner basis calculation shows that the primes of bad reduction of  $X_2$  divide

$$\begin{aligned} n := & 18468445386704774116897512713438756322646374324269134481315634355660 \\ & 59216198653927410468599212130905398491499555534045930594495263034981 \\ & 50100881353352665095649631677613412079293044973446406764509694053112 \\ & 10471631439070548340358668493117334582314574674926223315439909955021 \\ & 6973495867514854209929544319382116616140800 \end{aligned}$$

Again, standard factorization methods give a few small prime power factors of  $n$ :

$$n = 2^{11} \cdot 5^2 \cdot 7 \cdot 89 \cdot 173 \cdot 263 \cdot 461^2 \cdot 6547^2 \cdot n',$$

where  $n'$  has 290 decimal digits. Our observation says that we may reasonably expect that  $m'$  and  $n'$  have a large greatest common divisor (which is easily calculated using the Euclidean algorithm). This is indeed the case:

$$\begin{aligned} \gcd(m', n') := & 809147864157687938441948148614369785987783654943839689121548451 \\ & 788111145202992792430023470932052297439515068068797124401938255 \\ & 799311490342451172887433057574480263654457987109316488649107. \end{aligned}$$

Here a small miracle happens:  $\gcd(m', n')$  is a prime number! This claim is rigorously verified using elliptic curve primality proving algorithms [AM93], implemented in both SAGE and magma. We are now in a position to complete the factorization of  $m$ , and hence compute the primes of bad reduction for  $X_1$ , which are:

$$\begin{aligned} & 2, 5, 7, 89, 173, 257, 263, 650779, \\ & 521219738678096220868573969913582546660848099260319499224599922739, \\ & \gcd(m', n'). \end{aligned}$$

We note that the penultimate prime in the list above occurs with multiplicity 2 in the factorization of  $m$ . We will write  $q$  for this prime number in Table 1.

*Remark 5.1.* Our numerical experiments yield several “viable” pairs  $(X_1, \mathcal{A})$  that could be counter-examples to the Hasse principle explained by a transcendental Brauer-Manin obstruction arising from  $\mathcal{A}$ , in the following sense:  $X_1$  has geometric Picard rank 1, and we can control the real and 2-adic invariants of  $\mathcal{A}$  (using Corollary 4.6 and Lemma 4.7). Out of a dozen or so viable candidates that our initial search yielded, the example we present is the only one we found for which  $\gcd(m', n')$  is a prime number. One can obtain further examples by computing 2-adic invariants by “brute force” instead of using Lemma 4.7.

**5.2. Local points.** By the Weil Conjectures, if  $p > 22$  is a prime such that  $X_1$  has smooth reduction  $(X_1)_p$  at  $p$ , then  $(X_1)_p$  has a smooth  $\mathbb{F}_p$ -point, which can be lifted by Hensel’s lemma to a smooth  $\mathbb{Q}_p$ -point. Thus, to show  $X_1$  is locally soluble, it suffices to verify that  $X_1$  has local points at  $\mathbb{R}$  (clear), and at  $\mathbb{Q}_p$  for primes  $p \leq 19$  and primes  $p > 19$  where  $X_1$  has bad reduction. This is indeed the case: we substitute integers with small absolute value for  $x_0, x_1$ , and  $x_2$ , and check if  $-\frac{1}{2} \det(M)$  is a square in  $\mathbb{Q}_p$ . The results are recorded in Table 1.



$p$	$x_0$	$x_1$	$x_2$	$-\frac{1}{2} \det(M)$
2	0	0	-1	57872
3	-1	-1	1	1622952
5	-1	-1	-1	736256
7	-1	-1	0	256575
11	-1	-1	-1	736256
13	-1	-1	-1	736256
17	-1	-1	1	1622952
19	-1	-1	-1	736256
89	-1	0	-1	80019
173	-1	-1	0	256575
257	-1	-1	-1	736256
263	-1	-1	0	256575
650779	-1	-1	1	1622952
$q$	-1	-1	-1	736256
$\gcd(m', n')$	-1	-1	-1	736256

TABLE 1. Verifying  $X_1$  has  $\mathbb{Q}_p$ -points at small  $p$  and primes of bad reduction. The numerical entries in the rightmost column are all squares in the appropriate  $p$ -adic field.

**5.3. Picard Rank 1.** In this section we show  $X_1$  has (geometric) Picard rank 1. This will allow us to conclude that the obstruction to the Hasse principle arising from  $\mathcal{A}$  is genuinely transcendental. Until recently, the method to prove a K3 surface has odd Picard rank, devised by van Luijk and refined by Kloosterman, and Elsenhans and Jahnel [vL07, Klo07, EJ11b], required point counting over extensions of the residue field at two primes of good reduction. A recent result of Elsenhans and Jahnel allows us to prove odd Picard rank using information at two primes, but counting points over extensions of a single residue field.

**Theorem 5.2** ([EJ11c]). *Let  $f: X \rightarrow \text{Spec } \mathbb{Z}$  be a proper, flat morphism of schemes. Suppose there is a rational prime  $p \neq 2$  such that the fiber  $X_p$  of  $f$  at  $p$  satisfies  $H^1(X_p, \mathcal{O}_{X_p}) = 0$ . Then the specialization homomorphism  $\text{Pic}(X_{\mathbb{Q}}) \rightarrow \text{Pic}(X_{\mathbb{F}_p})$  has torsion-free cokernel.*

We deduce the following generalization of [EJ11c, Example 1.6].

**Proposition 5.3.** *Let  $X$  be a K3 surface of degree 2 over  $\mathbb{Q}$ , given as a double cover  $\pi: X \rightarrow \mathbb{P}^2$  ramified over a smooth plane sextic curve  $C$ . Let  $p$  and  $p'$  denote two odd primes of good reduction for  $X$ . Assume that there exists a line  $\ell$  that is tritangent to the curve  $C_p$ , and suppose further that  $\text{Pic}(\overline{X}_p)$  has rank 2 and is generated by the curves in  $\pi_p^{-1}(\ell)$ . If there are no tritangent lines to the curve  $C_{p'}$ , then  $\text{Pic}(\overline{X})$  has rank 1.*

*Proof.* Since  $\text{Pic}(\overline{X})$  injects into  $\text{Pic}(\overline{X}_p)$ , if  $\text{Pic}(\overline{X})$  has rank 2, then we claim the tritangent line  $\ell$  must lift to a tritangent line  $L$  in characteristic 0. To see this, note that by Theorem 5.2, the components of the pullback of  $\ell$  to  $\overline{X}_p$  lift to divisors  $C$  and  $C'$  on  $\overline{X}$  such that  $C^2 = C'^2 = -2$  and  $C \cdot h = C' \cdot h = 1$ , where  $h$  is the pullback of the class of a line from  $\mathbb{P}^2$ . By Riemann-Roch, either  $C$  or  $-C$  is effective. Since  $C \cdot h > 0$ , we conclude that  $C$  is effective;

likewise for  $C'$ . The projection  $L$  of  $C$  or  $C'$  to  $\mathbb{P}^2$  is a tritangent line in characteristic zero, as claimed.

Degree considerations show that  $L$  cannot break upon reduction modulo  $p'$ . This contradicts the assumption that the curve  $C_{p'}$  has no tritangent lines.  $\square$

*Remarks 5.4.* Proposition 5.3 is computationally useful because:

- (1) Checking the existence of a tritangent line modulo  $p'$  is an easy Groebner basis calculation; see [EJ08, Algorithm 8].
- (2) Given a K3 surface of degree 2 over  $\mathbb{Q}$ , we can quickly *search* for small primes  $p$  of good reduction over which the branch curve  $C_{p'}$  of the double cover  $X_{p'} \rightarrow \mathbb{P}_{\mathbb{F}_{p'}}^2$  has a tritangent line.

Our particular surface  $X_1$  reduces modulo 3 to the (smooth) K3 surface

$$w^2 = 2x_1^2(x_0^2 + 2x_0x_1 + 2x_1^2)^2 + (2x_0 + x_2)(x_0^5 + x_0^4x_1 + x_0^3x_1x_2 + x_0^2x_1^3 + x_0^2x_1^2x_2 + 2x_0^2x_2^3 + x_0x_1^4 + 2x_0x_1^3x_2 + x_0x_1^2x_2^2 + x_1^5 + 2x_1^4x_2 + 2x_1^3x_2^2 + 2x_2^5).$$

From the expression on the right hand side, it is clear that  $2x_0 + x_2 = 0$  is a tritangent line to the branch curve of the double cover. The components of the pullback of this line generate a rank 2 sublattice of  $\text{Pic}((X_1)_3)$ . Let  $N_n := \#X_1(\mathbb{F}_{3^n})$ ; counting points we find

$N_1$	$N_2$	$N_3$	$N_4$	$N_5$	$N_6$	$N_7$	$N_8$	$N_9$	$N_{10}$
7	79	703	6607	60427	532711	4792690	43068511	387466417	3486842479

This is enough information to determine the characteristic polynomial  $f$  of Frobenius on  $H^2((X_1)_{\overline{\mathbb{F}}_3}, \overline{\mathbb{Q}}_\ell)$ ; see, for example [vL07] (the sign of the functional equation for  $f$  is negative—a positive sign gives rise to roots of  $f$  of absolute value  $\neq 3$ ). Setting  $f_3(t) = 3^{-22}f(3t)$ , we obtain a factorization into irreducible factors as follows:

$$f_3(t) = \frac{1}{3}(t-1)(t+1)(3t^{20} + 3t^{19} + 5t^{18} + 5t^{17} + 6t^{16} + 2t^{15} + 2t^{14} - 3t^{13} - 4t^{12} - 8t^{11} - 6t^{10} - 8t^9 - 4t^8 - 3t^7 + 2t^6 + 2t^5 + 6t^4 + 5t^3 + 5t^2 + 3t + 3).$$

The number of roots of  $f_3(t)$  that are roots of unity give an upper bound for  $\text{Pic}((X_1)_{\overline{\mathbb{F}}_3})$  (see, e.g., [vL07, Corollary 2.3]). The roots of the degree 20 factor of  $f_3(t)$  are not integral, so they are not roots of unity. We conclude that  $\text{rk Pic}((X_1)_{\overline{\mathbb{F}}_3}) = 2$ .

A computation shows that  $X_1$  has no line tritangent to the branch curve when we reduce modulo  $p' = 11$  (see Remark 5.4(i)). Note that the surface is not smooth at  $p' = 5, 7$ . Applying Proposition 5.3, we obtain the following result.

**Proposition 5.5.** *The surface  $X_1$  has geometric Picard rank 1.*  $\square$

**5.4. Local invariants.** In this section we compute the local invariants of the algebra  $\mathcal{A}$  for our particular surface  $X_1$ .

**Proposition 5.6.** *Let  $p \leq \infty$  be a place of  $\mathbb{Q}$ . For any  $P \in X_1(\mathbb{Q}_p)$ , we have*

$$\text{inv}_p(\mathcal{A}(P)) = \begin{cases} 0, & \text{if } \mathbb{Q}_p \neq \mathbb{R}, \\ 1/2, & \text{if } \mathbb{Q}_p = \mathbb{R}. \end{cases}$$

*Proof.* Whenever  $p \neq 2$  is a finite prime of good reduction for  $X_1$ , we have  $\text{inv}_p(\mathcal{A}(P)) = 0$  for all  $P$ , by Lemma 4.4.

At every odd prime of bad reduction of  $X_1$ , the singular locus consists of  $r < 8$  ordinary double points: for most of these primes  $p$  the claim follows because the valuation at  $p$  of the discriminant of  $X_1$  is one, by our work in §5.1, so the singular locus consists of a single ordinary double point. For the remaining primes, a straightforward computer calculation does the job.

Together with Proposition 4.1 and Lemma 4.2, this implies that  $\text{inv}_p(\mathcal{A}(P))$  is independent of  $P$ ; it thus suffices to evaluate these invariants at a single point  $P$ . We use the local points listed in Table 1 to verify that all the local invariants vanish.

Finally, the quadrics (14) are readily seen to satisfy the hypotheses of Lemmas 4.5 and 4.7, which establishes the claim for real and 2-adic points of  $X_1$ , using Corollary 4.6.  $\square$

**5.5. Proof of Theorem 1.1.** The first part of the Theorem is just Proposition 3.3. We specialize now to the case  $k = \mathbb{Q}$ .

Let  $A, \dots, F$  be as in (14), so that  $X$  is the surface  $X_1$  considered throughout this section. The cohomology group  $H^1(\mathbb{Q}, \text{Pic}(\overline{X}))$  is trivial, because  $\text{Pic}(\overline{X}) \cong \mathbb{Z}$ , with trivial Galois action, by Proposition 5.5. By (1), we have  $\text{Br}_1(X) = \text{Br}_0(X)$ . Hence, the class  $\mathcal{A} \in \text{Br}(X)$  is transcendental, if it is not constant.

We established in §5.2 that  $X(\mathbb{A}) \neq \emptyset$ . On the other hand,  $X(\mathbb{A})^{\mathcal{A}} = \emptyset$ , by Proposition 5.6. This shows that  $\mathcal{A}$  is nonconstant, and that  $X(\mathbb{A})^{\text{Br}} = \emptyset$ .  $\square$

## 6. COMPUTATIONS

In the interest of transparency, we briefly outline the computations that led to the example witnessing the second part of Theorem 1.1. The basic idea is to construct “random” K3 surfaces of the form (2), and perform a series of tests that guarantee the statement of Theorem 1.1 holds. Any surface left over after Step 7 below is a witness to this theorem.

**Step 1: Seed polynomials.** Generate random homogeneous quadratic polynomials

$$A, B, C, D, E, \text{ and } F \in \mathbb{Z}[x_0, x_1, x_2],$$

with coefficients in a suitable range, subject to the constraints imposed by the hypotheses of Lemma 4.7. We also require that the signs of  $x_0^2$ ,  $x_1^2$  and  $x_2^2$  are positive for  $B$ ,  $C$  and  $E$ , and negative for  $A$ ,  $D$  and  $F$ , to improve the chances that the hypotheses of Lemma 4.5 are satisfied. If these hypotheses are not satisfied, then start over.

**Step 2: Smoothness.** Compute  $f := -\frac{1}{2} \det(M)$ , where  $M$  is the matrix in (8). This is an equation for the curve  $C_1$ . Use the Jacobian criterion to check smoothness of  $C_1$  over  $\mathbb{Q}$  and  $\mathbb{F}_3$  (the latter will be needed to certify that the K3 surface  $X_1$  has Picard rank 1). If either condition is not satisfied, then start over.

**Step 3: Tritangent lines.** Here we have the hypotheses of Proposition 5.3 in mind. Over  $\mathbb{F}_3$ , use [EJ08, Algorithm 8] to test for the existence of a tritangent line to  $C_1$ . Let

$$S := \{p : 5 \leq p \leq 100 \text{ a prime of good reduction for } C_1\}.$$

Find  $p \in S$ , such that  $C_1$  over  $\mathbb{F}_p$  has no tritangent line. If either test fails, then start over.

**Step 4: Local points.** For primes  $p \leq 22$  and  $p = \infty$ , test for  $\mathbb{Q}_p$ -points of  $X_1/\mathbb{Q} : w^2 = f$  by plugging in integers with small absolute value (typically 1 or 2) for  $x_0, x_1$  and  $x_2$ , and determining whether  $f$  is a  $p$ -adic square. If this test fails, then it is plausible that  $X_1$  has no local points (false negatives are certainly possible); start over.

**Step 5: Point Counting.** Use [EJ08, Algorithm 15] to determine  $X_1(\mathbb{F}_{3^n})$  for  $n = 1, \dots, 10$ . This algorithm counts *Galois orbits* of points, saving a factor of  $n$  when counting  $\mathbb{F}_{3^n}$ -points. Use [EJ08, Algorithms 21 and 23] to determine an upper bound  $\rho_{\text{up}}$  for the geometric Picard number of the surface  $X_1$  over  $\mathbb{F}_3$ . If  $\rho_{\text{up}} > 2$ , then start over. Otherwise, Proposition 5.3 guarantees that  $X_1$  has geometric Picard number 1, by our work in Step 3.

**Step 6: Primes of bad reduction.** Proceeding as in the beginning of §5.1, compute the integer  $m$  whose prime factors give the places of bad reduction of  $C_1$  (and hence those of  $X_1$ ). Compute an equation for  $C_2$ , as well as the analogous integer  $n$  giving its primes of bad reduction. Typically,  $m$  and  $n$  will be very large. Proceed as in §5.1 to factorize them. If the factorization is not feasible (e.g., the integer  $\gcd(m', n')$  as in §5.1 not is prime), then start over.

**Step 7: Computations at places of bad reduction.** At odd places of bad reduction, check for local points, as in Step 4. Determine the (geometric) singular locus. If at any prime in question the locus does not consist of  $r < 8$  ordinary double points, then start over. Use to the local points found to compute the (constant) value the invariants of  $\mathcal{A}$  takes at these places. If there is no Brauer-Manin obstruction, then start over.

## REFERENCES

- [AM93] A. O. L. Atkin and F. Morain. Elliptic curves and primality proving. *Math. Comp.*, 61(203):29–68, 1993.
- [BBFL07] M. J. Bright, N. Bruin, E. V. Flynn, and A. Logan. The Brauer-Manin obstruction and Sh[2]. *LMS J. Comput. Math.*, 10:354–377 (electronic), 2007.
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [Bea77] Arnaud Beauville. Variétés de Prym et jacobiniennes intermédiaires. *Ann. Sci. École Norm. Sup. (4)*, 10(3):309–391, 1977.
- [Bri06] Martin Bright. Brauer groups of diagonal quartic surfaces. *J. Symbolic Comput.*, 41(5):544–558, 2006.
- [BSD75] B. J. Birch and H. P. F. Swinnerton-Dyer. The Hasse problem for rational surfaces. *J. Reine Angew. Math.*, 274/275:164–174, 1975. Collection of articles dedicated to Helmut Hasse on his seventy-fifth birthday, III.
- [Cor07] Patrick Corn. The Brauer-Manin obstruction on del Pezzo surfaces of degree 2. *Proc. Lond. Math. Soc. (3)*, 95(3):735–777, 2007.
- [Cor10] Patrick Corn. Tate-Shafarevich groups and  $K3$  surfaces. *Math. Comp.*, 79(269):563–581, 2010.
- [CTCS80] Jean-Louis Colliot-Thélène, Daniel Coray, and Jean-Jacques Sansuc. Descente et principe de Hasse pour certaines variétés rationnelles. *J. Reine Angew. Math.*, 320:150–191, 1980.

- [CTKS87] Jean-Louis Colliot-Thélène, Dimitri Kanevsky, and Jean-Jacques Sansuc. Arithmétique des surfaces cubiques diagonales. In *Diophantine approximation and transcendence theory (Bonn, 1985)*, volume 1290 of *Lecture Notes in Math.*, pages 1–108. Springer, Berlin, 1987.
- [CTS] Jean-Louis Colliot-Thélène and Alexei N. Skorobogatov. Sur le groupe de Brauer transcendant. arXiv:1106.6312.
- [CTS13] Jean-Louis Colliot-Thélène and Alexei N. Skorobogatov. Good reduction of the Brauer–Manin obstruction. *Trans. Amer. Math. Soc.*, 365(2):579–590, 2013.
- [CTSSD87] Jean-Louis Colliot-Thélène, Jean-Jacques Sansuc, and Peter Swinnerton-Dyer. Intersections of two quadrics and Châtelet surfaces. II. *J. Reine Angew. Math.*, 374:72–168, 1987.
- [Cun07] Stephen Cunnane. Rational points on Enriques surfaces, 2007. Ph. D. thesis, Imperial College London.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [EJ08] Andreas-Stephan Elsenhans and Jörg Jahnel.  $K3$  surfaces of Picard rank one and degree two. In *Algorithmic number theory*, volume 5011 of *Lecture Notes in Comput. Sci.*, pages 212–225. Springer, Berlin, 2008.
- [EJ10a] Andreas-Stephan Elsenhans and Jörg Jahnel. Cubic surfaces with a Galois invariant double-six. *Cent. Eur. J. Math.*, 8(4):646–661, 2010.
- [EJ10b] Andreas-Stephan Elsenhans and Jörg Jahnel. On the Brauer–Manin obstruction for cubic surfaces. *J. Comb. Number Theory*, 2(2):107–128, 2010.
- [EJ11a] Andreas-Stephan Elsenhans and Jörg Jahnel. Cubic surfaces with a Galois invariant pair of Steiner trihedra. *Int. J. Number Theory*, 7(4):947–970, 2011.
- [EJ11b] Andreas-Stephan Elsenhans and Jörg Jahnel. On the computation of the Picard group for  $K3$  surfaces. *Math. Proc. Cambridge Philos. Soc.*, 151(2):263–270, 2011.
- [EJ11c] Andreas-Stephan Elsenhans and Jörg Jahnel. The Picard group of a  $K3$  surface and its reduction modulo  $p$ . *Algebra Number Theory*, 5(8):1027–1040, 2011.
- [Fuj02] Kazuhiro Fujiwara. A proof of the absolute purity conjecture (after Gabber). In *Algebraic geometry 2000, Azumino (Hotaka)*, volume 36 of *Adv. Stud. Pure Math.*, pages 153–183. Math. Soc. Japan, Tokyo, 2002.
- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, second edition, 1998.
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [GS06] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*, volume 101 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006.
- [Har96] David Harari. Obstructions de Manin transcendentes. In *Number theory (Paris, 1993–1994)*, volume 235 of *London Math. Soc. Lecture Note Ser.*, pages 75–87. Cambridge Univ. Press, Cambridge, 1996.
- [HS05] David Harari and Alexei Skorobogatov. Non-abelian descent and the arithmetic of Enriques surfaces. *Int. Math. Res. Not.*, 52:3203–3228, 2005.
- [HVAV11] Brendan Hassett, Anthony Várilly-Alvarado, and Patrick Varilly. Transcendental obstructions to weak approximation on general  $K3$  surfaces. *Adv. Math.*, 228(3):1377–1404, 2011.
- [Ier10] Evis Ieronymou. Diagonal quartic surfaces and transcendental elements of the Brauer groups. *J. Inst. Math. Jussieu*, 9(4):769–798, 2010.
- [ISZ11] Evis Ieronymou, Alexei N. Skorobogatov, and Yuri G. Zarhin. On the Brauer group of diagonal quartic surfaces. *J. London Math. Soc. (2)*, 83(3):659–672, 2011.
- [Kat86] Kazuya Kato. A Hasse principle for two-dimensional global fields. *J. Reine Angew. Math.*, 366:142–183, 1986. With an appendix by Jean-Louis Colliot-Thélène.
- [Klo07] Remke Kloosterman. Elliptic  $K3$  surfaces with geometric Mordell–Weil rank 15. *Canad. Math. Bull.*, 50(2):215–226, 2007.
- [KT04] Andrew Kresch and Yuri Tschinkel. On the arithmetic of del Pezzo surfaces of degree 2. *Proc. London Math. Soc. (3)*, 89(3):545–569, 2004.

- [KT08] Andrew Kresch and Yuri Tschinkel. Effectivity of Brauer-Manin obstructions. *Adv. Math.*, 218(1):1–27, 2008.
- [Lip69] Joseph Lipman. Rational singularities, with applications to algebraic surfaces and unique factorization. *Inst. Hautes Études Sci. Publ. Math.*, (36):195–279, 1969.
- [Log08] Adam Logan. The Brauer-Manin obstruction on del Pezzo surfaces of degree 2 branched along a plane section of a Kummer surface. *Math. Proc. Cambridge Philos. Soc.*, 144(3):603–622, 2008.
- [LP81] Eduard Looijenga and Chris Peters. Torelli theorems for Kähler  $K3$  surfaces. *Compositio Math.*, 42(2):145–186, 1980/81.
- [LvL09] Adam Logan and Ronald van Luijk. Nontrivial elements of Sha explained through  $K3$  surfaces. *Math. Comp.*, 78(265):441–483, 2009.
- [Man71] Y. I. Manin. Le groupe de Brauer-Grothendieck en géométrie diophantienne. In *Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1*, pages 401–411. Gauthier-Villars, Paris, 1971.
- [Man74] Yu. I. Manin. *Cubic forms: algebra, geometry, arithmetic*. North-Holland Publishing Co., Amsterdam, 1974. Translated from Russian by M. Hazewinkel, North-Holland Mathematical Library, Vol. 4.
- [Mil80] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [MP09] Davesh Maulik and Bjorn Poonen. Néron-Severi groups under specialization, 2009. arXiv:0907.4781.
- [Muk84] Shigeru Mukai. Symplectic structure of the moduli space of sheaves on an abelian or  $K3$  surface. *Invent. Math.*, 77(1):101–116, 1984.
- [Nik79] V. V. Nikulin. Integer symmetric bilinear forms and some of their geometric applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(1):111–177, 238, 1979.
- [Pre10] Thomas Preu. Transcendental Brauer-Manin obstruction for a diagonal quartic surface, 2010. Ph. D. thesis, Universität Zürich.
- [S<sup>+</sup>09] W. A. Stein et al. *Sage Mathematics Software (Version 4.2.1)*. The Sage Development Team, 2009. <http://www.sagemath.org>.
- [SD93] Peter Swinnerton-Dyer. The Brauer group of cubic surfaces. *Math. Proc. Cambridge Philos. Soc.*, 113(3):449–460, 1993.
- [SD99] Peter Swinnerton-Dyer. Brauer-Manin obstructions on some Del Pezzo surfaces. *Math. Proc. Cambridge Philos. Soc.*, 125(2):193–198, 1999.
- [Ser73] J.-P. Serre. *A course in arithmetic*. Springer-Verlag, New York, 1973. Translated from the French, Graduate Texts in Mathematics, No. 7.
- [SGA03] *Revêtements étales et groupe fondamental (SGA 1)*. Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 3. Société Mathématique de France, Paris, 2003. Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960–61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)].
- [Sko96] Alexei N. Skorobogatov. Descent on fibrations over the projective line. *Amer. J. Math.*, 118(5):905–923, 1996.
- [Sko01] Alexei Skorobogatov. *Torsors and rational points*, volume 144 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2001.
- [SSD05] Alexei Skorobogatov and Peter Swinnerton-Dyer. 2-descent on elliptic curves and rational points on certain Kummer surfaces. *Adv. Math.*, 198(2):448–483, 2005.
- [SZ08] Alexei N. Skorobogatov and Yuri G. Zarhin. A finiteness theorem for the Brauer group of abelian varieties and  $K3$  surfaces. *J. Algebraic Geom.*, 17(3):481–502, 2008.
- [SZ12] Alexei N. Skorobogatov and Yuri G. Zarhin. The Brauer group of Kummer surfaces and torsion of elliptic curves. *J. Reine Angew. Math.*, 666:115–140, 2012.
- [VA08] Anthony Várilly-Alvarado. Weak approximation on del Pezzo surfaces of degree 1. *Adv. Math.*, 219(6):2123–2145, 2008.

- [vG05] Bert van Geemen. Some remarks on Brauer groups of  $K3$  surfaces. *Adv. Math.*, 197(1):222–247, 2005.
- [vGS07] Bert van Geemen and Alessandra Sarti. Nikulin involutions on  $K3$  surfaces. *Math. Z.*, 255(4):731–753, 2007.
- [vL07] Ronald van Luijk.  $K3$  surfaces with Picard number one and infinitely many rational points. *Algebra Number Theory*, 1(1):1–15, 2007.
- [Voi86] Claire Voisin. Théorème de Torelli pour les cubiques de  $\mathbf{P}^5$ . *Invent. Math.*, 86(3):577–601, 1986.
- [Wit] Olivier Wittenberg. Personal letter, April 27th, 2010.
- [Wit04] Olivier Wittenberg. Transcendental Brauer-Manin obstruction on a pencil of elliptic curves. In *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002)*, volume 226 of *Progr. Math.*, pages 259–267. Birkhäuser Boston, Boston, MA, 2004.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX 77005, USA

*E-mail address:* `hassett@rice.edu`

*URL:* `http://www.math.rice.edu/~hassett`

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX 77005, USA

*E-mail address:* `varilly@rice.edu`

*URL:* `http://www.math.rice.edu/~av15`