

The Surface Subgroup Theorem

and

The Ehrenpreis Conjecture

by Jeremy Kahn

Joint with Vladimir Markovic

Theorem (KM) (Surface subgroup conjecture)


Let M be a closed hyperbolic 3-manifold. Then $\pi_1(M)$ has a surface subgroup (the isomorphic image of π_1 of a hyperbolic surface),

Theorem (KM) (Ehrenpreis conjecture)


Let S and T be closed hyperbolic Riemann surfaces. Then

S and T have nearby finite covers (in the Teichmüller metric).


2D




3D



Building Covers of the Circle




degree 2




degree 4


We can join immersed
1-manifolds with boundary
to form a cover of a
closed 1-manifold (a circle).

We can join immersed orientable n -manifolds with boundary to form a cover of a closed orientable n -manifold.



We can build more than one cover







We can
fail to
build a
cover

Given a collection $C = \sum n_i \alpha_i$ of 1-manifolds, immersed in a circle, We can assemble C to form a cover if and only if


$$\partial C \equiv \sum n_i \partial \alpha_i = 0$$




$$\partial C = 0$$




$$\partial C = 0$$









$$\partial C \neq 0$$




We can always build a piecewise immersed 1-manifold in a closed 2-manifold if we use the "doubling trick"






A set of points ~~at~~ on S' is
 "evenly distributed to the ~~set~~ scale ε "
 if every ε -interval of S' has
 the same number of points, ~~to~~ up
 to a factor of $1+\varepsilon$.




$$\frac{103}{98} < 1 + \varepsilon$$


We can make the piecewise immersed 1-manifold nearly geodesic (small bending) if the segments are "evenly distributed" around each meeting point:




A (hyperbolic) pair of pants
is a hyperbolic 2-manifold with
geodesic boundary that is
diffeomorphic to the following.




We obtain a closed hyperbolic surface when we join pairs of pants



Hyperbolic Pairs of Pants




double
→




Half-lengths L_1, L_2, L_3

Perfect Pants and Perfect Surfaces




$$hl(C) = R$$




$$hl(C) = R$$
$$s(C) = 1$$

Theorem Any two closed ~~surfaces~~ perfect surfaces have a common finite cover.

Good Pants and Good Surfaces



$$|h(c) - R| < \varepsilon$$



$$|s(c) - 1| < \varepsilon/R$$


Theorem (KM) ~~Ann~~

A good closed pants surface is close to a perfect surface.

Theorem (KM)

For all $\varepsilon < \varepsilon_0$, and $R > R_0$,
any (ε, R) good surface
is $10^{12}\varepsilon$ -close to
an R -perfect surface
in the Teichmüller metric.

Good Covers $\Rightarrow$ Ehrenpreis




An Immersed Pair of Pants



We can think of the pants as
being isometrically immersed, and
speak of the goodness of pants

Gluing Together Immersed Pants



We can ~~ind in~~ identify cuffs of
the parametrizing pants whenever
the cuffs map to the same ~~ge~~-geodesic γ ,
and the pants are on opposite sides
of γ .

Easy Theorem

Let $\mathcal{Q}$ be a finite set of immersed pants, such that for every closed geodesic γ on S we have the same number of pants in $\mathcal{Q}$ on both sides of γ . (That number can be zero). Then we can assemble $\mathcal{Q}$ to form a finite cover of S .


Good Pants and Good Curves

We let



$$\Gamma_{\varepsilon, R} = \left\{ \gamma \mid \begin{array}{l} \gamma \text{ is a closed geodesic on } S \\ |hl(\gamma) - R| < \varepsilon \end{array} \right\}$$

$$\Pi_{\varepsilon, R} = \left\{ P \mid \begin{array}{l} P \text{ is an immersed pair} \\ \text{of pants in } S, \text{ and} \\ \partial P \subseteq \Gamma_{\varepsilon, R} \end{array} \right\}$$

The square root of a geodesic



Shear Coordinates and $N^4(\sqrt{\gamma})$



The Equidistribution Theorem

For every good curve γ , and large R ,


$$\{\text{feet } p \mid p \in \pi_{\varepsilon, R} \text{ and } \gamma \in \partial p\} \leq N^1(\sqrt{\gamma})$$

is evenly distributed on $N^1(\sqrt{\gamma})$
to the scale e^{-gR} , for $g = g(S, \varepsilon)$.

(We assume that $R > R_0(S, \varepsilon)$).


Equidistributed & Balanced $\Rightarrow$ Good Cover

If there were exactly the same number of pants on both sides of each geodesic γ , then we would be able to assemble the pants in $\mathbb{T}_{\varepsilon, R}$ to form a good cover of S :




Equidistributed $\Rightarrow$ Nearly balanced

It follows from equidistribution that there are nearly the same number of points in $\Pi_{\varepsilon, R}$ on either side of a good geodesic γ



An Interlude in Three Dimensions


In a hyperbolic 3-manifold $N^1(\delta)$ (and $N^1(\sqrt{\delta})$) is connected, so we can use the "doubling trick" to build a nearly geodesic immersed surface (which is therefore essential).



$$\pi_1(f): \pi_1(S) \rightarrow \pi_1(M^3)$$

injective



A skew pair of pants



$\gamma_i \equiv f(c_i)$ is a closed geodesic

$\gamma_i \equiv f(h_i)$ is a geodesic segment
orthogonal to $f(\gamma_{i \pm 1})$.

The feet of a skew pair of pants




$$N^1(\gamma_i) \cong \mathbb{C}/2\pi i\mathbb{Z} + h\ell(\gamma)\mathbb{Z}$$



Two and Three Dimensions


2D



$$N^1(\gamma) \cong \mathbb{R} / \ell(\gamma) \cdot \mathbb{Z} \times \{-1, +1\}$$

$$\cong \mathbb{R}^* / \langle x \mapsto e^{\ell(\gamma)} x \rangle$$

3D




$$N^1(\gamma) \cong \mathbb{C} / \ell(\gamma) \cdot \mathbb{Z} + 2\pi i \mathbb{Z}$$

$$\cong \mathbb{C}^* / \langle z \mapsto e^{\ell(\gamma)} z \rangle$$


The meaning of equidistribution

The set $\{x \in \mathbb{R}^d : x \in \mathcal{P}, P \in \Pi_{\varepsilon, R}\}$

are $e^{-\varepsilon R}$ - evenly distributed
as points on $N^1(\sqrt{\varepsilon})$



$$x \mapsto x + (\pi + 1)$$



The "doubling trick" revisited

We take


$$A_\gamma = \{ \text{foot}_\gamma P : \gamma \in \partial P, P \in \Pi_{\varepsilon, R} \}$$

We can find

$\sigma: A_\gamma \rightarrow A_\gamma$ a permutation

such that, $|\sigma(x) - x - i\pi - 1| < \varepsilon/R$
(for all $x \in A_\gamma$)

and then define

$$\mathcal{I}: A_\gamma^+ \sqcup A_\gamma^- \rightarrow A_\gamma^+ \sqcup A_\gamma^-$$


The Idea of Self-Correction

We find $g: \mathbb{Q} \Gamma_{\varepsilon, R} \rightarrow \mathbb{Q} \Pi_{\varepsilon, R}$
such that

1. $\partial g(\alpha) = \alpha$ whenever $\alpha = \partial \pi$

2. $\|g(\alpha)\|_{\infty} \leq P(R) e^{-R} \|\alpha\|_{\infty}$ for all α .

(where $P(R) = C R^N$ is
a polynomial in R)

Counting Good Pants and Good Curves

$$\# \pi_{\varepsilon, R} \asymp e^{3R}$$

$$\# \Gamma_{\varepsilon, R} \asymp e^{2R/R}$$

$$\# \{P \in \pi_{\varepsilon, R} \mid \gamma \in \partial P\} \asymp e^{R/R}$$

for each $\gamma \in \Gamma_{\varepsilon, R}$

Then let $\pi = \sum_{P \in \Pi_{\varepsilon, R}} P$, and $\alpha = \partial \pi$.


Then $\|\alpha\|_{\infty} \leq e^{(1-\delta)R}$ by equidistribution

So $\|g(\alpha)\|_{\infty} \leq P(R)e^{-\delta R} \ll 1$.

So $\pi - g(\alpha)$ is positive and equi-distributed

And $\partial(\pi - g(\alpha)) = \partial \pi - \partial g(\partial \pi) = \partial \pi - \partial \pi = 0$.

After clearing denominators, we can assemble the pants of $\pi - g(\alpha)$ to form a good cover!



The Relative Surface Subgroup Theorem

Let M be a closed hyperbolic ~~3~~
3-manifold, and let γ be a closed
geodesic in M of length $l(\gamma) \geq R_0(M, \varepsilon)$.
(and suppose γ bounds a singular chain).

Then we can find a compact
hyperbolic surface S with boundary
and $f: S \rightarrow M$ that is ε -close to
being an isometric immersion, and
for which $f(\gamma) = \gamma$ for every component
 γ of ∂S .

The Good Pants Homology

We let $H_1^{\varepsilon, R}(S; \mathbb{Q}) = \mathbb{Q}\Gamma_{\varepsilon, R} / 2\mathbb{Q}\Pi_{\varepsilon, R}.$

We find a series of identities
leading to $H_1^{\varepsilon, R}(S; \mathbb{Q}) = H_1(S; \mathbb{Q}).$

The Good Pants Homology

We let $H_1^{\varepsilon, R}(S; \mathbb{Q}) = \mathbb{Q}\Gamma_{\varepsilon, R} / \partial \mathbb{Q}\Pi_{\varepsilon, R}.$

We find a series of identities
leading to $H_1^{\varepsilon, R}(S; \mathbb{Q}) = H_1(S; \mathbb{Q}).$

Then we observe that $g: \mathbb{Q}\Gamma_{\varepsilon, R} \rightarrow \mathbb{Q}\Pi_{\varepsilon, R}$
has been implicitly defined, such
that $\partial g(\alpha) = \alpha$ when $\alpha = 0$ in $H_1.$

The Algebraic Square Lemma

Under conditions of reasonable geometry,

$$\sum_{i,j=0,1} (-1)^{i+j} [A_i \cup B_j V] \equiv 0$$

in $H_1^{\varepsilon, R}$

(where $[X]$ denotes the closed
geodesic ^{freely} homotopic to X , for
 $X \in \pi_1(S, x)$).

We then define

$$A_T = \frac{1}{2}([TA\bar{T}B] - [T\bar{A}TB])$$

(which is independent of the choice
of B in $H_1^{\varepsilon, R}$)

and prove

$$(XY)_T = X_T + Y_T$$

in $H_1^{\varepsilon, R}$.